

# Appendices

**Appendix 1 – Glossary of terms used**

**Appendix 2 – Simulation modelling for the four risk assessment examples**

**Appendix 3 – Predictive microbiology: concepts, applications and sources**

**Appendix 4 – Prevalence and incidence of *Listeria monocytogenes* in fermented meat products**

**Appendix 5 – Background for the cold-smoked fish risk assessment**



# Appendix 1.

## Glossary of Terms

### Beta distribution

The Beta distribution is defined as

$$f(x) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1}(1-x)^{\beta-1},$$

where  $0 \leq x \leq 1$ ,  $\alpha > 0$  and  $\beta > 0$ . There are generalizations to a random variable defined on any interval [a, b].

(Source: <http://www.statsoft.com/textbook/glosfra.html>)

### Binomial distribution

The binomial distribution is used when each trial has exactly two mutually exclusive possible outcomes, often labelled success and failure. The binomial distribution is the probability of obtaining x successes in N trials where the probability of success on a single trial is  $\pi$ . The binomial distribution assumes that  $\pi$  is fixed for all trials. The formula for the binomial probability mass function is

$$p(x; n, \pi) = \binom{n}{x} \pi^x (1-\pi)^{n-x}, \text{ for } x = 0, 1, 2, \dots, n > 0.$$

(Source: [www.itl.nist.gov/div898/handbook/eda/section3/eda366h.htm](http://www.itl.nist.gov/div898/handbook/eda/section3/eda366h.htm))

### Confidence interval

A range of values believed to include an unknown population parameter. Associated with the interval is a measure of the confidence we have that the interval contains the parameter of interest, the *confidence level*, (depending on interpretation) the probability that the parameter of interest will fall within the specified confidence interval. Where a point estimate is a specific numerical value that estimates a parameter, an interval estimate such as a confidence interval is a numeric range that estimates a parameter, generally with an associated probability. A confidence interval for a parameter generalizes to a *confidence set* for more than one parameter at a time. (Source: [www2.spsu.edu/tmgt/richardson/Statistics/](http://www2.spsu.edu/tmgt/richardson/Statistics/))

It should be noted that, under the frequentist definition, the confidence level is the probability of the interval covering the true unknown value. The true value is fixed and it is the interval that is random in repeated experimentation.

### Continuous random variable

A continuous random variable is one that takes an infinite number of possible values. Continuous random variables are usually measurements. Examples include height, weight,

and the concentration of *L. monocytogenes* in a sample. Examples of probability distributions for continuous random variables are the Normal distribution and the Gamma distribution.

(Source: [www.stats.gla.ac.uk/steps/glossary/](http://www.stats.gla.ac.uk/steps/glossary/))

### Correlation

Correlation is a measure of the relation between two or more variables. Correlation coefficients can range from -1 to +1. The value of -1 represents a perfect negative correlation while a value of +1 represents a perfect positive correlation. A value of 0 represents a lack of correlation.

(Source: [www.statsoft.com/textbook/stathome.html](http://www.statsoft.com/textbook/stathome.html))

### Convolution

Consider  $X$  and  $Y$  are non-negative, independent, integer-valued random variables with probability distributions  $\Pr\{X=j\}=a_j$  and  $\Pr\{Y=j\}=b_j$  and  $\Pr\{X=y, Y=k\}=a_y b_k$ . The sum  $S=X+Y$  is a random variable also, and we recognize that the event  $S=r$  is the union of events  $(X=0, Y=r), (X=1, Y=r-1), \dots, (X=r, Y=0)$ . So, the probability distribution for  $S$  is  $\Pr\{S=r\}=c_r$  where  $c_r=a_0 b_r + a_1 b_{r-1} + \dots + a_r b_0 = \sum_{k=0}^r a_k b_{r-k}$ . Feller (1968: 266 et ff.) names this operation convolution (German *Faltung*, French *composition*) and extends the definition to any 2 sequences  $\{a_k\}$  and  $\{b_k\}$ , not necessarily probability distributions. Combinations like this appear in much of the simulation.

### Deterministic

Commonly, *deterministic* is an antonym for *stochastic*.

### Discrete random variable

A discrete random variable is one which may take on only a countable number of distinct values such as 0, 1, 2, 3, 4, ... Discrete random variables are usually, but not necessarily, counts. If a random variable can take only a finite number of distinct values, then it must be discrete. Examples of discrete random variables include the number of children in a family, the Friday night attendance at a cinema, the number of *L. monocytogenes* organisms in a serving of food. Examples of probability distributions for discrete random variables are the Binomial distribution and the Poisson distribution.

(Source: [www.stats.gla.ac.uk/steps/glossary/](http://www.stats.gla.ac.uk/steps/glossary/))

### Distribution function

The distribution of a variable is a description of the relative numbers of times each possible outcome in the domain of the variable will occur in a number of trials. The function describing the distribution is called the probability function (probability mass function if the random variable takes only discrete values; probability density function if the random variable is continuous). The *cumulative distribution function* describes the probability that a trial takes on a value less than or equal to a number, commonly  $F(x) = \Pr\{X \leq x\}$ . The cumulative distribution function is monotone increasing whereas the probability density is not.

(Source: [mathworld.wolfram.com](http://mathworld.wolfram.com))

### Empirical distribution function

Given data  $\{x_k, k = 1, \dots, n\}$  sorted from smallest to largest,  $\{x_{(k)}, k = 1, \dots, n\}$ ,  $x_{(1)} \leq x_{(2)} \leq \dots \leq x_{(n)}$ , the empirical (cumulative) distribution function (e.c.d.f. or e.d.f.) is the function

defined by  $\hat{F}(x) = \frac{\text{number of } x_k \leq x}{n}$ , a step function with steps of size  $1/n$ . The values

of the e.c.d.f. are the discrete set of cumulative probabilities  $\{0, 1/n, \dots, n/n\}$ . When used in a simulation, values between any two consecutive samples,  $x_{(k)}$  and  $x_{(k+1)}$  cannot be simulated, nor can a value smaller than the minimum, nor can a value larger than the maximum. The e.c.d.f. has mean equal to the sample mean, and variance equal to  $(n-1)/n$  times the sample variance. The e.c.d.f. tends to underestimate the true mean and variance when the underlying distribution is skewed to the right. Expected values of simulated e.c.d.f. quantiles are equal to the sample quantiles. Some variations on the e.c.d.f. appear in simulations: linearly extrapolating between observations; or adding lower and upper tails to the data to reflect a range of the variable outside the observed range, either through expert judgement or by postulating some shape to the tails beyond the sample extremes.

### Gamma distribution

The probability density of the Gamma distribution is defined as

$$f(x) = x^{\alpha-1} e^{-x/\beta} [\beta^\alpha \Gamma(\alpha)]^{-1},$$

where  $x \geq 0$ ,  $\alpha > 0$ ,  $\beta > 0$ .  $\alpha$  is referred to as the shape parameter.  $\beta$  is referred to as the scale parameter. For integral  $\alpha$ , one can recognize the Gamma distribution as the distribution of the waiting time for  $\alpha$  Poisson events. As a special case, when  $\alpha = 1$ , the Gamma distribution is the Exponential distribution.

(Source: [www.statsoft.com/textbook/glosfra.html](http://www.statsoft.com/textbook/glosfra.html))

### Latin Hypercube Sampling

Latin Hypercube Sampling (LHS) is a stratified sampling technique where the random variable distributions are divided into equal probability intervals. A probability is randomly selected from within each interval for each basic event. Generally, LHS will require fewer samples than simple Monte Carlo sampling for similar accuracy. LHS ensures that the entire range of each variable is sampled.

(Source: [http://saphire.inel.gov/guest\\_area/SAF00758.htm](http://saphire.inel.gov/guest_area/SAF00758.htm))

### Lognormal distribution

The lognormal distribution has the probability density function

$$f(x) = \frac{1}{[x\sigma\sqrt{2\pi}]} \exp\left(-\frac{1}{2}[\ln x - \mu]^2 / \sigma^2\right),$$

where  $0 \leq x < \infty$ ,  $\mu > 0$ ,  $\sigma > 0$ . If the distribution of a random variable  $X$  is lognormal, then the distribution of  $\ln(X)$  is Normal.

(Source: [www.statsoft.com/textbook/glosfra.html](http://www.statsoft.com/textbook/glosfra.html))

### Maximum likelihood

The method of maximum likelihood is a general method of estimating parameters of a population by values that maximize the *likelihood* ( $L$ ) of a sample. The likelihood  $L$  of a sample of  $n$  observations  $x_1, x_2, \dots, x_n$ , is the joint probability function  $p(x_1, x_2, \dots, x_n)$  when  $x_1, x_2, \dots, x_n$  are discrete random variables. If  $x_1, x_2, \dots, x_n$  are continuous random variables, then the likelihood  $L$  of a sample of  $n$  observations,  $x_1, x_2, \dots, x_n$ , is the joint density function  $f(x_1, x_2, \dots, x_n)$ . When  $L$  is a function of parameters, then the maximum likelihood estimates (m.l.e.) of the parameters are the values that maximize  $L$ .

(Source: [www.statsoft.com/textbook/stathome.html](http://www.statsoft.com/textbook/stathome.html))

### Method of moments

This method can be employed to determine parameter estimates for a distribution. The method of matching moments sets the distribution moments equal to the data moments and solves to obtain estimates for the distribution parameters. For example, for a distribution with two parameters, the first two moments of the distribution (the mean  $\mu$  and variance  $\sigma^2$  of the distribution) would be set equal to the first two moments of the data (the sample mean and variance, e.g. the unbiased estimators  $\bar{X}$  and  $s^2$ ) and solved for the parameter estimates.

(Source: [www.statsoft.com/textbook/glosfra.html](http://www.statsoft.com/textbook/glosfra.html))

### Monte Carlo

In Monte Carlo methods, the computer uses random number simulation techniques to mimic a statistical population. For each Monte Carlo replication, the computer: simulates a random sample from the population; analyses the sample; and stores the result. After many replications, the stored results will mimic the sampling distribution of the statistic.

(Source: [www.statsoft.com/textbook/stathome.html](http://www.statsoft.com/textbook/stathome.html)).

### Normal distribution

A continuous random variable  $X$  has a Normal distribution if its probability density function is  $f(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)$ ,  $-\infty < x < \infty$ ,  $\sigma > 0$ ,  $-\infty < \mu < \infty$ . The normal probability density function has two parameters:  $\mu$  (mean) and  $\sigma$  (standard deviation). The Normal distribution is sometimes called the *Gaussian* distribution.

(Source: [http://ce597n.www.ecn.purdue.edu/CE597N/1997F/students/michael.a.kropinski.1/project/tutorial#Normal Distribution](http://ce597n.www.ecn.purdue.edu/CE597N/1997F/students/michael.a.kropinski.1/project/tutorial#Normal%20Distribution))

### Quantile

The  $p^{\text{th}}$  quantile of a distribution of values is a number  $x_p$  such that a proportion  $p$  of the population values are less than or equal to  $x_p$ . In a simple random sample of  $n$  values, where the sample values ordered in ascending order are  $x_{(1)}, \dots, x_{(n)}$ , it is common to use the  $x_{(k)}$  as an estimate of the  $k/(n+1)^{\text{th}}$  quantile, although different software packages use variations of this,  $(k-\alpha)(n-\alpha-\beta)^{-1}$  for  $\alpha, \beta > 0$  (Hyndman and Fan, 1996).

(Source: [www.statsoft.com/textbook/glosfra.html](http://www.statsoft.com/textbook/glosfra.html))

## Quantitative risk assessment

If “risk assessment is generally regarded as a process to scientifically evaluate the probability and severity of known or potential adverse effects attributable to a hazardous agent, process or circumstance” (Cassin et al., 1998), then quantitative risk assessment “implies an estimation of the probability and impact of adverse health outcomes...” (Cassin et al., 1998).

## Poisson distribution

The Poisson distribution is defined as  $\Pr\{X=k\} = \mu^k e^{-\mu} / k!$ ,  $x = 0, 1, \dots$ , where  $\mu > 0$  is the average number of occurrences (count) per interval. A Poisson random variable  $X$  is a count, interpreted in the context of either distance, area, volume, time or other measure of size (interval) as follows:

- Each non-overlapping interval increment of interest is so small that only one event can occur within it (or at least, the probability of 2 or more events in the interval is negligible), but the sum of the individual increments comprises the entire interval or time period; and
- the probability of an event occurring in the given increment is constant. The number of events observed depends only on the length of the interval considered and not on its end points. If length of interval is 0 and time is 0, the number of events observed is 0. The numbers of changes in non-overlapping intervals are independent for all intervals.

Examples occur in many fields: the number of imperfections (gas trap or cracks) per square metre in rolls of metals; the number of telephone calls per hour received by an office; the number of cashews per can in one can of mixed nuts; the number of bacteria in a given culture; or the number of typing errors per page. The specified region can be an area, a volume, a segment of a line or even a piece of material.

(Sources: <http://engineering.uow.edu.au/Courses/Stats/File40.html>  
<http://mathworld.wolfram.com/PoissonDistribution.html>)

## Rank Correlation

A rank correlation coefficient is a correlation coefficient that is based on the ranks of the sample values and not the actual values. A rank is a consecutive number assigned to a specific observation in a sample of observations sorted by their values. So, ranks reflect the ordered relation of one observation to the others in the sample. The lowest value is assigned a rank of 1; the higher ranks represent the higher values.

(Source: [www.statsoft.com/textbook/stathome.html](http://www.statsoft.com/textbook/stathome.html))

## Simulation

Etymology: Middle English *simulation*, from Middle French, from Latin *simulation-*, *simulatio*, from *simulare*

1. the act or process of simulating.
2. a sham object.
- 3a. the imitative representation of the functioning of one system or process by means of the functioning of another *<a computer simulation of an industrial process>*.
- 3b. examination of a problem often not subject to direct experimentation by means of a simulating device.

See also: Monte Carlo. (Source: Merriam-Webster Collegiate Dictionary On-line.  
www.m-w.com/cgi-bin/mweb)

## Stochastic

Etymology: Greek *stochastikos* skilful in aiming, from *stochazesthai* to aim at, guess at, from *stochos* target, aim, guess.

- RANDOM; specifically: involving a random variable <a stochastic process>
- involving chance or probability: PROBABILISTIC <a stochastic model of radiation-induced mutation>.

(Source: Merriam-Webster Collegiate Dictionary On-line.  
<http://www.m-w.com/cgi-bin/mweb>)

A stochastic process is a family of random variables  $X(t)$  indexed by a parameter  $t$ , which usually takes values in the discrete set  $T = \{0, 1, 2, \dots\}$  or the continuous set  $T = [0, +\infty)$ . In many cases  $t$  represents time, and  $X(t)$  is a random variable observed at time  $t$ . Examples are the Poisson process, the Brownian motion process, and the Ornstein-Uhlenbeck process. Considered as a totality, the family of random variables  $\{X(t), t \in T\}$  constitutes a "random function".

(Source: [www.britannica.com/bcom/eb/article/3/0,5716,117323+26+109439,00.html](http://www.britannica.com/bcom/eb/article/3/0,5716,117323+26+109439,00.html))

Commonly, *deterministic* is an antonym for *stochastic*.

NOTE: A more extensive glossary of terms related to microbiological risk assessment can be found in MRA 3, an earlier volume in this series (FAO/WHO, 2003).

## REFERENCES CITED IN THE GLOSSARY OF TERMS

- Feller, W. 1968. *An Introduction to Probability Theory and its Applications*. 2nd edition. New York NY: Wiley.
- Hyndman, R.J. & Fan, Y. 1996. Sample quantiles in statistical packages. *American Statistician*, **50**: 361–365.
- Cassin, M.H., Lammerding, A.M., Todd, E.C.D., Ross, W. & McColl, R.S. 1998. Quantitative risk assessment for *Escherichia coli* O157:H7 in ground beef hamburgers. *International Journal of Food Microbiology*, **41**: 21–44.
- FAO/WHO. 2003. Hazard characterization for pathogens in food and water: Guidelines. *FAO/WHO Microbiological Risk Assessment Series*, No. 3. 61p.

# Appendix 2.

## Simulation modelling for the four risk assessment examples

### A2.1 INTRODUCTION

This appendix serves as documentation for the simulation modelling carried out for the pasteurized milk and ice cream examples, making the work transparent. The model documentation reflects methodological issues, but is not intended to explain the issues in detail. Specific issues related to the development of the two risk assessment models are addressed here. These include a description of the consumption characteristics used and the modelling of home storage conditions, and how non-susceptible and susceptible populations might be defined. How to combine independent data sets to describe prevalence of *L. monocytogenes* in foods and to describe concentration of *L. monocytogenes* in foods are also addressed. The appendix provides a list of references to support the documentation here, and to provide that vast amount of supplementary material that is the background for much of the work. Still other methodological material appears in the main body of the report (Part 3 – Exposure assessment, and Part 4 – Example risk assessments). Implementation of the Monte Carlo simulation for this exposure assessment was performed using Analytica™1.11, 2.0.1 or 2.0.5 (Lumina Decisions) software. Additional computations and preparation of graphs were done using Microsoft® Excel 97 and Microsoft® Excel 2000, and with S-Plus 4.5 Professional, S-Plus 2000 Professional and S-Plus 6 Professional (MathSoft, Inc.).


### A2.2 MODELLING THE EXPOSURE ASSESSMENT

#### A2.2.1 Overview

Objectives for the exposure assessment are to simulate the number of *L. monocytogenes* organisms in a serving of a particular RTE) food, *Lm ingested*, and to determine the annual frequency of servings for individuals in the consuming population, *Annual meals*. Every shape in the influence diagram (Figure A2.1) is termed a node. Different shaped nodes perform different functions. The hexagonal figures, *Lm ingested* and *Annual meals*, represent the stochastic results that answer the questions deriving from the objectives. Elliptical shapes, such as *Food amount eaten*, are chance (stochastic) nodes that hold intermediate calculations that form part of the modelling for the objective nodes. The round-cornered rectangular nodes, such as *Prevalence and concentration*, are organizing modules that contain other nodes. The hexagonal pennant boxes, *Discrete distributions* and *Study indices* are libraries of functions that support some calculations or that contain index nodes that structure the results. Arrows indicate influences and indicate the direction of the influence. For example, the number of *L. monocytogenes* organisms ingested in a serving when that number exceeds zero, *Lm ingested given >0*, depends on the *Concentration in ingested food* and the *Food amount eaten*. The values in *Food frequency* determine what values reside in *Annual meals*.

To model the exposure assessment, the following information is needed:

1. *Prevalence and concentration characteristics*, measured at the same consistent point in the farm-to-fork chain. Prevalence relates how often the food is contaminated with *L. monocytogenes*. The notion is generalized to consider it equivalent to the probability that a serving from a package or unit of product contains any contamination. Concentration defines how many *L. monocytogenes* organisms are in a contaminated portion.
2. *Storage characteristics* and *Growth characteristics* that determine the amount of *Growth* of *L. monocytogenes* in the product from that point in the process to the point of consumption.
3. *Consumption characteristics* that relate how much food consumers eat and how often they eat it. How large a serving the consumer eats determines how many *L. monocytogenes* organisms the consumer ingests.
4. *Non-susceptible and susceptible populations*. Hazard Identification generally indicates that some portions of the consuming population are more susceptible to infection or illness from *L. monocytogenes*.



**Figure A2.1** Influence diagram for *Listeria monocytogenes* exposure assessment.

Figure A2.1 clarifies the model structure and what information is needed from the exposure assessment. However, it does not make explicit the node's specific parameterization. Also, it does not show all of the interrelationships and dependencies. Accompanying documentation in this Appendix does that (such as Table A2.1), and specific methodological issues are addressed in the various sections.

**Table A2.1** Nodes for top level of *Listeria monocytogenes* in RTE eat foods exposure assessment model.


Title Identifier Structure	Description and Definition
Non-susceptible and susceptible populations Nonandsusceptible Module	Module holds characteristics that define the allocation of individuals from Gender $\times$ Age groups to non-susceptible and susceptible groups. Among adults, for whom we have some information about consumption characteristics, susceptible groups are defined to include all adults 65 and older, pregnant women (1.3% of the population) and individuals with suppressed immune systems and certain medical conditions such as cancer and recent organ transplantation (3.3% of the population) (Miller, Whiting and Smith, 1997).
Consumption characteristics Consumption_character Module	Consumption characteristics come from 24-hour recall data from CFPNS (1992–1995), which addressed the nutritional habits of non-institutionalized adults between 18 and 74 years old in Québec, Nova Scotia, Saskatchewan, Alberta and Prince Edward Island.
Prevalence and concentration Prevalence_and_conce Module	Prevalence and concentration module determines simulated distributions for the prevalence of <i>L. monocytogenes</i> in food and the concentration in contaminated food, nominally at retail.
Storage characteristics Storage_characterist Module	Storage characteristics module determines simulated distributions for the refrigerator temperature that the consumer stores the food at, and the length of time, measured as if from retail purchase, to the time of consumption.
Growth Growth2 Module	Growth module determines growth characteristics for <i>L. monocytogenes</i> in the food. Growth is characterized by exponential growth rates and stationary phase population size for each foodstuff.
Food amount eaten Serving_size1 (g) Chance node	Food amount eaten is the simulated distribution of the daily serving size for individuals from the non-susceptible group and for individuals from the susceptible group. Food amount eaten comes directly from the consumption amounts generated in the Consumption characteristics module.
Annual meals Annual_meals Objective node	The number of Annual meals for an individual is calculated from the Food frequency probability of consumption on a given day, by implementing Binomial sampling. The number of meals (population days with consumption) is calculated in the Consumption characteristics module, rounded here for display $\text{Table}(\text{Annualmealsreporting})(\text{Round}(\text{Binomial}(365, \text{Mealfrequency}[\text{Annualmealsreporting}=\text{'Individual'}])), \text{Round}(\text{Mealfrequency}[\text{Annualmealsreporting}=\text{'Population'}]))$
Prevalence in ingested food Prevalence_in_ingest Chance node	The Prevalence in ingested food node is the simulated distribution for how often a serving contains any <i>L. monocytogenes</i> contamination. $\text{For Icebox}=\text{Refrigerator\_studies Do For Person}=\text{Risk\_group\_definitio Do Correlatedprevalence} * (1-\text{Exp}(-10^{\wedge}\text{Finalconcentration}[\text{Refrigerator\_studies}=\text{Icebox}] * \text{Serving\_size1}[\text{Risk\_group\_definitio}=\text{Person}] ) )$
Contaminated serving or not There_or_not_there Chance node	Contaminated serving or not is a simple accounting of whether a serving is contaminated or not. It is generated by sampling from the outcomes Not contaminated and Contaminated, with probabilities Not contaminated $(1-\text{Prevalence\_in\_ingest})$ Contaminated $\text{Prevalence\_in\_ingest}$

Title Identifier Structure	Description and Definition
Concentration in ingested food Finalconcentration (log <sub>10</sub> CFU/g) Chance node	Concentration in ingested food is the simulated distribution of the concentration of <i>L. monocytogenes</i> in contaminated food at ingestion. Initial concentrations grow into final concentrations according to the growth determined in the Growth module. Final concentrations are restricted to theoretical maximum population densities or stationary phase densities. Using Calculatedfinal:= Initialconcentration + Unconstrgrowthamount Do ( if Maximum_population >0 then ( if Calculatedfinal <= logten(Maximum_population) then Calculatedfinal else logten(Maximum_population) ) else Calculatedfinal )
Lm ingested given >0 Dose (CFU) Chance node	The Lm ingested given >0 node records the simulated distribution of the number of <i>L. monocytogenes</i> organisms in a serving of food, in those cases where the number of organisms is larger than 0. The Prevalence in ingested food node lets us derive how often the number of organisms is 0. Lm ingested given >0 generates non-zero observations by sampling on [1,∞) with Poisson probabilities. For Icebox:=Refrigerator_studies Do For Person:=Risk_group_definitio Do logten(Round(Conditional_poisson(10^Finalconcentration[Refrigerator_studies=Icebox x] * Serving_size1[Risk_group_definitio=Person])))
Lm ingested Lm_ingested (CFU) Objective node	Lm ingested is one of the objective nodes for this exposure assessment. It is calculated by combining the simulated distributions for the Prevalence in ingested food and the Lm ingested given >0. The number of <i>L. monocytogenes</i> organisms ingested when the food is not contaminated is assumed to be 0. For temp:=Run Do if There_ot_not_there[Run=temp]='Not contaminated' then 0 else 10^Dose
Study indices Row_and_column_inde1 Library	Study indices is a collection of index information that structure the results.

### A2.2.2 Non-susceptible and susceptible populations

The susceptible population is determined by the fractions of persons who have one of the characteristics named: elderly, pregnant, otherwise susceptible, young. For the present implementation of the exposure assessment, where there are no consumption data for persons under 18 years old from Canadian data (CFPNS, 1992–1995), the fraction young is moot. In Figure A2.2, *Population age and gender* table holds domain estimates for the

Age and Gender groups used. For Canadian consumption data, estimates represent the population counts, in the years of the surveys, in five provinces: Alberta, Nova Scotia, Prince Edward Island, Quebec and Saskatchewan, for which consumption information is available (CFPNS, 1992–1995). *Fraction elderly*, *Fraction pregnant*, *Fraction otherwise susceptible*



**Figure A2.2** Influence diagram for non-susceptible and susceptible populations.

and *Fraction young* are tables that describe the fraction of the population, in Gender  $\times$  Age groups, who would be attributed to the susceptible group, for the named reason. For example, a fraction of Females, 18–34 and 35–49 would be attributed to the susceptible group in the *Fraction pregnant* table. All persons 65 and older, but no others, would be attributed to the susceptible group in the *Fraction elderly* table. Allocations were based on Miller, Whiting and Smith (1997), as applied to the Canadian population. The *Population susceptible group* table collects the fractions together to give the population size in the susceptible group, by Gender and by Age. The *Population non-susceptible group* table is derived by difference from the *Population age and gender* and *Population susceptible group* tables to give the population size in the non-susceptible group, by Gender and by Age. *Risk group characterization* is a useful summary of the attribution of individuals to the non-susceptible and susceptible groups. This module is also a natural holding place for parallelogram-shaped index nodes, *Risk groups*, *Gender* and *Age group*, that structure the results through many modules in the model. Details of the nodes are described in Table A2.2.

**Table A2.2** Nodes for non-susceptible and susceptible populations module.


Title, Identifier, Structure	Description and Definition
Population Age and gender Popn_age_gender Module	Population age and gender table holds domain estimates for the Age and Gender groups used.
Fraction elderly Fraction_elderly Variable node	
Fraction Pregnant Fraction_pregnant Variable node	Fraction elderly, Fraction Pregnant, Fraction otherwise susceptible and Fraction young are tables that describe the fraction of the population, in Gender $\times$ Age groups, whom we would attribute to the susceptible group, for the named reason.
Fraction otherwise susceptible Fraction_otherwise_s Variable node	
Fraction young Fraction_young Variable node	
Population non-susceptible risk group Population_normal Variable node	Population non-susceptible group table is derived by difference from the Population Age and gender and Population Susceptible group tables to give the population size in the non-susceptible group, by Gender and by Age. Popn_age_gender — Population_high_risk
Population susceptible group Population_high_risk Variable node	Population susceptible group table collects the fractions together to give the population size in the susceptible group, by Gender and by Age. Popn_age_gender * ( Fraction_elderly + Fraction_otherwise_s + Fraction_pregnant + Fraction_young )
Risk group characterization Riskgroupcharacteriz Variable node	Risk group characterization is a useful summary of the attribution of individuals to the risk groups.
Risk groups Risk_group_definitio Index node	Risk groups are defined as ['Non-susceptible', 'Susceptible']
Gender Gender_definition Index node	Gender is defined as ['Female', 'Male']
Age group Age_group_definition Index node	Age group is defined as ['18-34', '35-49', '50-64', '65-74']

### A2.2.3 Consumption characteristics

Consumption characteristics are the amount of food eaten in a serving, the daily probability of consuming the food and the annual number of meals (days with consumption) in the population. The nodes *Food amount eaten* and *Food frequency* are collectors for the characteristics calculated in each of the food-specific modules: *Ice cream*; and *Fluid milk, pasteurized*. *Food amount eaten* is the distribution of amounts eaten (g). *Food frequency* is the probability of consuming the food on a given day. *Survey results* and *Ecdf columns* are index nodes that structure data tables in the *Fluid milk, pasteurized* and *Ice cream* modules (Table A2.3, Figure A2.3). Exposure assessment examples for pasteurized milk and ice cream were implemented using the modules described here.

**Table A2.3** Nodes for Consumption characteristics module.

Title, Identifier, Structure	Description and Definition
Ice cream Consumption_char1 Module	The main consumption characteristics are the serving size and the frequency of consumption for ice cream.
Fluid milk, pasteurized Consumption_char3 Module	The main consumption characteristics are the serving size and the frequency of consumption for pasteurized fluid milk.
Food amount eaten Serving_size (g) Chance node	Food amount eaten is the distribution of amounts eaten (g). DetermTable(Food_groups)(Samp_cons_ice, Samp_cons_pmilk)
Food frequency Meal frequency Chance node	Food frequency is the probability of consuming the food on a given day and the annual meals (population consumption days) for population. Determtable(Food_groups, Annualmealsreporting)(Samp_freq_ice, Annualmealsicecream, Samp_freq_pmilk, Annualmealspmilk)
Survey results Survey_results Index node	Survey results structures data tables in Fluid milk, pasteurized and Ice cream modules. ['Respondents', 'Consumers']
Ecdf columns Ecdf_columns Index node	Ecdf columns structures data tables in Fluid milk, pasteurized and Ice cream modules. ['Amount', 'Fraction']



**Figure A2.3** Influence diagram for *Consumption characteristics* module.


### A2.2.4 Ice cream and Fluid milk, pasteurized modules

Consumption characteristics modules are specific to the RTE food considered. The structure of consumption characteristics modules for *Ice cream* and for *Fluid milk, pasteurized* are identical. Both are described together, referring to the *Fluid milk, pasteurized* example (Table A2.4, Figure A2.4).

**Table A2.4** Nodes for *Ice cream* and for *Fluid milk, pasteurized* modules.

Title, Identifier, Structure	Description and Definition
Gender, age consumption frequency Gender_age_freq6 Variable node	Gender, age consumption frequency is a table indexed by Age group and Gender, holding point estimates of daily consumption probabilities for Pasteurized milk. Other nodes in the module use the point estimates to determine what proportions of Age and Gender group characteristics to include in non-susceptible and susceptible populations.
Susceptible group proportions Gender_age_pro_high6 Variable node	Susceptible group proportions is a table that holds proportions of total eating episodes assigned to Gender and Age group for individuals in the susceptible group. We have adjusted Gender, age consumption frequency proportions to reflect membership in susceptible groups, Population: susceptible group.
Susceptible group gender Sampled_high_risk_g6 Chance node	Susceptible group gender and Susceptible group age are stochastic nodes that hold a Gender and an Age Group, sampled so that Gender $\times$ Age group proportions among consumers in the susceptible population are respected.
Susceptible group age Sampled_high_risk_a6 Chance node	
Non-susceptible group proportions Normal_intake_pr6 Variable node	Non-susceptible group proportions is a table that holds proportions of total eating episodes assigned to Gender and Age group for individuals in the non-susceptible group. We have adjusted Gender, age consumption frequency proportions to reflect membership in non-susceptible groups, Population: Non-susceptible group.
Non-susceptible group gender Sampled_gender6 Chance node	Non-susceptible group gender and Non-susceptible group age are stochastic nodes that hold a Gender and an Age Group, sampled so that Gender $\times$ Age group proportions among consumers in the non-susceptible population are respected.
Non-susceptible group age Sampled_age_group6 Chance node	
Nutrition survey results, milks Nutrition_survey_re6 Variable node	Nutrition survey results, milks is a table that holds inferential statistics from the nutrition surveys: the number of survey respondents and the number who reported consuming Pasteurized milk on a given day.
Pasteurized milk amounts Pmilk_amount Variable node	Pasteurized milk amounts is a table that holds empirical daily Pasteurized milk amounts collected from the nutrition surveys. The table has columns Amount, an amount consumed (g) and Fraction, the inverse of the design-based weights associated with Pasteurized milk consumers in the nutrition surveys. There is a separate table for each Gender $\times$ Age group.
Milk amount index Pmilk_amount_index Index node	Milk amount index structures the table of amounts, Pasteurized milk amounts. Range of sequence corresponds to a set of rows in an Excel spreadsheet. Sequence(5, 459, 1)
Beta, milks frequency Beta_frequency6 Chance node	Beta, milks frequency uses a Beta distribution to represent uncertainty or variability over a Gender $\times$ Age group, for consumers' consumption probability on a given day. Beta, milks frequency is assumed to be Beta( $x+1$ , $n-x+1$ ), where $n$ is the number of respondents, and $x$ is the estimated number of Pasteurized milk consumers, $n\pi$ .


Title, Identifier, Structure	Description and Definition
Annual meals, susceptible group Annualmealshighrisk6 Chance node	Annual meals, susceptible group simulates the annual meals (population days with consumption) in susceptible population. It incorporates the variability and uncertainty about the fraction of adults who consume Pasteurized milk on a given day. It samples binomially among population days (population × 365), but uses a Normal approximation. We truncate the Normal distribution at 0 and the total population days.  Sum(Sum(Using Beta_value:= Beta_frequency6[Risk_group_definitio='High risk'] Do Using People_days:=Population_high_risk*365 Do Using Interim:= -Truncate( -(Truncate(Normal(People_days*Beta_value, People_days*sqrt( Beta_value*(1-Beta_value))), 0)), -People_days) Do if Interim<=0 then 0 else if Interim>=People_days then People_days else Round(Interim), Gender_definition), Age_group_definition)



**Figure A2.4** Influence diagram for *Pasteurized milk* consumption characteristics module. Except for changes to node identifiers, the *Ice cream* consumption characteristics module is identical.

### A2.2.5 Prevalence and concentration

The *Prevalence and concentration* module simulates the *Prevalence characteristics* – nominally prevalence of *L. monocytogenes* contamination in foods at retail or source for the consumer – and simulates the *Concentration characteristics* – nominally the *L. monocytogenes* concentration in the food at that point. The module lets one specify a rank correlation coefficient between the prevalence and concentration. Last, it simulates the *Prevalence in ingested food* and the *Concentration in ingested food*, when the concentration is larger than 0 (Figure A2.5, Table A2.5).



**Figure A2.5** Influence diagram for *Prevalence and concentration* characteristics module.

**Table A2.5** Nodes for *Prevalence and concentration* module.

Title, Identifier, Structure	Description and Definition
Prevalence characteristics Prevalence_character Module	Prevalence characteristics simulates the prevalence of <i>L. monocytogenes</i> as measured at retail, in the packages or units that the consumer would purchase.
Concentration characteristics Concentration_charac Module	Concentration characteristics simulates the <i>L. monocytogenes</i> concentration as measured at retail or source, in the packages or units that the consumer would purchase.
Correlated prevalence & concentration Correlated_prevalenc Module	Correlated prevalence & concentration is the means to specify the rank correlation coefficient between the prevalence and concentration.
Prevalence in ingested food Prevalence_in_ingest Chance node	The Prevalence in ingested food node is the simulated distribution for how often a serving contains any <i>L. monocytogenes</i> contamination. Some servings from a contaminated package will carry no organisms ( $\exp(-m\mu)$ , where the serving size is mg and the concentration is $\mu\text{g}^{-1}$ ). Those probabilities adjust the prevalence estimates that emerge from the Prevalence and concentration characteristics module. Prevalence in ingested food is calculated as follows. For Icebox:=Refrigerator_studies Do For Person:=Risk_group_definitio Do Correlatedprevalence * (1-Exp(-10^Finalconcentration[Refrigerator_studies=Icebox] * Serving_size1[Risk_group_definitio=Person] ) )
Concentration in ingested food Finalconcentration (log <sub>10</sub> CFU/g) Chance node	Concentration in ingested food is the simulated distribution of the concentration of <i>L. monocytogenes</i> in contaminated food at ingestion. Initial concentrations grow into final concentrations according to the growth determined in the Growth module. Final concentrations are restricted to theoretical maximum population densities or stationary phase densities. Using Calculatedfinal:= Initialconcentration + Unconstrgrowthamount Do ( if Maximum_population >0 then ( if Calculatedfinal <= logten(Maximum_population) then Calculatedfinal else logten(Maximum_population) ) else Calculatedfinal )

### A2.2.6 Prevalence characteristics

This implementation models prevalence as measured at retail, in the product packages or units that the consumer would purchase. The *Prevalence parameters* table specifies the  $\alpha$  and  $\beta$  parameters of a Beta distribution. *Prevalence in packages* is defined as Beta ( $\alpha$ ,  $\beta$ ) parameters as appropriate for each food group (Table A2.6).


**Table A2.6 Nodes for Prevalence characteristics module.**

Title, Identifier, Structure	Description and Definition
Prevalence parameters	The Beta distribution, which has support on [0, 1], is a common way to characterize the heterogeneity in the prevalence.
Prevalenceparameters	
Variable node	Determtable(Food_groups,Prevdistrnparameters)(0.424, 0.55, 155.47)
Prevalence in packages	Prevalence in packages samples from the Beta distribution specified by
Prevalenceinpackage	Prevalence parameters and makes the Packaging adjustment required
Chance node	Packaging_adjustment * ( Using localalpha:=Prevalenceparameters[Prevdistrnparameters='alpha'] Do Using localbeta:=Prevalenceparameters[Prevdistrnparameters='beta'] Do Beta(localalpha, localbeta) )
Prevalence distribution parameters	Prevalence distribution parameters indexes the columns of the
Prevdistrnparameters	Prevalence parameters table.
Index node	['alpha', 'beta']

### A2.2.7 Concentration characteristics

Concentration distributions were derived from published studies for two groups of RTE foods. In Figure A2.6, rounded rectangular shapes are variables holding a table of data that describes the empirical distribution function or a set of quantiles, or parameters for a distribution function for the concentrations .

At each iteration, a value is sampled from the distribution and collected into one of the elliptical nodes. The *Initial concentration* node collects all results, still separate, together in the same place. The parallelogram at the bottom of the Figure A2.6, *Concentration table columns*, lists the columns that appear in concentrations tables: Concentration, and Quantile. Data collection and organization from referenced studies provide concentration distributions that represent levels of concentrations in recognizable packages or units of products (Table A2.7).



**Figure A2.6** Influence diagram for *Concentration characteristics* module.

**Table A2.7** Nodes for *Concentration characteristics* module.

Title, Identifier, Structure	Description and Definition
Ice cream concentrations Icecreamconctable Variable node	Ice cream concentrations holds quantile distribution for <i>L. monocytogenes</i> concentration in ice cream.
Ice cream concentration Icecreamconcc Chance node	Ice cream concentration samples from the cumulative distribution that is specified in Ice cream concentrations. Cumdist(Icecreamconctable[Concentration_tables='Quantile'], Icecreamconctable[Concentration_tables='Concentration'])
Pasteurized milk concentrations Pastmilkconctable Variable node	Pasteurized milk concentrations holds quantile distribution for <i>L. monocytogenes</i> concentration in pasteurized milk
Pasteurized milk concentration Pastmilkconcc Chance node	Pasteurized milk concentration samples from the cumulative distribution that is specified in Pasteurized milk concentrations. Cumdist(Pastmilkconctable[Concentration_tables='Quantile'], Pastmilkconctable[Concentration_tables='Concentration'])
Concentration table columns Concentration_tables Index node	Concentration table columns structures the concentration tables. ['Concentration', 'Quantile']
Initial concentration Initialconcentration Chance node	Initial concentration node collects all results, still separate, together in the same place. Determtable(Food_groups)(Icecreamconcc, Pastmilkconcc)

### A2.2.8 Correlated prevalence and concentration

The Analytica™2.0.1 and 2.0.5 software does not directly implement built-in methods for generating random variables with a desired correlation structure. The installation does provide a Library module, *Correlated Distributions*, which provides the mechanics to achieve the desired result. The *Correlated Distributions* library module implements the method of Iman and Conover (1982), which makes the rank correlation between specified variables meet the desired result. So, *Prevalence in packages*, from the *Prevalence characteristics* module and *Initial concentration*, from the *Concentration characteristics* module, are re-ordered to produce rank-correlated *Correlated prevalence* and *Correlated concentration*.

### A2.2.9 Storage characteristics

*Storage temperature* and *Storage time* characterize storage conditions in this exposure assessment. Storage temperature is intended to represent storage in the consumer's refrigerator, after purchase of the food product from retail. Storage time is intended to represent the length of time that the food product is stored at that temperature, measured from retail purchase until the consumer eats a portion. A simple implementation assumes constant temperature. More complicated implementations that depend on quantitative data lacking here could incorporate time and temperature integration.

*Storage temperature* comes from four separate sources in different countries, which were included through the whole exposure to examine the effects of different assumptions about temperatures, or different distributions of temperatures. It is assumed that refrigerator storage temperatures are the same for any food product – unrealistic, but simplifying (Table A2.8).

**Table A2.8** Quantiles for refrigerator temperature (°C) distributions, showing point estimates for cumulative probabilities from four studies.

Audits International (2000)		Johnson et al. (1998)		Sergelidis et al. (1997)		O'Brien (1997)	
°C	Cum. Prob.	°C	Cum. Prob.	°C	Cum. Prob.	°C	Cum. Prob.
0	0	-2.5	0	0	0	0	0
0.14	0.03	-2	0.002	9	0.45	4	0.40
0.83	0.06	-1	0.002	10	0.75	11	1
1.94	0.15	0	0.01	13	1		
3.05	0.37	1	0.02				
4.16	0.74	2	0.07				
5.28	0.80	3	0.11				
6.39	0.91	4	0.18				
7.50	0.98	5	0.30				
8.61	0.99	6	0.44				
9.72	1.00	7	0.76				
10.28	1	8	0.92				
		9	0.96				
		10	0.99				
		11	0.997				
		12	0.998				
		13	1				

*Storage time* represents the length of time that the consumer stores the product before eating a serving from it. Storage time distributions are modelled as specific to the food product under consideration. Following FDA/FSIS (2001), *Minimum time*, *Mode time* and *Maximum time* parameterize storage time distributions, via Triangular(Minimum time, Mode time, Maximum time). Minimum time is set to a constant 0.5 days for all products, but Mode and Maximum are intended to depend on the food. Mode time and Maximum time are allowed to be stochastic. Mode time varies as Uniform( $\pm 20\%$  nominal). Maximum time varies as Uniform( $\pm 50\%$  nominal). Nominal values are listed in Table A2.9. Mode time and Maximum time are strictly related, so that nonsensical values are not generated. Consequently, this implementation calculates an *Indep. Storage time* that aligns the smallest *Mode time* with the smallest *Indep. Maximum* (Table 2.10).

Storage life for pasteurized milk depends on the growth of spoilage bacteria, which depends on temperatures. The effect would be to truncate the time distribution differently at different temperature values. General tendencies would be

the same. Distribution shapes would change. The storage life for pasteurized milk is assumed to be 12 days at 4°C, with storage life at other temperatures determined by the relationship  $Life(T) = 12 \times \left[ \frac{4+7.7}{T+7.7} \right]$  in Neumeier, Ross and McMeekin (1997) and Neumeier et al. (1997). The influence diagram is shown in Figure A2.7.

**Table A2.9**

Post-retail storage times (days).

	Min	Mode	Max
Ice cream	0.5	7	30
Fluid milk, pasteurized	1	5	12

**Table A2.10** Nodes for Storage characteristics module.

Title, Identifier, Structure	Description and Definition
Audit International 2000 Audit2000 Variable node	Audits International 2000 is 1 of 4 sources of refrigerator storage temperatures (Table A2.8).
Refrigerator characteristics Refrigerator Index node	Refrigerator characteristics node structures the data in the refrigerator temperature table. ['Temperature', 'Frequency', 'Cumulative frequency']
Refrigerator studies Refrigerator_studies Index node	Refrigerator studies node maintains a consistent list of the 4 refrigerator temperature source studies. ['Audits International 2000', 'Johnson et al., 1998', 'Sergelidis et al., 1997', 'O'Brien 1997']
Storage temperature Storage_temperature Chance node	Sstorage temperature is sampled from the information in the four refrigerator temperature studies. Cumdist(Audit2000[Refrigerator='Cumulative frequency'], Audit2000[Refrigerator='Temperature'])
Post-retail storage time Post_retail_storage_ Variable node	Post-retail storage time holds minimum, mode and maximum storage time for each Food groups label and for each Updates label. Storage time represents the length of time that the consumer stores the product before eating a serving from it. Storage time distributions are modelled as specific to the food product under consideration. Following FDA/FSIS (2001), Minimum time, Mode time and Maximum time parameterize storage time distributions, via Triangular(Minimum time, Mode time, Maximum time) (Table A2. 9).
Minimum time Minimumtime Chance node	Minimum time extracts the minimum time from Post-retail storage time appropriate to the selected Food groups. Post_retail_storage_[Post_retail_storage_='Minimum']
Mode time Modetime Chance node	Mode time extracts the nominal mode time from Post-retail storage time appropriate to the selected Food groups. Mode time varies as Uniform ( $\pm 20\%$ nominal). Using Temp:=Post_retail_storage_[Post_retail_storage_='Mode'] Do Uniform(0.8*Temp, 1.2*Temp)
Indep. maximum Maximumindeptime Chance node	Indep. maximum extracts the nominal maximum time from Post-retail storage time appropriate to the selected Food groups. Maximum time varies as Uniform( $\pm 50\%$ nominal). Using Temp:=Post_retail_storage_[Post_retail_storage_='Maximum'] Do Uniform(0.5*Temp, 1.5*Temp)
Corr. Maximum Maximumtime Chance node	The Mode time is assumed to follow a Uniform(0.8*mode, 1.2*mode) and the Indep. maximum to follow a Uniform(0.5*maximum, 1.5*maximum). To avoid nonsensical parameter combinations, and to represent what would be a sensible set of conditions, the random mode and random maximum have a correlation coefficient of 1. For Onebyone:=Updates Do Using Another:=Rank(Maximumindeptime[Updates=Onebyone],Run) Do Using Sortedmaximum:=Maximumindeptime[Updates=Onebyone, Run=Sortindex(Another,Run)] Do Sortedmaximum[Run=Rank(Modetime[Updates=Onebyone], Run)]
Storage time Preliminarytime Chance node	Storage time is the storage time before acting to make the time and temperature related. Triangular(Minimumtime, Modetime, Maximumtime)
Truncated storage time Storage_time Chance node	The storage life for pasteurized milk is assumed to be 12 days at 4°C, with storage life at other temperatures determined by the relationship in Neumeyer, Ross and McMeekin (1997) and Neumeyer et al. (1997). (Using local1 := (1643/((Storagetemperature+7.7)^2)) Do Using local2 := (Storage_time1>local1) Do ((Storage_time1*(1-local2))+local2*For local3 := Run Do (If local2[Run=local3] Then (-Truncate((-Storage_time1),(-local1[Run=local3]))) Else 0))) )




Figure A2.7 Influence diagram for *Storage characteristics* module.

### A2.2.10 Growth

The *Growth* module uses simulated *Growth characteristics* to determine the *Growth per day*. If it is assumed that every day, or every part day, has the same *Storage conditions* and the same *Growth characteristics*, then the *Unconstrained growth and die off* can be determined in a straightforward manner, constraining that *Unconstrained growth* by the stationary phase maximum population density. The influence diagram is shown in Figure A2.8 and the noted are described in Table A2.11.




Figure A2.8 Influence diagram for *Growth* module.

**Table A2.11** Nodes for *Growth* module.

Title, Identifier, Structure	Description and Definition
Growth characteristics Growth_characteristi Module	Growth characteristics are characterized by exponential growth rates at 5°C and stationary phase population size for each foodstuff.
Growth conditions Growth_conditions Module	Growth conditions summarizes the growth characteristics and growth conditions.
Unconstrained growth and die off Unconstrgrowthamount (log <sub>10</sub> CFU/g) Chance node	Unconstrained growth and die off is the simple product of Storage time and Growth per day, giving the amount of growth (log <sub>10</sub> CFU/g) over the whole storage time, were growth not constrained in any way. Storage_time * Growthdaily
Growth per day Growthdaily (log <sub>10</sub> CFU/g/day) Chance node	Growth per day adjusts calculated growth (5°C) to Storage temperature. Convert to an EGR at some other temperature via McMeekin et al. (1993). In no growth conditions – zero growth rate or Storage temperature below minimum growth temperature – the zero growth rate remains as is. For Icebox:=Refrigerator_studies Do Using localtemperature:= Storage_temperature[Refrigerator_studies=Icebox] Do if localtemperature <= Minimum_growth_tempe then ( if Growth_rate<=0 then Growth_rate else 0 ) else if Growth_rate<=0 then Growth_rate else Growth_rate * (localtemperature-Minimum_growth_tempe)^2/(5-Minimum_growth_tempe)^2

### A2.2.11 Growth characteristics

Growth characteristics are the exponential *Growth rates*, the *Minimum Growth Temperature* and the *Stationary Phase Population*. The influence diagram is shown in Figure A2.9. The Growth rates node is a table of means and standard deviations for the growth rate, log<sub>10</sub>/day, at 5°C, gleaned from versions of FDA/FSIS (2001). Storage temperature is explicitly accounted for as a dependent condition for growth. It is assumed that the range of growth rates (FDA/FSIS, 2001) samples among the other dependent conditions (a<sub>w</sub>, pH, NaCl, NO<sub>3</sub>). Values are shown in Table A2.12. Growth rate selects values according to Normal(mean, standard deviation) for this exposure assessment. FDA/FSIS (2001) provides maximum Stationary phase population values that change with temperature. Minimum growth temperature is implemented as Triangular(1°C, 1.1°C, 2°C) for this exposure assessment.

### A2.2.12 Growth conditions

The *Growth conditions* module gives a summary of growth and survival. For example, it converts the *Growth rate* simulated for 5°C into a *Generation time*, via log<sub>10</sub>(2)/Growth rate. Also, it summarizes the growth situations, by tabulating from the simulated growth, to report the fraction of cases where the conditions jointly point to growth, no growth and die-off of the population. Nodes on the left-hand side of the diagram (Figure A2.10) are defined elsewhere, but are displayed here for continuity. Storage temperature and Storage time are defined in the Storage conditions module. Growth rate and Minimum growth temperature are defined in the Growth characteristics module. Growth per day is defined in the Growth module. Details of the notes are presented in Table A2.13.




Figure A2.9 Influence diagram for *Growth characteristics* module.




Figure A2.10 Influence diagram for *Growth conditions* module.

**Table A2.12** Nodes for *Growth characteristics* module.

Title, Identifier, Structure	Description and Definition
Growth rates Growth_rates Variable node	FDA/FSIS 2001 suggests these Growth rates. We summarize the growth rates by the mean and standard deviation. Other summaries of growth rates appear in such as Farber and Peterkin (2000: Table 44-10). In order by Food groups, the definition specifies the mean, variance and number of studies. Determtable(Food_groups,Growthtableparameter)(0,Uniform(0.092, 0.434))
FDA/FSIS growth rates Fda_fsis_growth_rate Chance node	FDA/FSIS growth rates Determtable( Food_groups ) (0,Uniform(0.092, 0.434))
Stationary phase population intermediate Stationary_phase_int Chance node	Stationary phase population intermediate is modelled as different for milks and for other foods of interest. Also, it varies with a range of Storage temperature. Using Allotherfoods:=( if Storage_temperature<5 then 10^5 else if Storage_temperature>7 then 10^8 else 10^6.5) Do Using Milks:=(if Storage_temperature<5 then 10^7 else if Storage_temperature>7 then 10^8 else 10^7.5) Do Table(Food_groups) ( Allotherfoods, Allotherfoods, Milks, Milks, Allotherfoods, Allotherfoods, Allotherfoods)
Minimum growth temperature Minimum_growth_tempe Chance node	Farber and Peterkin (2000: Table 44-9) and its references suggest Minimum growth temperature between 1°C and 2°C for foods. The structure of the definition leaves room for the Minimum growth temperature to be different for each Food group, but leaves it the same, regardless of the set of Growth rates. Determtable(Food_groups,Updates)(Triangular(1, 1.1, 2), -1.18, Triangular(1, 1.1, 2), -1.18)
Growth rate Growth_rate Chance node	Though indexed by Updates (WHO/FAO 2000.06.17, FDA/FSIS 2000.05.19), Growth rate uses the FDA/FSIS growth rates for both. Growth rate has the simulated distribution of growth rate, for the food of interest, at 5°C. Table(Updates) (Fda_fsis_growth_rate, Fda_fsis_growth_rate)
Stationary phase population Maximum_population Chance node	Stationary phase population selects only the maximum density from Stationary phase population intermediate, for the selected Food groups. Stationary_phase_int[Food_groups=Food_groups]
Growth table parameters Growthtableparameter Index node	Growth table parameters structures the Growth rates table. ['mean', 'std. dev.', '# studies']

**Table A2.13** Nodes for *Growth conditions* module.

<b>Title, Identifier, Structure</b>	<b>Description and Definition</b>
Generation times Generation_times (h) Chance node	Generation times calculated at the exponential growth rates at 5°C. $24 * \log_{10}(2)/\text{Growth\_rate}$
Growth, no growth, die-off Growth1 Chance node	Growth, no growth, die-off is a simple summary. if Growth_rate<0 then 'Die-off' else if Growth_rate=0 Or Storage_temperature<Minimum_growth_tempe then 'No growth' else 'Growth'
Growth at maximum Growth_at_maximum Chance node	Growth at maximum tabulates how the maximum population density constrains growth. Sometimes, simulated growth is "Below maximum". Sometimes, simulated growth is constrained by the maximum population density set by the lower Storage temperature range ('At Lower'), set by the middle Storage temperature range ('At Mid'), and set by the upper Storage temperature range ('At Higher'). if $10^{\text{Finalconcentration}} < \text{Maximum\_population}$ then 'Below maximum' else if Storage_temperature<5 then 'At Lower' else if Storage_temperature>7 then 'At Higher' else 'At Mid'
Generation times Generation_times1 Chance node	Generation times

### A2.2.13 Study indices

The *Study indices* library module stores 4 index nodes that structure results from the exposure assessment (Table A2.14).

**Table A2.14** Nodes for *Study indices* module.

<b>Title, Identifier, Structure</b>	<b>Description and Definition</b>
Food groups Food_groups Index node	Food groups lists the food commodities that the exposure assessment addresses. ['Ice cream', 'Fluid milk, pasteurized']
Updates Updates Index node	Updates lets the exposure assessment address different sets of storage, time and growth conditions. ['WHO/FAO 2000.06.17', 'FDA/FSIS 2000.05.19']
Annual meals reporting Annualmealsreporting Index node	Annual meals reporting indexes the Annual meals objective node ['Individual', 'Population']
Contaminated or not Contaminated_or_not Index node	Contaminated or not defines the domain of the Chance node Contaminated or not. ['Not contaminated', 'Contaminated']

## A2.3 CONSUMPTION CHARACTERISTICS

### A2.3.1 Overview

The exposure assessments characterize consumption by meal size and meal frequency, noting and reporting differences in consumption patterns in the population sub-groups with different susceptibility. The meal or serving size is the estimated portion that people eat and has a

distribution estimated from survey data. Similar surveys derive the frequency of eating specific RTE foods. Sources of consumption data are discussed earlier in Sections 3.1, 4.1.2 and 4.1.4.3 of this report. For the pasteurized milk and ice cream assessments, data describe the consumption characteristics of adult Canadians.

### **A2.3.2 The data**

Consumption characteristics are derived from 24-hour recall data from Canadian Federal-Provincial Nutrition Surveys (CFPNS, 1992–1995), which addressed the nutritional habits of non-institutionalized adults between 18 and 74 years old in the Provinces of Québec, Nova Scotia, Saskatchewan, Alberta and Prince Edward Island. Results are based on data from a total of 10 162 individual respondents from the 1990 Nova Scotia Nutrition Survey (2212 respondents), the 1990 Québec Survey (2118 respondents), the 1993/94 Saskatchewan Survey (1798 respondents), the 1994 Alberta Survey (2039 respondents) and the 1995 Prince Edward Island Survey (1995 respondents). Detailed one-day 24-hour recall data were used to examine consumption of various foods that would help to describe the consumption frequency and amounts eaten for the food groups relevant to this exposure assessment. All survey estimates are weighted to adjust for the sample design, and balance the ages and provinces according to their representation in the populations of those provinces. It is assumed that the remainder of the Canadian adult population eats like this group.

By using single occasion or daily consumption, estimates represent the fraction of the total population consuming the selected food on a given day, essentially a day at random. Food intakes are subject to day-to-day variation among individuals. Thus, the estimates are not indicative of “usual” intake, but are more indicative of the episodic intake with which would be associated foodborne illness. Distributions of “usual” intakes are unobservable in the 24-hour, one-day recall data that the Nutrition Surveys provide. Bureau of Biostatistics and Computer Applications has developed methods to remove the day-to-day within person variability (Junkins and Laffey, 2000; Junkins, Laffey and Weston, 2001; Hayward, [2001]). Those synthesized distributions of “usual” intakes are less heavily tailed than distributions of intakes that retain inter- and intra-person variability as is appropriate for the consumption distributions for these exposure assessments.

There is some uncertainty due to extrapolation of the results to 365 days’ experience, when simulating factors such as annual consumption in the population, or to any reference period. The consumption of milk or ice cream were represented by reference to consumption of any of several foodcode categories, a classification system that the surveys employed. Selection of foodcodes from the nutrition surveys’ databases was intended to reflect both consumption frequency and amount consumed on eating occasions. The information from individual all-eating episodes that included the food was used, except when the eating episode involved preparation such as cooking. When the food was an ingredient in the serving, an appropriate amount of the food to include was derived or estimated. There are uncertainties associated with this representation of intended foods by particular foods identified in the surveys. Additionally, there might be underreporting or overreporting errors associated with respondent errors and misclassification errors. Trained interviewers estimated amounts consumed on respondents’ eating occasions. This is methodologically preferable to a practice that lets a respondent estimate the amounts consumed. However, it is recognized that the amounts recorded contain reporting errors and variability due to the interviewers’ estimation methods.

It was difficult to adequately identify how to appropriately aggregate the sometimes many individual foods into the same eating occasion within the respondent's reference day. Therefore, all eating episodes for a food on the same day were aggregated into a daily consumption amount. That practice loses the distinction that one might wish to strike among occasions when the food was consumed alone, as an ingredient in a recipe or as only one element among several elements in a meal. Although the Nutrition Surveys distinguish consumption at an individual's home from consumption at another establishment outside the home, the distinction has been ignored for this exposure assessment. Past work suggests that consumption frequency differences and consumption amounts differences do exist between home and away consumption. Consequently, the consumption that is incorporated is assumed to represent a combination of all eating occasions. Combined independently with foodborne contamination, it is implicitly assumed that there are no differences in contamination rates and concentrations between food consumed at home and food consumed away from home (E.A. Junkins, pers. comm., 2000; M. Vigneault, pers. comm., 2000).

### **Gender and Age groups**

The Nutrition Surveys do not classify respondents into groups to which might be attributed characteristics like higher susceptibility to foodborne illness. Rather, membership in non-susceptible and susceptible groups is imputed from Gender and Age attributes. Consumption characteristics of susceptible and non-susceptible groups of individuals, then, are different only in the manner that constituent Gender and Age characteristics are present in those groups. A susceptible group that is represented by elderly consumers would therefore possess consumption characteristics that differ from the non-susceptible group, solely because of differences between the consumption characteristics of elderly consumers and other consumers.

### **Simulating consumption amounts**

To make it easier to specify the consumption distribution, some conventions were followed. Distributions were described by sampling in the correct proportions, from distributions that describe consumption in 4 Age  $\times$  2 Gender ranges, both for frequency of consumption and consumption amount. Eating episodes, both at home and away from home, were combined into the same distribution, capturing some variability but not distinguishing their separate influences. Non-susceptible and susceptible populations were defined by assuming that some fraction of each Gender  $\times$  Age group is more susceptible. It is assumed also that the consumption characteristics of all persons of the same age and gender are the same, whether the person is in the susceptible or non-

**Table A2.15** Fraction of population in Non-susceptible and Susceptible risk groups attributed to Gender  $\times$  Age groups.

<b>Susceptible group</b>			
Age	Female	Male	Total
18–34	0.16	0.04	0.20
35–49	0.08	0.03	0.12
50–64	0.02	0.02	0.04
65–74	0.35	0.29	0.64
	0.61	0.39	1.00
<b>Non-susceptible group</b>			
Age	Female	Male	Total
18–34	0.20	0.23	0.43
35–49	0.17	0.18	0.35
50–64	0.11	0.11	0.22
65–74	0.00	0.00	0.00
	0.48	0.52	1.00

susceptible group. So, consumption in the susceptible and non-susceptible groups can be correctly simulated by sampling in the correct proportions from consumption characteristics captured for the Gender  $\times$  Age groups. However, differences in the distributions of consumption for persons from the non-susceptible and susceptible groups are attributable only to the different Gender  $\times$  Age group make-up of the groups. The values used are presented in Tables A2.15 and A2.16.


**Table A2.16** Fraction of population in Gender  $\times$  Age group attributed to Susceptible group.

Age	Female	Male
18–34	0.12	0.03
35–49	0.08	0.03
50–64	0.03	0.03
65–74	1.00	1.00

## A2.4 NON-SUSCEPTIBLE AND SUSCEPTIBLE POPULATIONS

Among adults, susceptible groups are defined to include all adults 65 and older, pregnant women (1.3% of the population) and individuals with suppressed immune systems and certain medical conditions, such as cancer and recent organ transplantation (3.3% of the population) (Miller, Whiting and Smith, 1997). Different fractions of the population in Age and Gender groups are attributed to the susceptible group (Table A2.15), so non-susceptible and susceptible groups include gender and age groups in different fractions (Table A2.16). Fractions attributed to a Gender  $\times$  Age group depend on the population size for that group, and are criteria for attributing risk categorization to that group. When individual food products are considered, the fraction of individuals who are susceptible depends, too, on consumption characteristics for the population. Susceptible groups would include also all children under 6 months, or perhaps 0–4 years old (J.M. Farber, pers. comm., 2000), some fraction of individuals under 18 years old, and all persons older than 74 years. Among adults aged 18–74, 15% would fit into the susceptible group and 85% would fit into the non-susceptible group when the definition described here is applied (Tables A2.15, A2.16 and A2.17).

An alternative approach follows one suggested in FDA/FSIS (2001). Observations from the FoodNet database describe listeriosis in the United States of America by age group. One might hypothesize that the incidence in an age group depends on, particularly, susceptibility, consumption characteristics and population representation. United States of America incidence data were used and combined with Australian populations in different age groups. Figure A2.11 scales those incidence data so that the incidence in the 10–19-year-old age group corresponds to 1. The relative incidence in the <30 days age group is 300 times the incidence in the 10–19 age group. Similarly, consumption characteristics among populations could be used to account for consumption differences. The remaining differences would affect the different age groups' susceptibility to listeriosis, perhaps forming a surrogate representation for the contrast between susceptible and non-susceptible populations.



**Figure A2.11** Relative per capita incidence of listeriosis (See text for details of calculation).

**Table A2.17** Population of Canada allocated to susceptible and non-susceptible groups, using factors from Table A2.15.

Population of Canada				Susceptible population size				Non-susceptible population size			
Age	Male	Female	Total	Age	Male	Female	Total	Age	Male	Female	Total
0-4	911 028	866 302	1 777 330	0-4	911 028	866 302	1 777 330	0-4	0	0	0
5-17	2 738 162	2 598 365	5 336 526	5-17	82 145	77 951	160 096	5-17	2 656 017	2 520 414	5 176 431
18-34	3 711 154	3 591 613	7 302 768	18-34	111 335	430 994	542 328	18-34	3 599 820	3 160 620	6 760 439
35-49	3 823 789	3 802 863	7 626 652	35-49	114 714	304 229	418 943	35-49	3 709 075	3 498 634	7 207 709
50-64	2 403 311	2 453 603	4 856 914	50-64	72 099	73 608	145 707	50-64	2 331 212	2 379 995	4 711 207
65-74	1 000 723	1 134 443	2 135 166	65-74	1 000 723	1 134 443	2 135 166	65-74	0	0	0
74+	644 742	1 069 989	1 714 731	74+	644 742	1 069 989	1 714 731	74+	0	0	0
	15 232 909	15 517 178	30 750 087		2 936 785	3 957 516	6 894 301		12 296 124	11 559 662	23 855 786

SOURCE: Adapted from: Statistics Canada, [www.statcan.ca/english/Pgdb/People/Population/demo10a.htm](http://www.statcan.ca/english/Pgdb/People/Population/demo10a.htm) (November 2000), except for 15–19 years age group prorated into 5–17 years and 18–34 years age groups in table above.

## A2.5 HOME STORAGE CHARACTERISTICS

### A2.5.1 Home refrigeration temperatures

Four studies<sup>1</sup> contributed information about the distribution of refrigeration temperatures, important as one of the main determinants of growth of *L. monocytogenes* during storage. Audits International (2000) surveyed homes in the United States. Johnson et al. (1998) surveyed persons 65 years and older in the United Kingdom. Sergelidis et al. (1997) published results from a survey of homes in Athens, Greece. O'Brien (1997) also considered homes in the United States of America (Figure A2.12). Quantiles from Johnson et al. (1998)

1. A comment made on a late draft of this report pointed to two other references. Notermans et al. (1997) report refrigerator temperatures for households in the Netherlands, for pasteurized milk. Willocx, Hendrickx and Tobbacq (1993) report, *inter alia*, refrigerator temperatures for Belgian residences.

are approximately 3°C higher in the middle of the distribution than the ones in Audits International (2000). Based on limited presentation, though, the quantiles from Sergelidis et al. (1997) are approximately 3°C higher still in the middle of the distribution. O'Brien (1997) and Sergelidis et al. (1997) report 1 or 2 quantiles from the core of their temperature distributions (Table A2.8). It has been assumed that the same storage temperature distribution is appropriate for all RTE foods of interest. Further, it has been assumed that the food is stored at the same temperature throughout its shelf life.




Figure A2.12 Four storage temperature distributions.

### A2.5.2 Home storage times

No specific references that describe the length of time that consumers store foods in the home before eating were found<sup>2</sup>. Several characteristics that simulated results should try to emulate, at least qualitatively, might be considered. First, there should be at least a short, minimum storage time associated with all food consumed, representing, at least, the time from retail purchase to the individual's home. Second, storage time distributions that describe variability should have some maximum time that should be constrained by when the consumer would no longer accept the product. The maximum time would be related to the product's shelf-life, but should also reflect variability among individuals' practices of choosing whether to consume foods that have been stored beyond recommended limits. Some authors have studied the relationship between food spoilage, as represented by growth of some spoilage bacteria to high concentrations, and the organoleptic qualities of the food – qualities that help individuals to decide whether to eat a food or not (Priepke, Wei and Nelson, 1976; King, Henderson and Lill, 1986; Garcia-Gimeno and Zurera-Cosana, 1997). Several countries use  $10^6$  CFU/g concentrations of mesophiles as a guideline for acceptability.

2. A comment on a late draft of this report pointed to Notermans et al. (1997), which reports summary results for the storage time (after pasteurization of the milk) for pasteurized milk in households in the Netherlands.

In another manner, FDA/FSIS (2001) provides some useful information about storage time to consumption among individuals (Table A2.18). For the present, this exposure assessment uses those same criteria. FDA/FSIS (2001) accounts for individual variability in the storage time by describing a distribution specified by Triangular(Minimum, Mode, Maximum). That work also introduces variability and uncertainty in their representation by varying the Mode uniformly in the interval Mode  $\pm 20\%$  and the Maximum in the interval Maximum  $\pm 50\%$ .

**Table A2.18** Storage time distribution parameters.

	Minimum	Mode	Maximum
Ice cream	0.5	7	30
Fluid milk, pasteurized	1	5	12

### A2.5.3 Storage time and temperature

Clearly, storage time and storage temperature are not independent. Spoilage actions would severely truncate storage time, forcing shorter storage times to happen with higher storage temperatures. No studies that describe such a relationship directly were found. Studies that directly relate spoilage bacteria to organoleptic qualities might be useful, but have not been explored. A simple implementation for pasteurized milk assumes that organoleptic preferences that would truncate storage time can be related to storage life. Storage life for pasteurized milk depends on the growth of spoilage bacteria, which depends on temperatures. The storage life for pasteurized milk is assumed to be 12 days at 4°C, with storage life at other temperatures determined by the relationship  $Life(T) = 12 \times \left[ \frac{4+7.7}{T+7.7} \right]$  (Neumeyer, Ross and McMeekin, 1997; Neumeyer et al., 1997). To account for variability among individuals, the relationship  $time \propto temperature^{-1}$  would not be deterministic and this relationship would be directed only to constrain the most extreme storage length at a given temperature.

## A2.6 GROWTH CHARACTERISTICS

### A2.6.1 Introduction

Extensive discussions about the microbial ecology of *L. monocytogenes*, predictive microbiology and growth characteristics are discussed elsewhere in this report. For the assessment examples, it has been assumed that the *L. monocytogenes* organisms were in the food sufficiently long for the lag phase to have passed. With that assumption, growth dynamics can be described with the exponential growth rate alone. It is assumed, further, that growth characteristics that must be explicitly accounted for are: the exponential growth rates, the minimum growth temperature and the stationary phase population.

### A2.6.2 Growth rates

It is assumed that there is no growth or decline of *L. monocytogenes* populations in contaminated ice cream. For the milk exposure assessment example, growth rates reported in FDA/FSIS (2001) have been used to capture level and variability in growth of *L. monocytogenes*. Specific parameterization is given in Table A2.19, where growth rates

refer to growth at 5°C. FDA/FSIS (2001) reports that in their referenced studies, study-specific growth rates were converted from the growth rate at a specific study temperature other than 5°C using relationships from McMeekin et al., 1993. At a temperature T°C other than 5°C, growth is calculated using the relationship  $\sqrt{\mu_T} = \sqrt{\mu_5}(T - T_{\min})(5 - T_{\min})^{-1}$ , where  $\mu$  is the growth rate and  $T_{\min}$  is the minimum growth temperature. Storage temperature as a dependent condition for the growth rate has been explicitly accounted for. However, it is assumed that the assumed distribution of growth rates effectively samples among the other dependent conditions ( $a_w$ , pH, NaCl, NO<sub>3</sub>) in the same proportions that would occur in real environments. Bovill et al. (2000) note that competitive flora in the growth environment and the physiological state of the *L. monocytogenes* organisms might also be considered to be growth conditions. There is additional uncertainty in the estimated growth rates, not explicitly accounted for. Study methods and measurements contribute generally random effects that increase variability in replicated results for a given set of conditions, and therefore contribute uncertainty regarding what the true rate would be at those conditions.

Growth rates in Table A2.19 have been converted to refer to growth at 5°C, from the growth rate at a specific study temperature, using the square root relationship (McMeekin et al., 1993). So, growth rates at 5°C, as a baseline, do not explicitly account for variability and uncertainty in model extrapolation (or interpolation) from a study temperature back to 5°C.

### A2.6.3 Stationary phase population

This implementation of maximum population densities is straightforward. FDA/FSIS (2001) reports stationary phase population values that change with storage temperature (Table A2.19). The stationary phase population is viewed as one of many constraints on the growth of *L. monocytogenes*. Other characteristics include competition with other microorganisms and growth of total spoilage bacteria populations to the extent that the food is not organoleptically acceptable, but these other characteristics are not accounted for.

### A2.6.4 Minimum growth temperature

Based on Farber and Peterkin (2000), minimum growth temperature is implemented as Triangular(1°C, 1.1°C, 2°C) for this exposure assessment. Alternatives, such as -1.18°C (FDA/FSIS, 2001), set lower minimum growth temperature for *L. monocytogenes* than are implemented here. It is assumed also that minimum growth temperature is the same for the example foods to which it is applied.

### A2.6.5 Implementation of microbial growth

The amount of growth using *daily growth* × *days storage* is calculated and applied to initial concentrations using  $Concentration_{Final} = Concentration_{Initial} + Growth$  to get final concentrations, when quantities are expressed on a log<sub>10</sub> scale. Growth rates at a stochastic storage temperature are adjusted for, using the relationship  $\sqrt{\mu_T} = \sqrt{\mu_5}(T - T_{\min})(5 - T_{\min})^{-1}$  (McMeekin et al., 1993). Doing so incorporates variability associated with storage temperature, but does not explicitly incorporate uncertainty in extrapolating from a growth rate at 5°C to a growth rate at another temperature. The amount of growth until consumption of a portion is the simple product of the daily growth rate and the length of the storage time. This incorporates variability associated with the storage time, but assumes constant growth

rate over the whole storage time. Growth is constrained so that final concentrations cannot be simulated to exceed the maximum population density.

**Table A2.19** Population growth characteristics for *L. monocytogenes*, giving growth rates at 5°C and stationary phase populations at various temperatures.

	Growth rate distribution	Stationary population		
		<5°C	5°C–7°C	>7°C
Ice cream	0			
Milks	Uniform(0.092, 0.434)	Milks 10 <sup>7</sup>	10 <sup>7.5</sup>	10 <sup>8</sup>

SOURCE: FDA/FSIS, 2001.

## A2.7 PREVALENCE AND CONCENTRATION

This exposure assessment model copies the practice that separates prevalence of contaminated servings and the concentration of *L. monocytogenes* in contaminated servings. This practice is similar to some published quantitative risk assessments (Cassin, Paoli and Lammerding, 1998; Lindqvist and Westöö, 2000), but differs from others (Bemrah et al., 1998, in part; FDA/FSIS, 2001). The practice separates concentration zeros (non-prevalence) from concentration non-zeros. First, the literature presents large data sets that count qualitatively positive and qualitatively negative samples. Concentrations, when presented, come from the typically small number of qualitatively positive samples. Second, it makes the simulation more efficient. The same 10<sup>k</sup> iterations can define the probability of contaminated product and the distribution of concentrations, given contaminated product. Were both zero and non-zero concentrations combined, then that 10<sup>k</sup> simulated observations would generate ~10<sup>k-m</sup>, with *m* generally smaller than 1, number of zeros and only ~10<sup>m</sup> non-zeros, reducing the amount of precision that the simulation generates about the concentration distribution.

This implementation acts as if the declarations that positive and negative samples make are exact. Hence it calls the concentration in qualitatively negative samples exactly 0 CFU/g. The concentration in qualitatively negative samples should be modelled as random variables on [0, ∞).

## A2.8 COMBINING INDEPENDENT PREVALENCE ESTIMATES

### A2.8.1 Introduction

Prevalence estimates for the presence of *L. monocytogenes* in foods come from surveys that typically provide summary information that includes the number of samples detected positive and the total number of samples tested. In some cases, though rarely, a detailed study design is also provided. Most studies give some context for the source of the samples – geography, food types or textures, points of origin, raw materials used or motivation. Most studies describe the methods used to test for *L. monocytogenes* presence. Most often, research has come via the microbiological literature. Some research has come from reports issued by national agencies. The food industry has also provided data sets. One could include prevalence estimates whose source is a modelled estimate, as is common in a quantitative risk

assessment. When several studies are available, it is useful to take advantage of the observed variability between study estimates to provide a proxy model for including uncertainty and variability in a probabilistic risk assessment. To this end, it is assumed that  $k$  combinable studies are available, providing summary data  $\{(Y_i, n_i); i = 1, \dots, k\}$ , where  $Y$  is the number of samples positive for *L. monocytogenes* and  $n$  is the number of samples. This assumes that, within studies, sample designs behave as simple random samples, that samples are independent, and that there is constant probability of a positive sample.

### A2.8.2 Beta-binomial model for combining prevalence estimates

A simple assumption about the stochastic structure of a collection of studies gives binomial variability to the individual study estimates and a Beta distribution to the between-study variability of the true study prevalences. Also, this assumes that there are no overriding factors that are present that would group the studies into subsets, part of the assumption that the studies can be combined with a simple mixing distribution. More formally, the following two-stage model is assumed:  $Y_i | n_i, \pi_i \sim \text{Binomial}(n_i, \pi_i)$ ,  $i = 1, \dots, k$  and  $\pi_i \sim \text{Beta}(\alpha, \beta)$ . For risk assessments, the mixing distribution is of importance. The role played by the distribution of the true study prevalence values is understood in the following way. If the mixing distribution is primarily a description of uncertainty, with a common fixed underlying prevalence value, then information from these several studies could be simply combined to give a more precise estimate of that single, fixed, true prevalence than the individual studies give. In this case, the Beta distribution plays the role of a prior density on the prevalence parameter. However, if the distribution of true study prevalence values also reflects variability in the prevalence value, then increasing the number of observations does not reduce this variability, though it can improve knowledge of the underlying distribution. In this case, the Beta distribution is an intrinsic component of the variability of the phenomenon under consideration among circumstances, situations or scenarios. Information from the studies can be appropriately combined to estimate the unknown parameters of that Beta distribution (Ross, pers. comm., 2000). There are a number of approaches available for estimating the parameters  $\alpha$  and  $\beta$  of the Beta mixing distribution (Vose, 2000).

### A2.8.3 Other alternatives appropriate to some circumstances

Alternatives can be found appropriate to some circumstances that Lindqvist and Westöo (2000) and Vose (2000) illustrate. Lindqvist and Westöo (2000) present prevalence data that are proportions  $\{p_j, j = 1, \dots, k\}$ , where the sample sizes and the numbers of positive samples are either ignored or not reported. Those authors pool the observed data, treating them as independently and identically distributed from a distribution that they describe by the quantiles of that pooled sample. There, quantiles are defined by associating the  $j$ th largest observed fraction  $p_j$  with the  $j(k+1)^{-1}$ th point of the distribution. Such a derivation is appropriate when the sample sizes used to estimate the individual fractions are the same or nearly the same, so that they are ignorable. Retaining the sample sizes, nevertheless, is useful to properly account for the uncertainty that one would associate with the fractions as estimates of true fraction values drawn from that empirically defined distribution. The true fraction values play the same role as described above. They can be understood as describing the variability among the true values of the fraction obtained under the conditions that the pooled sample describes. Alternatively, they can be understood as an expression of the uncertainty about the single true prevalence for that same population, from which the samples

form independent observations. Vose (2000) discusses several methods to use with an example data set of fractions  $\{p_j, j = 1, \dots, k\}$ , where the sample sizes and the numbers of positive samples are not reported. He specifies a Beta mixing distribution for the true values that the data set observes and describes how to estimate the parameters,  $\alpha$  and  $\beta$ , by maximum likelihood methods and by the method of moments. He also describes in brief a procedure similar to the one that Lindqvist and Westöo (2000) carry out with their pooled data.

Alternatively, some knowledge of the population might be constructed by considering how to appropriately mix the various conditions of the studies sampled from, rather than basing the mixing distribution for the variability or uncertainty about the true prevalence values on the Beta distribution. One might consider mixing distributions for recognizable parts of a food supply: geographical, food type, point of origin, or raw materials used.

## **A2.9 DISTRIBUTIONS FOR *L. MONOCYTOGENES* CONCENTRATIONS IN FOODS**

### **A2.9.1 Empirical distribution functions and fitted distributions**

Small samples of observations for *L. monocytogenes* concentration in contaminated samples capture the distribution only with some uncertainty, both in the centre of the distribution and in the tails. Of particular concern is the upper tail of the distribution, where large concentrations sit. Studies that were reviewed seldom record high concentrations, or only under exceptional circumstances, making it difficult to model the thickness of the tail. Theoretical constraints on the length of the tail probably can be derived from predictive microbiology, but these require knowledge of growth conditions such as temperature and medium, and might be so much larger than empirical data produce that they would be somewhat unrealistic for practical use. Empirical distributions, too, are somewhat limited in their ability to capture the distribution very precisely in the upper tails. Confidence intervals can capture some notion of uncertainty, but will be a constant width in the tails, above the largest recorded observation. Uncertainty about the whole distribution can be captured non-parametrically by determining confidence intervals about the empirical density function or the empirical distribution function, or as a summary of the empirical distribution at selected quantiles. Last, given some assumptions, one can capture the shape of the distribution by fitting parametric distributions to the data. Uncertainty can be captured by varying the parameters among a confidence set, encompassing all combinations of the parameters that produce distributions that are consistent with the data. The distributions themselves capture variability among *L. monocytogenes* concentrations in different conditions. Parameter uncertainty and confidence intervals may be considered to describe some combination of that variability and uncertainty about that variability.

### **A2.9.2 Families of distributions**

One alternative is to fit an analytical distribution to the data. The families of distributions considered as candidates for describing the concentration distribution should, first, respect the domain of the distribution. As used here, concentrations in contaminated foods have support on  $(0, \infty)$  or on a subset, truncating  $(0, \infty)$  at a minimum and at a maximum value. Second, consideration of candidate probability distributions would be restricted to ones that refer to a continuous random variable, and not a discrete random variable. FDA/FSIS (2001), for

example, considered candidates like the lognormal distribution, the logistic distribution (folded or half-logistic, since the logistic distribution is defined on  $(-\infty, \infty)$ ) and the Beta distribution. Method of moments could be used to estimate parameters for the analytical distributions. Preferable would be to use maximum likelihood methods. Nevertheless, some attention to the fit that the analytical distribution provides for the data, and in which parts of the domain, and goodness-of-fit criteria for the fit over the whole range of the data, should be considered. Distributions selected should also represent what is known about the sampling methods. That is, a point estimate is made about the concentration in an amount of product based on a small sample.

### **Minimum and maximum concentrations**

Setting limits on the length of upper and lower tails can be straightforward and heuristic. When working with concentration distributions in this exposure assessment, the extent of the lower limit of contamination has been set to 0.04 CFU/g (1 CFU per 25 g), a lower detection limit, in effect, for every foodstuff. Upper limits are often set based on authors' suggestions, or set a judged limit larger than the largest observation. In some studies, the largest observed concentration stands as the upper limit, though this might be considered to be unrealistic. A more rigorous approach to setting maxima would consider the operating characteristic curve that is associated with the sample size and sample design of the studies that form the data sets. Minimum and maximum concentrations might also be used in conjunction with fitted distribution functions. The distribution function would define the shape of the distribution; the limits would define the domain of the distribution.

### **Heterogeneity of the organism in the package**

Data collection and organization from referenced studies provide concentration distributions that represent levels of concentrations in recognizable packages or units of products, or give measurements from which one makes an inference about the concentration in the package or unit of product.

## **A2.10 REFERENCES CITED IN APPENDIX 2**

- Audits International. 2000. 1999 U.S. Food Temperature Evaluation. Design and Summary Pages. Audits International and U.S. Food and Drug Administration. 13p.
- Bemrah, N., Sana, M., Cassin, M.H., Griffiths, M.W. & Cerf, O. 1998. Quantitative risk assessment of human listeriosis from consumption of soft cheese made from raw milk. *Preventative Veterinary Medicine*, **37**: 129–145.
- Bovill, R., Bew, J., Cook, N., D'Agostino, M., Wilkinson, N. & Baranyi, J. 2000. Predictions of growth for *Listeria monocytogenes* and *Salmonella* during fluctuating temperature. *International Journal of Food Microbiology*, **59**: 157–165.
- Cassin, M.H., Paoli, G.M. & Lammerding, A.M. 1998. Simulation modeling for microbial risk assessment. *Journal of Food Protection*, **61**(11): 1560-1566.
- CFPNS [Canadian Federal-Provincial Nutrition Surveys]. 1992–1995. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada. See also: Karpinski & Nargundkar, 1992; Junkins & Karpinski, 1994; Junkins, 1994; Junkins & Laffey, 2000; Junkins, Laffey & Weston, 2001.

- Farber, J. M., & Peterkin, P.I. 2000. *Listeria monocytogenes*. pp. 1178–1232, in: B.M. Lund, T.C. Baird-Parker and G.W. Gould (eds). *The Microbiological Safety of Food*. Vol. 2. Gaithersburg, Maryland: Aspen Publishers.
- FDA/FSIS [U.S. Food and Drug Administration/Food Safety and Inspection Agency (USDA)]. 2001. Draft Assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. Center for Food Safety and Applied Nutrition (FDA) and Food Safety Inspection Service (USDA) (Available at: [www.foodsafety.gov/~dms/lmrisk.html](http://www.foodsafety.gov/~dms/lmrisk.html)). [Report published September 2003 as: Quantitative assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. Available at: [www.foodsafety.gov/~dms/lmr2-toc.html](http://www.foodsafety.gov/~dms/lmr2-toc.html)].
- Garcia-Gimeno, R.M. & Zurera-Cosana, G. 1997. Determination of ready-to-eat vegetable salad shelf-life. *International Journal of Food Microbiology*, **36**: 31–38.
- Hayward, S. [2001]. Multivariate adjustment of nutrient intakes. Draft Technical Report, Bureau of Biostatistics and Computer Applications, Food Directorate, Health Products and Food Branch, Health Canada, Ottawa.
- Iman, R.L. & Conover, W.J. 1982. A distribution-free approach to inducing rank correlation among input variables. *Communications in Statistics*, B. **11**(3): 311–334.
- Johnson, A.E., Donkin, A.J., Morgan, K., Lilley, J.M., Neale, R.J., Page, R.M. & Silburn, R. 1998. Food safety knowledge and practice among elderly people living at home. *Journal of Epidemiology and Community Health*, **52**: 745–748.
- Junkins, E. 1994. Saskatchewan Nutrition Survey 1993/94. Methodology for estimating usual intake. BBCA Technical Report E451311-005. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada.
- Junkins, E. & Karpinski, K. 1994. Enquête québécoise sur la nutrition. Méthodologie pour estimer l'apport habituel, les statistiques sommaires et les erreurs-types. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada.
- Junkins, E. & Laffey, P. 2000. Alberta Nutrition Survey 1994. Methodology for estimating usual intake. BBCA Technical Report E451311-006. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada.
- Junkins, E., Laffey, P. & Weston, T. 2001. Prince Edward Island Nutrition Survey 1995. Methodology for estimating usual intake. BBCA Technical Report E451311-007. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada.
- Karpinski, K. & Nargundkar, M. 1992. Nova Scotia Nutrition Survey. Methodology Report. BBCA Technical Report E451311-001. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada.
- King, G.A., Henderson, K.G. & Lill, R.E. 1986. Asparagus: effect of controlled atmosphere storage on shelf-life of four cultivars. *New Zealand Journal of Experimental Agriculture*, **14**: 421–424.
- Lindqvist, R. & Westö, A. 2000. Quantitative risk assessment for *Listeria monocytogenes* in smoked or gravad salmon/rainbow trout in Sweden. *International Journal of Food Microbiology*, **58**: 181–196.
- McMeekin, T.A., Olley, J., Ross, T. & Ratkowsky, D.A. 1993. *Predictive Microbiology. Theory and Application*. Taunton, UK: Research Studies Press. 340p.
- Miller, A.J., Whiting, R.C. & Smith, J.L. 1997. Use of risk assessment to reduce listeriosis incidence. *Food Technology*, **51**: 100–103.

- Neumeier, K., Ross, T. & McMeekin, T.A. 1997. Development of a predictive model to describe the effects of temperature and water activity on the growth of spoilage pseudomonads. *International Journal of Food Microbiology*, **38**: 45–54.
- Neumeier, K., Ross, T., Thompson, G. & McMeekin, T.A. 1997. Validation of a model describing the effects of temperature and water activity on the growth of psychotrophic pseudomonads. *International Journal of Food Microbiology*, **38**: 55–63.
- Notermans, S., Dufrenne, J., Teunis, P., Beumer, R., te Giffel, M. & Weem, P. 1997. A risk assessment study of *Bacillus cereus* present in pasteurized milk. *Food Microbiology*, **14**: 143–151.
- O'Brien, G.D. 1997. Domestic refrigerator air temperatures and the public's awareness of refrigerator use. *International Journal of Environmental Health Research*, **7**: 141–148.
- Priepke, P.E., Wei L.S. & Nelson, A.I. 1976. Refrigerated storage of prepackaged salad vegetables. *Journal of Food Science*, **41**: 379–382.
- Sergelidis, D., Abraham, A., Sarimvei, A., Panoulis, C., Karaioannoglou, P. & Genigeorgis, C. 1997. Temperature distribution and prevalence of *Listeria* spp. in domestic, retail and industrial refrigerators in Greece. *International Journal of Food Microbiology*, **34**: 171–177.
- Statistics Canada (2000), "Population by sex and age", <http://www.statcan.ca/english/Pgdb/People/Population/demo10a.htm>, (2000.November)
- Vose, D. 2000. *Risk analysis: a quantitative guide*. 2nd edition. Chichester, United Kingdom: Wiley.
- Willox, F., Hendrickx, M. & Tobback, P. 1993. Temperatures in the distribution chain. pp. 80–99 [oral presentation], in: Proceedings of the First European "Sous vide" cooking symposium. Leuven, Belgium, 25–26 March 1993.



## Appendix 3.

# Predictive microbiology: concepts, application and sources

Predictive microbiology involves the systematic study and quantification of microbial responses to environments in foods and may be considered as the application of research concerned with the quantitative microbial ecology of foods. It is based upon the premise that the responses of populations of microorganisms in a defined environment are reproducible. By characterizing environments in terms of those factors that most affect microbial growth and survival, it is possible from past observations and experience to predict the responses of those microorganisms in other, similar environments. Ideally, the patterns of microbial behaviour are integrated with knowledge of the physiology of microbes. This knowledge can be expressed very succinctly using the language of mathematics, in the form of mathematical models. Those models can be considered as “condensed knowledge”.

Predictive microbiology models provide a way to estimate changes in *L. monocytogenes* levels in foods as the product moves through the production-to-consumption chain. To make those estimates, periods, temperatures, product composition and concentrations of *L. monocytogenes* at some other point in the chain are required.

This section provides practical guidance for the application of predictive microbiology models in exposure assessment. Predictive microbiology has been extensively reviewed (Farber, 1986; Ross and McMeekin, 1994; Buchanan and Whiting, 1997; Ross, Baranyi and McMeekin, 1999; McDonald and Sun, 1999). McMeekin et al. (1993) provide a good introduction to the concept and its practical application.

The information below is drawn largely from Ross, Baranyi and McMeekin (1999).

### A3.1 SOURCES OF GROWTH RATE MODELS AND DATA

Many data (*see* ICMSF, 1996) and many models for prediction of the growth rate of *L. monocytogenes* are available (*see* Table A3.1, at the end of this appendix). In general, *L. monocytogenes* responds to environmental factors with the same patterns of response as other vegetative microorganisms and can be described by the same forms of model that describe growth rate responses of other organisms (Ross, 1993; Wijnz et al., 1993; Tienungoon, 1998). However, it is reported that the temperature–growth rate relationship of *L. monocytogenes* is not as well described by existing models as it is described for other organisms, particularly at low temperatures that cause slow growth rates (Bajard et al., 1996; Ross, 1999). Generation times of *L. monocytogenes* under a range of conditions can be estimated easily using models such as most of those listed in Table A3.1. One can easily incorporate a published model into spreadsheet software.

### A3.2 PRACTICAL CONSIDERATIONS

Models used in risk assessment must adequately reflect reality. Thus, before predictive models are used in exposure assessment, their appropriateness to that exposure assessment and overall reliability should be assessed. Users of models must be aware of the predictive limits of models, both in terms of the range of conditions that a model's interpolation region encompasses (Baranyi et al., 1996) and the variables that the model considers. Completeness error arises in model predictions when the model does not explicitly consider the effect of factors in a food that will affect the growth response of the microorganism modelled. The models referred to in Table A3.1 were developed to test different modelling strategies or, in the later published models, to include the effect of specific variables not included in earlier models. Ideally, a single model could encompass all the variables of relevance in all foods and is the ultimate aim of the scientific approach to predictive microbiology as the basis of a quantitative understanding of the microbial ecology of foods. However, creating such a model and scientific framework is time consuming. Alternatively, an iterative approach for development of product-oriented models i.e. based on observations in a system closely related to the food of interest, may satisfy the current technological needs of the food industry (Dalgaard, 1997; Dalgaard, Mejlholm and Huss, 1997).

Where completely appropriate models are not available, the limitations of the models should be documented and the implications of those limitations discussed as sources of uncertainty.

This section will consider assessment of model performance and limits. The discussion will be presented under the following headings:

- limits to application (i.e. interpolation or extrapolation);
- sources of variability and uncertainty; and
- performance evaluation.

### A3.3 INTERPOLATION OR EXTRAPOLATION

No predictive models currently in use have a sound basis in theory, i.e. they are empirical descriptions and summaries of observations. A simple rule of modelling is that models without theoretical bases cannot be used reliably to make predictions by extrapolation, but only by interpolation. Interpolation relates to prediction made “between” the observations that the model is based on, while extrapolation is when predictions are made for conditions outside the range of those studied in the development of the models. A common interpretation of the interpolation region is that any combination of variables (e.g. temperature, water activity, pH, phenol, nitrite, etc.) that falls within the respective ranges of variables tested in the development of the model is within the interpolation region.

Certainly, microbial growth or death in a food cannot be predicted reliably when the conditions are *outside* the range of any individual factor tested in the model. However, the interpolation region is usually smaller than the simple interpretation suggested above. Few models are based on full factorial experimental designs. Unfortunately, the regions with fewest observations are usually those at the extremes of the ranges, where growth is slowest or may not occur at all due to the interaction of inhibitory factors (this is considered further in the section below on growth/no-growth models). However, these regions are often of most relevance when modelling because they are the conditions normally used to extend the shelf-

life and safety of foods. As a result, users of models may inadvertently make predictions by extrapolation, particularly for conditions under which growth may be slow but of direct relevance in determining exposure to *L. monocytogenes* in RTE foods.

The determination of the true interpolation region, and the consequences of extrapolation, were discussed by Baranyi et al. (1996). Those authors concluded that models using a large number of parameters, e.g. higher order polynomial models, were more prone to unreliability resulting from inadvertent extrapolation because the predictions of the model often changed dramatically near the limits of the interpolation region.

Inadvertent extrapolation can also occur when using stochastic modelling techniques to describe effects of fluctuating temperature. Inadvertent extrapolation may also occur for other factors, but temperature is the factor most likely to fluctuate. Distributions can have infinitely long “tails”, so it is important that the tails of the distributions used to model temperatures are truncated to match the interpolation range of the predictive microbiology model used.

### A3.4 GROWTH/NO-GROWTH MODELS

Growth/no-growth models are a relatively new area of predictive microbiology. They aim to define the sets of combinations of factors that permit the growth of a modelled organism and those that do not. While there are absolute limits to growth of *L. monocytogenes* (see Table 3.1 in the main report) combinations of inhibitory factors can also prevent growth under milder conditions of each factor, a phenomenon widely employed in the food industry. These combinations of growth-preventing factors form a smooth surface in multi-dimensional space, or a smooth curve if one considers the interaction of two factors at a time as shown in Figure A3.2. There are relatively few growth/no-growth models currently available (Table A3.1). On the growth side of the interface, models can predict growth. On the no-growth side of the boundary, death occurs. Thus, growth/no-growth models provide additional information on the interpolation region of models.

### A3.5 SOURCES OF VARIABILITY AND UNCERTAINTY

Model predictions can never perfectly match observations. Each step in the model construction process introduces some error, as outlined below, and presented in order of the magnitude of their contribution to the overall error in the models predictions.

- **Homogeneity error** arises because either some foods are clearly not homogeneous, or, at the scale of a microorganism, foods of apparently uniform consistency may comprise many different microenvironments. Current predictive models do not account for this non-homogeneity of foods.
- **Completeness error** arises because the model is a simplification, i.e., in practice, not all relevant factors can be included in the model.
- **Model function error** is similar to completeness error, and arises mainly from the compromise made when using empirical models, i.e. that the model is only an approximation to reality.
- **Measurement error** originates from inaccuracy in the raw data used to estimate the parameters of a model, i.e. due to methodological limitations in our ability to measure accurately the environment and the microbial response.


- **Numerical procedure error** includes all errors that are the consequences of the numerical procedures used for model fitting and evaluation, some of which are methods of approximation only. Generally, numerical procedure errors are negligible in comparison with the other types of errors.

The error in the estimate of maximum specific growth rate (or doubling time) of an organism determined from measurement of growth in laboratory media is ~10% per independent variable. As a rule of thumb, each additional environmental factor (pH,  $a_w$ , etc.) adds at least another 10% relative error to the model, assuming that the interpolation region of the model is comparable to the whole growth region. (Models with a small interpolation region have smaller error). An example of the interaction of factors limiting the growth of *L. monocytogenes* and the use of a model to predict those interactions is presented in Figure A3.1. Thus, the best performance that might be expected from a kinetic model encompassing the effect of three environmental factors on growth rate is ~30%.

### A3.6 DISTRIBUTION OF RESPONSE TIMES

It is recognized that there is variation in the ecology of strains of *L. monocytogenes*. Begot, Lebert and Lebert (1997) reported variability in the growth rate responses of 58 strains. Peleg and Cole (1998) hypothesized that non-linear inactivation curves result from the natural variability that exists in microbial populations.

There has been lively discussion in the literature concerning the variability of bacterial growth rates. Using the limited amount of replicated published data concerning growth rate estimates under varying environmental conditions, Ratkowsky et al. (1991) concluded that growth rate responses became increasingly variable at slower growth rates, an observation confirmed by others (Fehlhaber and Krüger, 1998).



**Figure A3.1** An example of the interaction of factors limiting the growth of *Listeria monocytogenes* and the use of a model to predict those interactions. The smooth lines are predictions of a model for the probability of growth of *Listeria monocytogenes* L5 (Tienungoon et al., 2000). The symbols (filled circles: growth observed; open circles, no growth observed) are the data of George, Lund and Brocklehurst (1988) for the effect of temperature and pH on the growth of *L. monocytogenes* NCTC 10357. In the figure, the lines indicate the predicted limits of growth at various levels of confidence ( $P = 0.9$ : lower curve;  $P = 0.5$ : middle curve;  $P = 0.1$ : lower curve).

In the data presented by Ratkowsky et al. (1991) variance in the square root of growth rate ( $\text{var}\sqrt{\text{rate}}$ ) is constant, regardless of the magnitude of *rate*. Alber and Schaffner (1992) showed that for a strain of *Yersinia enterocolitica* (serotype 08), a logarithmic transformation of rate better “homogenizes” or “stabilizes” the variance. Dalgaard et al. (1994) reported that a transformation intermediate between the square root of rate and the logarithm of rate was required to normalize the variability in growth rate responses. Ratkowsky et al. (1996) reported similar observations, depending on the data used.

Ratkowsky (1992) presented the following general relationship between the variance in growth response times and the mean of those responses for a range of possible distribution types:

$$V = c\mu^n$$

where  $\mu$  is the mean of the probability distribution,  $V$  is the variance of the probability distribution,  $n$  is an integer exponent having values 0, 1, 2 or 3, corresponding to the normal, Poisson, Gamma (logarithm of rate) and Inverse Gaussian (square root of rate) distributions, respectively, and  $c$  is a constant.

It is important to characterize the variability in responses, and to recognize that those responses are not normally distributed if that information is to be used within stochastic models for risk assessment.

### A3.7 EVALUATION OF MODEL PERFORMANCE

A number of authors (Buchanan and Phillips, 1990; Wijtzes et al., 1993; George, Richardson and Peck, 1996; Fernandez, George and Peck, 1997; Walls and Scott, 1997; McClure et al., 1997; te Giffel and Zwietering, 1999) have evaluated the reliability of predictive microbiology models for *L. monocytogenes* growth rate and many have concluded that the models perform satisfactorily. However, in most of those assessments, no objective criterion for “satisfactory” was given.

Evaluation of model performance typically involves the comparison of model predictions to analogous observations not used to develop the model. Various measures have been used. Wijtzes et al. (1993) plotted literature values for the generation time of *L. monocytogenes* against the corresponding predictions of a model derived from studies in laboratory broth. From this plot, predictions that would be unsafe in practice could be visualized readily, and the overall reliability of the model assessed. Duh and Schaffner (1993) developed predictive equations for *Listeria* growth rate based on measurements in brain-heart infusion broth. Complementary literature values for the growth of the organism in food were then added to the data set and regression analysis of the supplemented dataset performed. The close similarity in mean square error (MSE) and correlation coefficient ( $r^2$ ) values of the equations fitted to either data set were taken as an indication of the reliability of the models when applied to foods. Another measure of the accuracy of predictive equations was introduced by McClure, Zwietering and Roberts (1993), who compared their models on the basis of the sum of the squares of the differences of the natural logarithm of observed and predicted values. Zwietering et al. (1994) introduced the use of the *F*-ratio test. In this method the MSE of the models when assessed against data that are not used to generate the model was compared to the measurement error of the model itself, i.e. the model compared to the data used to generate it. If the measurement error is not significantly different from the prediction error,

the model is considered to be satisfactory. Giffel and Zwietering (1999) reviewed these measures in greater detail.

Two additional complementary measures of model performance can be used to assess the “validity” of models and are claimed to have the advantage of being readily interpretable (Ross, 1996), namely a bias factor and an accuracy factor.

The “bias factor” ( $B_f$ ) is a multiplicative factor by which the model, on average, over- or under-predicts the response time. Thus, a bias factor of 1.1 indicates not only that a growth model is “fail-dangerous” because it predicts longer generation times than are observed, but also that the predictions exceed the observations, on average, by 10%. Conversely, a bias factor less than unity indicates that a model is, in general, “fail-safe”, but a bias factor of 0.5 indicates a poor model that is overly conservative because it predicts generation times, on average, half of that actually observed. Perfect agreement between predictions and observations would lead to a bias factor of 1.

The “accuracy factor” ( $A_f$ ) is also a simple multiplicative factor indicating the *spread* of observations about the model’s predictions. An accuracy factor of two, for example, indicates that the prediction is, on average, a factor of two different from the observed value, i.e. either half as large or twice as large. The bias and accuracy factors can equally well be used for any time-based response, e.g. lag time, time to an  $n$ -fold increase, death rate, D value, etc. Modifications to the factors were proposed by Baranyi, Pin and Ross (1999).

Ideally, predictive models would have  $A_f = B_f = 1$ , but, typically, the accuracy factor will increase by 0.10–0.15 for every variable in the model. Thus, an acceptable model that predicts the effect of temperature, pH and water activity on *Listeria* growth rate could be expected to have  $A_f = 1.3$ –1.5. Satisfactory  $B_f$  limits are more difficult to specify because limits of acceptability are related to the specific application of the model.  $B_f$  is a measure of the extent of under- or over-prediction of the observed response rates by the model. Thus, a bias factor of 1.1 indicates not only that a generation time model is “fail-dangerous” not only because it predicts longer generation times than are observed, but also because the observations exceed the predictions, on average, by 10% in terms of  $\log_{10}$  CFU. Conversely,  $B_f < 1$  indicates that a model is, in general, “fail-safe”. Note, however, that when applied to rate-based data,  $B_f > 1$  indicates the model *under*-predicts the observed rate, potentially leading to “fail-dangerous” predictions.

Armas, Wynne and Sutherland (1996) considered that  $B_f$  values in the range 0.6–3.99 were acceptable for the growth rates of pathogens and spoilage organisms when compared with independently published data. Giffel and Zwietering (1999) assessed the performance of many models for *L. monocytogenes* against seven datasets, and found Bias factors in the range 2–4, which they considered to be acceptable, allowing predictions of the order of magnitude of changes to be made.

Other workers have adopted higher standards. Dalgaard (2000) suggested that  $B_f$  values for successful validations of seafood *spoilage* models should be in the range 0.8–1.3. Ross (1999) considered that, for pathogens, less tolerance should be allowed for  $B_f > 1$  because that corresponds to under-predictions of the extent of growth and could lead to “fail-dangerous” predictions. Thus, Ross (1999) recommended that for models describing *pathogen* growth rate,  $B_f$  in the range 0.9–1.05 could be considered good, in the range 0.7–0.9 or 1.06–1.15 considered acceptable, and less than ~0.7 or greater than 1.15 considered unacceptable.

### A3.8 SPECIFIC MODELS VERSUS GENERAL MODELS

The results of te Giffel and Zwietering (1999) and Ross (1999) showed that model performance is dependent on the data used to assess them. Differences in the performance of individual models were observed when the test datasets were disaggregated into food groups, or into ranges of growth rates. Some of these differences stem from the quality of the data used to assess the models, and the shortcomings of assessing models against data derived from the published literature have been commented on in several studies (Sutherland, Bayliss and Roberts, 1994; Ross, 1996; Walls and Scott, 1997; te Giffel and Zwietering, 1999). A second reason for poor performance may stem from completeness error. While te Giffel and Zwietering (1999) endorsed the performance of general models, Dalgaard (1997) and Dalgaard, Mejlholm and Huss (1997) proposed that strategies for model development based on observations in a system closely related to the food of interest will provide better performance for that specific product.

### A3.9 PRACTICAL MICROBIAL ECOLOGY MODELLING IN RISK ASSESSMENTS

#### A3.9.1 Temperature distributions

Foods are rarely held under completely controlled temperature during their entire shelf-life. A common technique is to model the average temperature, based on temperature records obtained from surveys. The growth rate response of bacteria to temperature is complex and is not directly proportional to temperature. As noted by Cassin et al. (1998) the question arises whether the use of the average temperature over a time interval systematically biases the estimate of growth. This issue was addressed by Ross (1999) who used 246 temperature histories obtained using electronic temperature data-loggers for meat processing, transport and storage in Australia. Typically, the time interval between temperature recordings was a few minutes long.

Three methods were used to calculate the amount of microbial growth for each temperature history. In the first, the estimate of growth was based on the average temperature of all the temperatures recorded over the monitoring period. In addition, estimates were also generated for the *worst* case, i.e. based on the highest temperature recorded in each temperature record. The average and highest temperature values were substituted into models to predict the number of generations of pseudomonads and *E. coli* for each temperature history, respectively, by the two approaches. In the third method, the growth was determined using “time temperature function integration”. For each time interval in the temperature history the growth rate of both pseudomonads and *E. coli* at the beginning and end of each time interval was calculated. The average of those growth rates was substituted into predictive models to calculate the number of generations over each recording interval, and the calculated number of generations for each time interval added to estimate the growth (i.e. number of generations) over the *entire* time monitored for each of the 246 temperature histories used.

In all methods, any temperature outside the ranges specified for each model were calculated to correspond to no growth, whether based on the average temperature over the interval, or full time-temperature integration.

For each organism-sector combination, the histograms of the distributions predicted by each method were plotted on a single graph to enable direct comparison of the effect of the three calculation methods. Representative plots of pseudomonad growth are shown in Figure A3.2.

The relationship between specific sets of predictions is lost in the preparation and presentation of the frequency distribution graph.

Ross (1999) showed mathematically that the average rate of growth at two temperatures *in the sub-optimal temperature region* is always greater than or equal to the growth rate at the average of two temperatures and that the difference between the two calculation methods is a function of the magnitude of the difference between the two temperatures. Using the dataset described the results indicated that *in practice* the difference between the two estimation methods is typically of the order of -0.1 to 0.2 log<sub>10</sub> CFU, presumably because in most cases the range of temperatures experienced is small. This is a very small difference, particularly bearing in mind that the limits of accuracy of current microbial enumeration methods is approximately 0.3 log<sub>10</sub> CFU (Jarvis, 1989).

However, there are certain situations and temperature ranges in which differences due to estimation method become more pronounced. If the temperatures experienced transcend growth boundary values, e.g. maximum or minimum temperatures for growth, estimates of the predicted growth by the two methods can differ significantly and lead to different frequency distributions of predicted growth. They are unlikely to be important for prediction of the growth of *Listeria monocytogenes* in RTE foods, however, because the lower (= 0°C) or upper (= 45°C) temperature thresholds for *L. monocytogenes* are unlikely to be experienced in normal refrigerated storage.

Thus, the results of that study (Ross, 1999) suggest that the use of the average temperature approach can provide a reasonable prediction of the extent of the growth of *L. monocytogenes* under real conditions of storage and distribution.

### **A3.9.2 Upper and lower limits**

When distributions of temperatures are defined, they should reflect reality, i.e. the distributions should be truncated at realistic values. Similarly, when the range of temperature defined in the exposure model exceeds the minimum and optimum or maximum temperatures for growth of the organism, the growth model used must model the response of *L. monocytogenes*, i.e. the decline in growth rate as temperatures increase above that optimal for growth rate; and the cessation of growth at temperatures above or below the limits for growth.

Further pitfalls may occur in the use of unbounded temperature distributions. If the temperature distribution exceeds the range of the predictive model, nonsense predictions can occur, and may not be revealed by the simulation software used. While the effects might be subtle, they are likely to increase the range of uncertainty in the final model prediction.

### **A3.9.3 Lag time response**

Microbial lag time is dependent both on the environment and its effect on growth rate, and the amount of “work” the cell has to do before it can initiate growth. This has presented problems for modellers, because models are developed under sets of constant conditions and

it has been difficult to relate one set of conditions to another. The use of the relative lag time (RLT) concept, and RLT distributions, provides a way to overcome these problems in developing exposure assessments.

Ross (1999), Mellefont, McMeekin and Ross (2003) and Mellefont and Ross (2003) combined lag time data from experiments deliberately intended to induce long lag times with the published observations of other workers to investigate the distribution of lag times that are observed. When the lag time is expressed as an equivalent number of generation times of the organism in the same environment, i.e. lag time divided by generation time, or RLT, the distribution of RLTs observed has a sharp peak in the range 3–6. Augustin and Carlier (2000) also collated relative lag time distributions. The results are shown in Figures A3.3 and suggest that in many situations there is a practical upper limit to the lag time duration.

The number of generations of growth is predicted from the time and environmental conditions. The relative lag time is sampled from the RLT distribution and deducted from the predicted growth. If the predicted generations of growth do not exceed the lag time, no growth is predicted. If it does, the growth predicted to have occurred is given by the difference between the predicted generations of growth less the RLT.


#### **A3.9.4 Jameson Effect**

There has been very little work done to include in predictive models factors that contribute to the “Jameson effect” (Stephens et al., 1997), i.e. the suppression of growth of all microorganisms in the food by high *total* microbial loads. In some products, this effect may greatly reduce the health risk from *L. monocytogenes* predicted on the basis of models currently available. Example 4 in this report (cold-smoked fish) introduces a method for inclusion of the Jameson effect in exposure assessment modelling. It models the increase in spoilage or other microorganisms, or both, on the product simultaneously with the growth of *L. monocytogenes*. If the predicted growth of other microbiota is predicted to exceed  $10^9$  CFU/g, the predicted growth of *L. monocytogenes* is modified accordingly. Full details are given in the example.


#### **A3.9.5 Physiological state of cells**

Environmental and physiological factors during food processing or present in foods are reported to affect the infectivity or virulence, or both, of *L. monocytogenes* (Buchanan et al., 1994; Zemser and Martin, 1998). These have been reviewed (Rees et al., 1995; Archer, 1996; Rowan, 1999), and also specifically in relation to *L. monocytogenes* in foods (Lou and Yousef, 1999). Conversely other workers (Conte et al., 1994; Gahan and Hill, 1999) have found little effect of environmental conditions on virulence.

Harsh environments in foods that stress the microbial cell produce a response that makes the cell more resistant to subsequent stressful or potentially lethal conditions, extending the survival of the cell under those conditions. Cells that are in stationary phase will also have increased tolerance to potentially lethal environments. This phenomenon has been suggested as increasing the chance that pathogenic bacteria, including *L. monocytogenes*, will survive passage through the acid environment of the stomach, thereby effectively increasing their virulence.



**Figures A3.2** Graphs showing the distribution of the predicted number of generations of growth of pseudomonads during (a) transport from retail to home, (b) foodservice, and (c) domestic storage (home refrigerators) The heavy dashed line represents the predictions based on the average temperature; the solid line represents predictions based on time-temperature function integration; and the light dotted line represents predictions based on the maximum temperature recorded.



**Figure A3.3** Distribution of relative lag times reported for *Listeria monocytogenes*: (a) grown in broth under laboratory conditions (collated in Ross, 1999); (b) in foods (collated in Ross, 1999); and (c) collated by Augustin and Carlier (2000) for all sources and plotted as  $\ln(\text{RLT})$ . In figures (a) and (b), the dotted lines represent the cumulative frequency.

While these effects on virulence, and the chance of infection, are recognized, they are not well characterized and may be specific to strains and conditions (Buncic and Avery, 1996; Buncic, Avery and Rogers, 1996).

There is also uncertainty about the virulence of foodborne strains of *L. monocytogenes*. Other than that most cases of foodborne illness have been associated with serotypes 4a and 1/2a/b, specifically virulent strains cannot yet be differentiated. Notermans et al. (1998) studied the infectivity of more than 20 foodborne strains of *L. monocytogenes*, using a mouse bioassay and chick embryo test. Despite observing differences in virulence between strains, they concluded that almost all *L. monocytogenes* serovars present in foods have clear virulent properties, and should be considered potentially pathogenic, a view shared by McLauchlin (1996). Conversely, some exposure assessments (Farber, Ross and Harwig, 1996; Bemrah et al., 1998) have assumed that only 1–10% of foodborne *L. monocytogenes* are pathogenic.

The above issues are also discussed in detail in Part 2 of the main report.

### **A3.10 MODELLING CONTAMINATION AND RE-CONTAMINATION**

There is little data available upon which to enable cross-contamination, or its effects, to be modelled quantitatively. There are a number of variables that might be considered:

- (i) If contact with contaminated material occurs, how often does cross-contamination result?
- (ii) At what point in the food chain does it occur?
- (iii) What is the potential for growth on fomites, such as cutting equipment?
- (iv) How much material is transferred and does the nature of the source affect the amount transferred?

#### **A3.10.1 Source and amount of material transferred**

FAO/WHO (2002) cites the results of Zhao et al. (1998), who developed a model system to enumerate bacteria transferred during common food preparation practices. Zhao et al. (1998) found that chicken meat and skin inoculated with  $10^6$  bacteria transferred  $10^5$  to a chopping board and hands and then  $10^3$ – $10^4$  to vegetables chopped on the unclean board. It should be noted that chicken skin is likely to be wet and this might facilitate the transfer of bacteria, suspended in a surface film of moisture, compared with what might be transferred from RTE foods, which are often “drier” to the touch, e.g. cheeses, processed meats, smoked fish, etc.

#### **A3.10.2 Potential for growth**

In risk assessments, because of the assumption in some dose-response models that the risk of infection is directly proportional to dose for the low- to medium-dose range, the estimate of the microbiological risk to a population is largely governed by the estimate of the total numbers of the pathogen in the food supply. How that number of pathogens is distributed among individual packages of foods has less effect on the risk estimate. Accordingly, simple transfer of contamination from one unit of food to another will not affect the risk estimate, unless that transfer is subsequently accompanied by growth (i.e. multiplication) of the pathogen on a contact surface that contaminates uncontaminated material or growth in the (now) contaminated product itself.

In many RTE products, *L. monocytogenes* cells that contaminate the product will remain effectively immobilized at the site of contamination unless there is free liquid in the package, or other modes of transfer, to transport them to other parts of the package. Consequently, *L. monocytogenes* in some foods may be highly localized and exist as discrete pockets of contamination. This may limit the potential for growth of the organism, as nutrients are utilized and wastes accumulate at that site of contamination. Transfer to a new environment provides new potential for growth and potential increase in risk.

Grau (1993) traced the flow of *L. monocytogenes* through meat processing plants and found it to be transferred to many sites, such as trolleys, door handles and the surface packaging of finished products by contact contamination and cross-contamination. In these cases, paucity of nutrients and moisture may inhibit the growth of the organism, and limit the numbers of *L. monocytogenes* transferred to any product.

*L. monocytogenes* is known to colonize processing plants and, in particular, wet areas in plants. In these areas, if organic matter is present, growth can be expected to occur given sufficient time. Investigations in the United States of America suggest that listeriosis outbreaks often arise when virulent strains “colonize” a production line (Tompkin, 2002). Sites of colonization include hard-to-clean processing equipment. Hollow rollers on production lines are also known to deteriorate and crack, allowing water, nutrients and bacteria to colonize the interior. These niches are very difficult to clean, and provide a reservoir of pathogenic contaminants.

On equipment that is in direct contact with food and becomes fouled with food, growth would be expected to occur. The amount of growth that could occur would be determined by the product composition, the temperature of that part of the plant, and the time before the contamination was removed by cleaning. As an example, if the processing line were operating at 15°C and slicing a processed meat product (e.g. pH 6.2,  $a_w$  0.975, 100 ppm nitrite), *L. monocytogenes* growing in a residual material on contact surfaces could double in numbers approximately every 5 hours.

### **A3.10.3 Point in food chain at which contamination occurs**

As stated above, there is no increase in risk as a consequence of cross-contamination unless there is increased potential for microbial growth as a result. The amount of increase will depend on the product, its storage conditions, and the time between the contamination event and consumption. If the integrity of the chill chain between the point of production and consumption were uniform, the potential consequences of contamination would be expected to be greater for contamination at the point of production than at the point of retail sale or in the consumer’s home. This is because of the increased time available for growth to high numbers before consumption.

### **A3.10.4 Likelihood of transfer**

Even if uncontaminated material comes into contact with contaminated material, the probability of cross-contamination is not absolute, but would be expected to depend on the concentration of pathogens, and their distribution on, or in, the contaminated material.

### A3.11 RELATIVE RATE FUNCTIONS

Growth rate modelling can be simplified enormously by using relative rate functions, particularly when combined with square root-type models, or cardinal parameter models. The simple square root model (Ratkowsky et al., 1982) describes the effect of temperature on the growth rate of almost all bacteria. The square root model is:

$$\sqrt{\mu} = b(T - T_{min})$$

where  $\mu$  is the rate of growth  
 $T$  is temperature (°C)  
 $b$  is a constant to be fitted related to the maximum growth rate of the organism  
 $T_{min}$  is the temperature at which the growth rate is predicted to be zero.

It should be noted that  $T_{min}$  is a notional temperature, and is usually several degrees below the minimum temperature at which growth is observed to occur. It should also be noted that the simple square root model above applies only to the sub-optimal temperature region for growth, up to ~35–37°C for *L. monocytogenes*.

If:

- temperature is the only factor affecting the growth rate of a bacterium in a food that varies during the storage and distribution of the product (i.e. if pH, water activity, etc., are constant),
- $T_{min}$  is known for the organism, and
- the growth rate in a product of interest is known at one temperature,

then the growth rate of the organism in that product at any other temperature can be derived using the following relationship, based on the simple square root model (McMeekin et al., 1993):

$$\mu_T = \mu_{REF} * \frac{(T - T_{min})}{(T_{REF} - T_{min})}$$

where  $\mu_{REF}$  is the known growth rate at some temperature  $T_{REF}$ ,  
 $\mu_T$  is the unknown growth rate at some temperature  $T$ ,  
and the other parameters are as previously defined.

For example, FDA/FSIS (2001) collated data for the growth rate at 5°C of *L. monocytogenes* in many RTE food products. Growth at a temperature other than 5°C was calculated using the relationship  $\sqrt{\mu_T} = \sqrt{\mu_5}(T - T_{min})(5 - T_{min})^{-1}$ . This approach has been adopted in several of the exposure assessments in Part 4 of the main report.

The use of the relative rate function is a simplification. As conditions become less favourable for microbial growth, e.g. due to decreased water activity or increased acidity, the difference between  $T_{min}$  and the minimum temperature at which growth is possible increases. This was discussed above in relation to models for growth limits under multiple hurdles to growth (e.g. Tienungoon et al., 2000). Bajard et al. (1996) suggested that the simple square root model does not describe the growth rate response to temperature of *L. monocytogenes* as well as it does for other organisms. Nonetheless, in the context of the other sources of

uncertainty that arise in microbial risk assessments, these are considered to be relatively minor deficiencies.

### A3.12 PREDICTIVE MICROBIOLOGY MODELS

A summary of some currently available predictive models for *L. monocytogenes* is presented in Table A3.1.

**Table A3.1** Summary of predictive models available for the growth, survival and inactivation of *Listeria monocytogenes* in foods.

1.	2.	3. (°C)	4. (a <sub>w</sub> or %)	5. pH	6.	7. (µg/ml)	8.	9.	10.
<b>GROWTH</b>									
Y	broth	5–37	0.5–4.5%	4.5–7.5	–	0–1000	aerobic/ anaerobic	N	[1]
Y	food	3–35	2–8%	4.5–7.5	–	–	aerobic	Y	[2]
Y	broth	5–35	0.5–8%	4.6–7.4	–	–	aerobic	Y	[3]
N	broth	4–20 1–20	–	4.5–7.0 4.3–7.2	acetic 0–10 000 lactic 0–20 000	–	aerobic	Y	[4]
Y	broth	9	1.0–4.0%	5.5–6.5	lactic 0–0.6%; acetic 0–0.6%	70	aerobic	Y	[5]
Y	broth	2–46	–	–	–	–	aerobic	N	[6]
Y	broth	4–20	0.5–8%	4.5–7.0	–	–	CO <sub>2</sub> : 0–100%, balance N <sub>2</sub>	Y	[7]
Y	food	3, 7, 11	0.5	–	–	–	Air: 0.03% CO <sub>2</sub> , 78.03% N <sub>2</sub> , 20.99% O <sub>2</sub> ; Modified atmosphere #1: 76% CO <sub>2</sub> , 13.3% N <sub>2</sub> , 10.7% O <sub>2</sub> ; Modified atmosphere #2: 80% CO <sub>2</sub> , 20% N <sub>2</sub>	N	[8]
Y	meat broth	4–30	0.992–0.960 (a <sub>w</sub> )	5.4–7	–	–	aerobic	Y	[9]
N	broth	20–35	2–10%	4–8.5	–	–	–	N	[10]
N	broth	1.0–35	0.5–11.5%	4.0–7.2	–	0–200	–	N	[11]
Y	lean beef and fatty beef tissue	0–30.6	–	5.46–6.98	–	–	aerobic	Y	[12]
Y	lean beef fat beef	0–43 0–31	~0.99	5.6–6.7	–	–	aerobic	Y	[13]
Y	broth	3–37	0.5–13%	4.2–7.3	lactic 0–450 mM	–	aerobic	Y	[14]
Y	broth	4–37	0.5–13%	5.6–7	lactic 0, 200 mM	–	aerobic	Y	[15]
N	roast beef	-1.5 & 3	–	6.1	–	–	vac. pack and saturated CO <sub>2</sub>	N	[16]
N	broth	5–30	0.5–8%	4.6–7.4	–	0–400	–	N	[17]
Y	broth	5–35	0.95–0.997	#1 4.6–6.7 #2 4.6–7.4	–	–	–	Y	[18]
Y	broth	5–37	0.5 & 4.5	6.0 & 7.5	–	0–1000	aerobic and anaerobic	N	[19]

1.	2.	3. (°C)	4. (a <sub>w</sub> or %)	5. pH	6.	7. (µg/ml)	8.	9.	10.
<i>Dynamic growth</i>									
Y	fluid whole milk		4, 6, 8, 10, 15, 20, 25, 30, 35	–	–	–	–	Y	[20]
<i>Repair of heat injury</i>									
Y	broth	4–43	0.5–10.0	4.2–9.6	–	–	–	N	[21]
<b>SURVIVAL/GROWTH LIMITS/GROWTH INITIATION</b>									
<i>Probability of growth initiation in defined period of time</i>									
N	broth	4–30	0.5–12.5	>5.9	–	See Note (1)	–	N	[22]
<i>Survival and ongrowth</i>									
		5, 10, 30	0–18	4.19–4.83	–	–	aerobic	N	[23]
<i>Growth limits</i>									
	broth	3.1–35.8 3.1–36.4	0.5–13% 0.5–13%	3.9–7.3 3.9–7.7	lactic 0–500 lactic 0–450	–	aerobic	Y	[24]
<i>Effect of heat stress</i>									
Y	broth	53–60	–	–	–	–	Stationary phase cells	N	[25]
<b>INACTIVATION</b>									
<i>Thermal</i>									
Y	milk (bovine)	60.5–69.5 for 3–60 secs (HTST pasteurization process)	–	–	–	–	–	Y	[26]
Y	food	55, 60, 65	–	5, 6, 7	–	–	milkfat 0, 2.5, 5%	Y	[27]
Y	food	55–65	0–6%	4–8	–	–	sodiumpyrophosphate 0–0.3%	Y	[28]
Y	food (infant formula)	55, 60, 65	0, 2, 4%	5, 6, 7	–	–	physiological states (lag, exponential, stationary) of test cultures	Y	[29] using data from [41]
<i>Heating rate and thermal inactivation</i>									
Y	broth	50–64	–	–	–	–	Sodiumpyrophosphate 0–0.3%	N	[30]
<i>Heat resistance</i>									
Y	broth	50, 60, 65	–	–	–	–	Physiological state of cells (end of log phase cells; heat shocked cells; cells resistant to prolonged heat)	Y	[31]
Y	buffer	50, 55, 60	–	–	–	–	–	Y	[32]
Y	broth	30, 10, 5	0–18%	4.19–4.83	–	–	–	Y	[33]
<i>Non-thermal</i>									
N	broth	4 to 42	0.5–19%	3.3–7.3	lactic 0–2%	0–200	–	N	[34]

1.	2.	3. (°C)	4. (a <sub>w</sub> or %)	5. pH	6.	7. (µg/ml)	8.	9.	10.
N	broth	5 to 42	0.5–19%	3.3–7.4	lactic 0–2%	0–200	O <sub>2</sub> levels reduced, N <sub>2</sub> flushed vessels	N	[35]
Y	broth	28	–	4–7	lactic 0–18%; acetic 0–12%	–	–	Y	[36]
Y	broth	4, 19, 28	–	3–4.5	acetic: 0–2.0%; ascorbic: 0–2.0%	–	aerobic	N	[37]
Y	broth	4–42	0.5–19%	–	lactic 0–1%	0–200	–	Y	[38]
<b>COMBINED</b>									
<i>Growth survival death</i>									
Y	broth	4–12	2–4%	6.2	–	–	phenol: 5, 12.5, 20 ppm	Y	[39]
<i>Biotic interactions</i>									
Y	broth	10	2%	5–5.8	0–5 mM protonated lactic acid	–	<i>Lactococcus lactis</i> (non-nisin producing)	Y	[40]

KEY TO COLUMNS: (1) Model given? Y = Yes; N = No. (2) Medium. (3) Temperature (°C). (4) Water activity (a<sub>w</sub>) or salt (NaCl) percentage. (5) pH. (6) Organic acids. (7) Nitrite, expressed as µg/ml (= ppm). (8) Other. (9) Validation data? Y = Yes; N = No. (10) Source (see below).

NOTES: (1) methyl paraben 0–2%; sodium propionate 0.3%; sodium benzoate 0.1%; potassium sorbate 0.3%; inoculum size 0.01–100 000 CFU/ml; *Listeria* spp. (*L. monocytogenes*, *L. innocua*, *L. seeligeri* and *L. ivanovii*).

SOURCES: [1] Buchanan and Phillips, 1990. [2] Murphy, Rae and Harrington, 1996. [3] Wijtzes et al., 1993. [4] George, Richardson and Peck, 1996. [5] Nerbrink et al., 1999. [6] Duh and Schaffner, 1993. [7] Fernandez, George and Peck, 1997. [8] Zhao, Wells and Marshall, 1992. [9] Lebert, Bégot and Lebert, 1998. [10] McClure, Roberts and Otto Oguru, 1989. [11] McClure et al., 1997. [12] Grau and Vanderlinde, 1993. [13] Grau and Vanderlinde, 1992. [14] Tienungoon, 1998. [15] Ross, 1993. [16] Hudson, Mott and Penny, 1994. [17] McClure, Kelly and Roberts, 1991. [18] McClure, Zwietering and Roberts, 1993. [19] Buchanan, Stahl and Whiting, 1989. [20] Alavi et al., 1999. [21] Chawla, Chen and Donnelly, 1996. [22] Razavilar and Genigeorgis, 1998. [23] Cole, Jones and Holyoak, 1990. [24] Tienungoon et al., 2000. [25] Breand et al., 1998. [26] Piyasena, Liou and McKellar, 1998. [27] Chabra et al., 1999. [28] Juneja and Eblen, 1999. [29] Xiong et al., 1999. [30] Stephens, Cole and Jones, 1994. [31] Augustin, Carlier and Rozier, 1998. [32] Linton et al., 1995. [33] Cole, Jones and Holyoak, 1990. [34] Buchanan and Golden, 1995. [35] Buchanan, Golden and Phillips, 1997. [36] Buchanan and Golden, 1995. [37] Golden, Buchanan and Whiting, 1995. [38] Buchanan et al., 1994. [39] Farber, Cai and Ross, 1996. [39] Membre, Thurette and Cateau, 1997. [40] Breidt and Fleming, 1998. [41] Linton et al., 1996.

**A3.13 REFERENCES CITED IN APPENDIX 3**

- Alavi, S.H., Puri, V.M., Knabel, S.J., Mohtar, R.H. & Whiting, R.C. 1999. Development and validation of a dynamic growth model for *Listeria monocytogenes* in fluid whole milk. *Journal of Food Protection*, **62**: 170–176.
- Alber, S.A. & Schaffner, D.W. 1992. Evaluation of data transformations used with the Square Root and Schoolfield models for predicting bacterial growth rate. *Applied and Environmental Microbiology*, **58**: 3337–3342.
- Archer, D.L. 1996. *Listeria monocytogenes* – the science and policy. *Food Control*, **7**: 181–182.
- Armas, A.D., Wynne, A. & Sutherland, J.P. 1996. Validation of predictive models using independently published data. Poster/Abstract. 2nd International Conference of Predictive Microbiology. Hobart, Tasmania, 18–22 February 1996.
- Augustin, J.C. & Carlier, V. 2000. Mathematical modelling of the growth rate and lag time for *Listeria monocytogenes*. *International Journal of Food Microbiology*, **56**: 29–51.
- Augustin, J.C., Carlier, V. & Rozier, J. 1998. Mathematical modelling of the heat resistance of *Listeria monocytogenes*. *Journal of Applied Microbiology*, **84**: 185–191.
- Bajard, S., Rosso, L., Fardel, G. & Flandrois, J.P. 1996. The particular behaviour of *Listeria monocytogenes* under sub-optimal conditions. *International Journal of Food Microbiology*, **29**: 201–211.
- Baranyi, J., Ross, T., McMeekin, T.A. & Roberts, T.A. 1996. Effects of parameterization on the performance of empirical models used in ‘Predictive Microbiology’. *Food Microbiology*, **13**: 83–91.
- Baranyi, J., Pin, C. & Ross, T. 1999. Validating and comparing predictive models. *International Journal of Food Microbiology*, **48**: 159–166.
- Begot, C., Lebert, I. & Lebert, A. 1997. Variability of the response of 66 *Listeria monocytogenes* and *Listeria innocua* strains to different growth conditions. *Food Microbiology*, **14**: 403–412.
- Bemrah, N., Sana, M., Cassin, M.H., Griffiths, M.W. & Cerf, O. 1998. Quantitative risk assessment of human listeriosis from consumption of soft cheese made from raw milk. *Preventative Veterinary Medicine*, **37**: 129–145.
- Breand, S., Farde, G., Flandrois, J.P., Rosso, L. & Tomassone, R. 1998. Model of the influence of time and mild temperature on *Listeria monocytogenes* non-linear survival curves. *International Journal of Food Microbiology*, **40**: 185–195.
- Breidt, F. & Fleming, H.P. 1998. Modeling of the competitive growth of *Listeria monocytogenes* and *Lactococcus lactis* in vegetable broth. *Applied and Environmental Microbiology*, **64**: 3159–3165.
- Buchanan, R.L. & Golden, M.H. 1995. Model for the non-thermal inactivation of *Listeria monocytogenes* in a reduced oxygen environment. *Food Microbiology*, **12**: 203–212.
- Buchanan, R.L. & Phillips, J.G. 1990. Response surface model for predicting the effects of temperature, pH, sodium chloride content, sodium nitrite concentration and atmosphere on the growth of *Listeria monocytogenes*. *Journal of Food Protection*, **53**: 370–376.
- Buchanan, R.L. & Whiting, R.C. 1997. Risk assessment - A means for linking HACCP plans and public health. *Journal of Food Protection*, **61**: 1531–1534.
- Buchanan, R.L., Golden, M.H. & Phillips, J.G. 1997. Expanded models for the non-thermal inactivation of *Listeria monocytogenes*. *Journal of Applied Microbiology*, **82**: 567–577.
- Buchanan, R.L., Golden, M.H., Whiting, R.C., Phillips, J.G. & Smith, J.L. 1994. Non-thermal inactivation models for *Listeria monocytogenes*. *Journal of Food Science*, **59**: 179–188.

- Buchanan, R.L., Stahl, H.G. & Whiting, R.C. 1989. Effects and interactions of temperature, pH, atmosphere, sodium chloride and sodium nitrite on the growth of *Listeria monocytogenes*. *Journal of Food Protection*, **52**: 884–851.
- Buncic, S. & Avery, S.M. 1996. Relationship between variations in pathogenicity and lag phase at 37°C of *Listeria monocytogenes* previously stored at 4°C. *Letters in Applied Microbiology*, **23**: 18–22.
- Buncic, S., Avery, S.M. & Rogers, A.R. 1996. Listeriolysin O production and pathogenicity of non-growing *Listeria monocytogenes* stored at refrigeration temperature. *International Journal of Food Microbiology*, **31**: 133–147.
- Cassin, M.H., Lammerding, A.M., Todd, E.C.D., Ross, W. & McColl, R.S. 1998. Quantitative risk assessment for *Escherichia coli* O157:H7 in ground beef hamburgers. *International Journal of Food Microbiology*, **41**: 21–44.
- Chabra, A.T., Carter, W.H., Linton, R.H. & Cousin, M.A. 1999. A predictive model to determine the effects of pH, milkfat, and temperature on thermal inactivation of *Listeria monocytogenes*. *Journal of Food Protection*, **62**: 1143–1149.
- Chawla, C.S., Chen, H. & Donnelly, C.W. 1996. Mathematically modeling the repair of heat-injured *Listeria monocytogenes* as affected by temperature, pH, and salt concentration. *International Journal of Food Microbiology*, **30**: 231–242.
- Cole, M.B., Davies, K.W., Munro, G., Holyoak, C.D. & Kilsby, D.C. 1993. A vitalistic model to describe the thermal inactivation of *Listeria monocytogenes*. *Journal of Industrial Microbiology*, **12**: 232–239.
- Cole, M.B., Jones, M.V. & Holyoak, C. 1990. The effect of pH, salt concentration and temperature on the survival and growth of *Listeria monocytogenes*. *Journal of Applied Bacteriology*, **69**: 63–72.
- Conte, M.P., Longhi, C., Petrone, G., Polidoro, M., Valenti, P. & Seganti, L. 1994. *Listeria monocytogenes* infection of caco-2 cells - role of growth temperature. *Research Microbiology*, **145**: 677–682.
- Dalgaard, P. 1997. Predictive microbiological modelling and seafood quality. pp. 431–443, in: J. Luten, T. Børresen and J. Oehlenschläger (eds). *Seafood from Producer to Consumer, Integrated Approach to Quality*. Amsterdam, The Netherlands: Elsevier.
- Dalgaard, P. 2000. Fresh and lightly preserved seafood. pp. 110–139, in: C.M.D. Man and A.A. Jones (eds). *Shelf life evaluation of foods*. 2nd edition. Maryland, USA: Aspen Publishing.
- Dalgaard, P., Ross, T., Kamperman, L., Neumeyer, K. & McMeekin, T.A. 1994. Estimation of bacterial growth rates from turbidimetric and viable count data. *International Journal of Food Microbiology*, **23**: 391–404.
- Dalgaard, P., Mejlholm, O. & Huss, H.H. 1997. Application of an iterative approach for development of a microbial model predicting the shelf-life of packed fish. *International Journal of Food Microbiology*, **38**: 169–179.
- Duh, Y.-H. & Schaffner, D.W. 1993. Modeling the effect of temperature on the growth rate and lag time of *Listeria innocua* and *Listeria monocytogenes*. *Journal of Food Protection*, **56**: 205–210.
- FAO/WHO. 2002. Risk assessments of *Salmonella* in eggs and broiler chickens. *FAO/WHO Microbiological Risk Assessment Series*, No. 2. 300p.
- Farber, J.M. 1986. Predictive modeling of food deterioration and safety. pp. 57–90, in: M.D. Pierson and N.J. Stern (eds). *Foodborne Microorganisms and their Toxins*. New York NY: Marcel Dekker.

- Farber, J.M., Cai, Y. & Ross, W.H. 1996. Predictive modelling of the growth of *Listeria monocytogenes* in CO<sub>2</sub> environments. *International Journal of Food Microbiology*, **32**: 133–144.
- Farber, J.M., Ross, W.H. & Harwig, J. 1996. Health risk assessment of *Listeria monocytogenes* in Canada. *International Journal of Food Microbiology*, **30**: 145–156.
- FDA/FSIS [U.S. Food and Drug Administration/Food Safety and Inspection Agency (USDA)]. 2001. Draft Assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. Center for Food Safety and Applied Nutrition (FDA) and Food Safety Inspection Service (USDA) (Available at: [www.foodsafety.gov/~dms/lmrisk.html](http://www.foodsafety.gov/~dms/lmrisk.html)). [Report published September 2003 as: Quantitative assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. Available at: [www.foodsafety.gov/~dms/lmr2-toc.html](http://www.foodsafety.gov/~dms/lmr2-toc.html)].
- Fehlhaber, K. & Krüger, G. 1998. The study of *Salmonella enteritidis* growth kinetics using rapid automated bacterial impedance technique. *Journal Applied Microbiology*, **84**: 945–949.
- Fernandez, P.S., George, S.M. & Peck, M.W. 1997. Predictive model of the effect of CO<sub>2</sub>, pH, temperature and NaCl on the growth of *Listeria monocytogenes*. *International Journal of Food Microbiology*, **37**: 37–45.
- Gahan, C.G.M. & Hill, C. 1999. The relationship between acid stress responses and virulence in *Salmonella typhimurium* and *Listeria monocytogenes*. *International Journal of Food Microbiology*, **50**: 93–100.
- George, S.M., Lund, B.M. & Brocklehurst, T.F. 1988. The effect of pH and temperature on initiation of growth of *Listeria monocytogenes*. *Letters in Applied Microbiology*, **6**: 153–156.
- George, S.M., Richardson, L.C.C. & Peck, M.W. 1996. Predictive models of the effect of temperature, pH and acetic and lactic acids on the growth of *Listeria monocytogenes*. *International Journal of Food Microbiology*, **32**: 73–90.
- Golden, M.H., Buchanan, R.L. & Whiting, R.C. 1995. Effect of sodium acetate or sodium propionate with EDTA and ascorbic acids on the inactivation of *Listeria monocytogenes*. *Journal of Food Safety*, **15**: 53–65.
- Grau, F.H. 1993. Processed meats and *Listeria monocytogenes*. pp. 13–24, in: Prevention of *Listeria* in Processed Meats. Proceedings of a series of workshops. CSIRO Division of Food Science and Technology, Meat Research Laboratory, Queensland, Australia.
- Grau, F.H. & Vanderlinde, P.B. 1992. Occurrence, numbers, and growth of *Listeria monocytogenes* on some vacuum-packaged processed meats. *Journal of Food Protection*, **55**: 4–7.
- Grau, F.H. & Vanderlinde, P.B. 1993. Aerobic growth of *Listeria monocytogenes* on beef lean and fatty tissue: equations describing the effects of temperature and pH. *Journal of Food Protection*, **56**: 96–101.
- Hudson J.A., Mott, S.J. & Penny N. 1994. Growth of *Listeria monocytogenes*, *Aeromonas hydrophila* and *Yersinia enterocolitica* on vacuum and saturated carbon dioxide controlled atmosphere-packaged sliced roast beef. *Journal of Food Protection*, **57**: 204–208.
- ICMSF. 1996. *Microorganisms in Foods, Microbiological Specifications of Food Pathogens*. Vol. 5. London: Blackie Academic and Professional. 513p.
- Jarvis, B. 1989. *Statistical Aspects of the Microbiological Analysis of Foods*. Amsterdam, The Netherlands: Elsevier.
- Juneja, V.K. & Eblen, B.S. 1999. Predictive thermal inactivation model for *Listeria monocytogenes* with temperature, pH, NaCl, and sodium pyrophosphate as controlling factors. *Journal of Food Protection*, **62**: 986–993.

- Lebert, I., Bégot, C. & Lebert, A. 1998. Development of two *Listeria monocytogenes* growth models in a meat broth and their application to beef meat. *Food Microbiology*, **15**: 499–509.
- Linton, R.H., Carter, W.H., Pierson, M.D. & Hackney, C.R. 1995. Use of a modified Gompertz equation to model non-linear survival curves for *Listeria monocytogenes* Scott A. *Journal of Food Protection*, **58**: 946–954.
- Linton, R.H., Carter, W.H., Pierson, M.D., Hackney, C.R. & Eifert, J.D. 1996. Use of a modified Gompertz equation to predict the effects of temperature, pH and NaCl on the inactivation of *Listeria monocytogenes* Scott A heated in infant formula. *Journal of Food Protection*, **59**: 16–23.
- Lou, Y. & Yousef, A.E. 1999. Characteristics of *Listeria monocytogenes* important to food processors. pp. 131–225, in: Ryser & Marth, 1999, q.v.
- McDonald, K. & Sun, D.-W. 1999. Predictive food microbiology for the meat industry: a review. *International Journal of Food Microbiology*, **52**: 1–27.
- McClure, P.J., Kelly, T.M. & Roberts, T.A. 1991. The effects of temperature, pH, sodium chloride and sodium nitrite on the growth of *Listeria monocytogenes*. *International Journal of Food Microbiology*, **14**: 77–92.
- McClure, P.J., Roberts, T.A. & Otto Oguru, P. 1989. Comparison of the effects of sodium chloride, pH and temperature on the growth of *Listeria monocytogenes* on gradient plates and in liquid medium. *Letters in Applied Microbiology*, **9**: 95–99.
- McClure, P.J., Zwietering, M.H. & Roberts, T.A. 1993. Modelling bacterial growth of *Listeria monocytogenes* as a function of water activity, pH and Temperature. *International Journal of Food Microbiology*, **18**: 139–149.
- McClure, P.J., Beaumont, A.L., Sutherland, J.P. & Roberts, T.A. 1997. Predictive modelling of growth of *Listeria monocytogenes*. The effects on growth of NaCl, pH, storage temperature and NaNO<sub>2</sub>. *International Journal of Food Microbiology*, **3**: 221–232.
- McDonald, K. & Sun, D.-W. 1999. Predictive food microbiology for the meat industry: a review. *International Journal of Food Microbiology*, **52**: 1–27.
- McLauchlin, J. 1996. The relationship between *Listeria* and listeriosis. *Food Control*, **7**: 187–193.
- McMeekin, T.A., Olley, J., Ross, T. & Ratkowsky, D.A. 1993. *Predictive Microbiology. Theory and Application*. Taunton, UK: Research Studies Press. 340p.
- Mellefont, L.A., McMeekin, T.A. & Ross, T. 2003. The effect of abrupt osmotic shifts on the lag phase duration of foodborne bacteria. *International Journal of Food Microbiology*, **83**: 281–293.
- Mellefont, L.A. & Ross, T. 2003. The effect of abrupt shifts in temperature on the lag phase duration of *Escherichia coli* and *Klebsiella oxytoca*. *International Journal of Food Microbiology*, **83**: 295–305.
- Membre, J.M., Thurette, J. & Cateau, M. 1997. Modelling the growth, survival and death of *Listeria monocytogenes*. *Journal of Applied Microbiology*, **82**: 345–350.
- Murphy, P.M., Rae, M.C. & Harrington, D. 1996. Development of a predictive model for growth of *Listeria monocytogenes* in a skim milk medium and validation studies in a range of dairy products. *Journal of Applied Microbiology*, **80**: 557–564.
- Nerbrink, E., Borch, E., Blom, H. & Nesbakken, T. 1999. A model based on absorbance data on the growth rate of *Listeria monocytogenes* and including the effect of pH, NaCl, Na-lactate and Na-acetate. *International Journal of Food Microbiology*, **47**: 99–109.
- Notermans, S., Dufrenne, J., Teunis, P. & Chackraborty, T. 1998. Studies on the risk assessment of *Listeria monocytogenes*. *Journal of Food Protection*, **61**: 244–248.
- Peleg, M. & Cole, M.B. 1998. Reinterpretation of microbial survival curves. *Critical Reviews in Food Science and Nutrition*, **38**: 353–380.

- Piyasena, P., Liou, S. & McKellar, R.C. 1998. Predictive modelling of inactivation of *Listeria* spp. in bovine milk during high-temperature short-time pasteurization. *International Journal of Food Microbiology*, **39**: 167–173.
- Ratkowsky, D.A. 1992. Predicting response times in predictive food microbiology. Department of Primary Industry, Fisheries & Energy, Tasmania, Research and Development Unit, Biometrics Section, Occasional Paper No. 1992/1.
- Ratkowsky, D.A., Olley, J., McMeekin, T.A. & Ball, A. 1982. Relationship between temperature and growth rate of bacterial cultures. *Journal of Bacteriology*, **149**: 1–5.
- Ratkowsky, D.A., Ross, T., McMeekin, T.A. & Olley, J. 1991. Comparison of Arrhenius-type and Belehradek-type models for prediction of bacterial growth in foods. *Journal of Applied Bacteriology*, **71**: 452–459.
- Ratkowsky, D.A., Ross, T., Macario, T.W. & Kamperman, L. 1996. Choosing probability distributions for modelling generation time variability. *Journal of Applied Bacteriology*, **80**: 131–137.
- Razavilar, V. & Genigeorgis, C. 1998. Prediction of *Listeria* spp. growth as affected by various levels of chemicals, pH, temperatures and storage time in a model broth. *International Journal of Food Microbiology*, **40**: 149–157.
- Rees, C.E.D., Dodd, C.E.R., Gibson, P.T., Booth, I.R. & Stewart, G.S.A.B. 1995. The significance of bacteria in stationary phase to food microbiology. *International Journal of Food Microbiology*, **28**: 263–275.
- Ross, T. 1993. A philosophy for the development of kinetic models in predictive microbiology. Ph.D. Thesis, University of Tasmania, Hobart, Australia.
- Ross, T. 1996. Indices for performance evaluation of predictive models in food microbiology. *Journal of Applied Bacteriology*, **81**: 501–508.
- Ross, T. 1999. Predictive food microbiology models in the meat industry. Meat and Livestock Australia, Sydney, Australia. 196p.
- Ross, T., Baranyi, J. & McMeekin, T.A. 1999. Predictive Microbiology and Food Safety. pp. 1699–1710, in: R. Robinson, C.A. Batt and P. Patel (eds). *Encyclopaedia of Food Microbiology*. London: Academic Press.
- Ross, T. & McMeekin, T.A. 1994. Predictive microbiology - a review. *International Journal of Food Microbiology*, **23**: 241–264.
- Rowan, N.J. 1999. Evidence that inimical food-preservation barriers alter microbial resistance, cell morphology and virulence. *Trends in Food Science Technology*, **10**: 261–270.
- Ryser, E.T., & Marth, E.H. (eds). 1999. *Listeria, Listeriosis, and Food Safety*. 2nd edition, revised and expanded. New York NY: Marcel Dekker. 738p.
- Stephens, P.J., Cole, M.B. & Jones, M.V. 1994. Effect of heating rate on the thermal inactivation of *Listeria monocytogenes*. *Journal of Applied Bacteriology*, **77**: 702–708.
- Stephens, P.J., Joynson, J.A., Davies, K.W., Holbrook, R., Lappin-Scott, H.M. & Humphrey, T.J. 1997. The use of an automated growth analyser to measure recovery times of single heat injured *Salmonella* cells. *Journal of Applied Microbiology*, **83**: 445–455.
- Sutherland, J.P., Bayliss, A.P. & Roberts, T.A. 1994. Predictive modelling of the growth of *Staphylococcus aureus*: The effects of temperature, pH and sodium chloride. *International Journal of Food Microbiology*, **21**: 217–236.
- te Giffel, M.C. & Zwietering, M.H. 1999. Validation of predictive models describing the growth of *Listeria monocytogenes*. *International Journal of Food Microbiology*, **46**: 135–149.
- Tienungoon, S. 1998. Some aspects of the ecology of *Listeria monocytogenes* in salmonid aquaculture. Ph.D. Thesis, University of Tasmania, Hobart, Australia.

- Tienungoon, S., Ratkowsky, D. A., McMeekin, T. A. & Ross, T. 2000. Growth limits of *Listeria monocytogenes* as a function of temperature, pH, NaCl and lactic acid. *Applied and Environmental Microbiology*, **66**: 4979–4987.
- Tompkin, R.B. 2002. Control of *Listeria monocytogenes* in the food-processing environment. *Journal of Food Protection*, **65**: 709–725.
- Walls, I. & Scott, V.N. 1997. Validation of predictive mathematical models describing the growth of *Listeria monocytogenes*. *Journal of Food Protection*, **60**: 1142–1145.
- Wijtzes, T., McClure, P.J., Zwietering, M.H. & Roberts, T.A. 1993. Modelling bacterial growth of *Listeria monocytogenes* as a function of water activity, pH and temperature. *International Journal of Food Microbiology*, **18**: 139–149.
- Xiong, R., Xie, G., Edmondson, A.S., Linton, R.H. & Sheard, M.A. 1999. Comparison of the Baranyi model with the modified Gompertz equation for modelling thermal inactivation of *Listeria monocytogenes* Scott A. *Food Microbiology*, **16**: 269–279.
- Zhao, Y., Wells, J.H. & Marshall, D.L. 1992. Description of log phase growth for selected microorganisms during modified atmosphere storage. *Journal of Food Process Engineering*, **15**: 299–317.
- Zhao, P., Zhao, T., Doyle, M., Rubino, J., & Meng, J. 1998. Development of a model for evaluation of microbial cross-contamination in the kitchen. *Journal of Food Protection*, **61**: 960–963.
- Zwietering, M.H., Cuppers, H.G.A.M., deWit, J.C. & van't Riet, K. 1994. Evaluation of data transformation and validation of a model for the effect of temperature on bacterial growth. *Applied and Environmental Microbiology*, **60**: 195–203.
- Zemser, R.B. & Martin, S.E. 1998. Heat stability of virulence-associated enzymes from *Listeria monocytogenes* SLCC 5764. *Journal of Food Protection*, **61**: 899–902.



## Appendix 4.

# Prevalence and incidence of *Listeria monocytogenes* in Fermented Meat Products

### A4.1 REPORTED PREVALENCE AND INCIDENCE

The prevalence and incidence of *Listeria monocytogenes* in fermented meat products (FMPs) as reported in the literature is summarized in Table A4.1. It is to be noted, as mentioned in Section A4.2, below, that some products in the list in Table A4.1 might have a single name but represent very different products and processes in different countries. The authors have not attempted to distinguish these in the risk assessment modelling, but have instead treated all prevalence and concentration data as representative of all FMPs.

**Table A4.1** Reported prevalence and incidence of *Listeria monocytogenes* in fermented meat products

Product Description	Positive (samples or proportion)	Samples	% positive	Conc.	Location of Survey	Ref.
Fermented sausages	up to 0.20	5	20.00%		Various countries	[1] [2]
Fermented sausages					Austria	[3]
Fermented sausages	4	21	19.05%		Yugoslavia	[4]
Raw sausage	16	20	80.00%		Brazil	[5]
Fermented sausage	0.22 to 0.83				Spain	[6]
Dry sausages	0.22 to 0.83	18	44.00%		Various countries	[7] [8] [9] [10] [11]
Fermented sausages	6	30	20.00%		Canada	[12]
Raw sausage	13	25	52.00%		UK	[13]
Mettwurst with onion, fresh	1	11	9.09%		Germany	[14]
Sausages	2	8	25.00%		Hungary	[15]
Mettwurst with onion	27	245	11.00%		Germany	[16]
Spreadable, fermented	43	381	11.30%		Germany	[16]
Sliceable, fermented	11	228	4.80%		Germany	[16]
Raw sausage	30	120	25.00%	<100 CFU/g	Germany	[17]
Mettwurst, coarse	6	30	20.00%	<1000 CFU/g	Germany	[18]
Mettwurst, fresh	18	30	60.00%	<1000 CFU/g	Germany	[18]
Raw sausage, salami type	5	30	16.67%	<100 CFU/g	Germany	[18]
Beef sausage	0	1	0.00%		UK	[19]
Sausage	0	3	0.00%		UK	[19]
Raw fresh sausages	4	98	4.08%		France	[20]
Raw sausage	12	68	17.65%		Germany	[21]
Mettwurst, fresh	22	132	16.67%		Germany	[22]

Product Description	Positive (samples or proportion)	Samples	% positive	Conc.	Location of Survey	Ref.
Raw sausage, sliced	2	126	1.59%		Germany	[22]
Salsiccia	6	52	11.54%		Italy	[23]
Fermented sausages, salami type	0	70	0.00%		Norway	[24]
Ground/minced muscle (dry fermented sausages)	36	308	11.69%		Belgium	[25]
Fermented sausages	up to 0.20	5	20.00%	less than in non-fermented RTE cooked meats		[26]
Salami		128	10.00%		UK	[27]
Salami		67	16.00%		UK	[28]
Salami		59	5.00%	20 CFU/g	Switzerland	[29]
Mettwurst		14	0.00%		Switzerland	[30]
Dry cured		136	10.00%		Hungary	[31]
Fermented		21	10.00%		Hungary	[31]
Smoked		23	13.00%		Hungary	[31]
Cervelat		44	0.00%		South Africa	[32]
Vacuum-packed salami		19	0.00%		Australia	[33]
Salami		132	40.00%		Australia	[34]
Uncooked, preserved meat products						
1994/5 data	77	328	23.50%	1.8% > 10 (& <100) CFU/g; 0.6% >100 CFU/g		
1997 data from retail-level processors	19	132	14.40%	13.6% >10 (& <100) CFU/g; 0.8% > 100 CFU/g	Denmark	[35]
1998 data from retail-level processors	37	225	16.50%	14.7% >10 (& <100) CFU/g; 1.8% > 100 CFU/g		

SOURCES: [1] Breer and Schopfer, 1989. [2] Farber, Sanders and Johnston, 1989. [3] Breuer and Prandl, 1988. [4] Buncic, 1991. [5] Destro, Serrano and Kabuki, 1991. [6] Encinas et al., 1999. [7] Farber, Sanders and Johnston, 1989. [8] Nicolas and Vidaud, 1985. [9] McClain and Lee, 1988. [10] Breuer and Prandl, 1988. [11] Schmidt et al., 1988. [12] Farber, Sanders and Johnston, 1989. [13] Gilbert, Hall and Taylor, 1989. [14] Karches and Teufel, 1988. [15] Kiss et al., 1996. [16] 1991–2 data supplied to FAO/WHO by BgVV, Germany, in response to a call for data, 2000. [17] Leistner and Schmidt, 1992. [18] Leistner, Schmidt and Kaya, 1989. [19] MacGowan et al., 1994. [20] Nicolas and Vidaud, 1985. [21] Noack and Jockel, 1993. [22] Ozari and Stolle, 1990. [23] Pacini et al., 1995. [24] Rørvik and Yndestad, 1991. [25] Uyttendaele, Troy and Debevere, 1999. [26] WHO, 1988. [27] Velani and Gilbert, 1990. [28] Gilbert, 1991. [29] Trüssel, 1989. [30] Trüssel, 1989. [31] Kovacs-Domjan, 1991. [32] Vorster, Greebe and Nortje, 1993. [33] Grau and Vanderlinde, 1992. [34] Varabioff, 1992. [35] Nørnung, Andersen and Schlundt, 1999.

## A4.2 PRODUCTION METHODS AND STYLES OF FERMENTED MEATS

### A4.2.1 Introduction to fermented meat products

Fermented meats, including salami, have been manufactured for centuries (Lücke, 1985; Leistner, 1995; Ricke and Keeton, 1997). European sausages have been produced since the Middle Ages and per capita production and consumption of fermented meat products (FMPs) is still greatest in Europe. European migrants to North America, and elsewhere, took their FMPs methods and styles with them to their new homelands, where new variations evolved, i.e. these traditional products in some cases were changed to suit conditions in the New World. It is important, then, to recognize that FMPs products from the Old and New Worlds that have the same name may often differ in composition and processing. For example, all “Mettwurst” and “Teewurst” in the United States of America is cooked, and NaCl levels in United States of America products are normally higher than their European counterparts, due to regulations for the control of the parasite, *Trichinella*. United States of America producers typically use nitrite only, with no nitrate added (B. Tompkin, pers. comm., 2001).

Dry sausages include chorizo (Spanish, smoked, highly spiced), frizzes (similar to pepperoni, but not smoked), pepperoni (not cooked, air dried), Lola or Lolita and Lyons sausage (mildly seasoned pork with garlic), and Genoa salami (Italian, usually made from pork but may have a small amount of beef; in the preparation process it is moistened with wine or grape juice and seasoned with garlic).

Chinese-style fermented sausages, with pork as the main ingredient, are also common in Asia and date back thousands of years (Leistner, 1995; Yu and Chou, 1997). The Thai fermented sausage Nham is also receiving attention in the scientific literature (ASCA, 1986; Petchsing and Woodburn, 1990).

Most FMPs products have long shelf lives due to the combination of acidification (through fermentation), removal of oxygen, addition of compounds that favour the growth of some microbes while retarding the growth of others, and, ultimately, the removal of water.

Semi-dry sausages are usually heated in a smokehouse to fully cook the product and partially dry it. Semi-dry sausages are semi-soft sausages with good keeping qualities due to their lactic acid fermentation. "Summer Sausage" (another word for cervelat) is the general classification for mildly seasoned, smoked, semi-dry sausages like mortadella and Lebanon bologna.

Unless otherwise noted the information in these sections is drawn from Lücke (1985), Leistner (1995), Lücke (1995) and Ricke and Keeton (1997).

### A4.2.1 Processing

The fundamental steps involved in the production of FMPs are:

- chopping and mixing of ingredients, and filling into casing;
- fermentation; and
- drying (or maturation).

### A4.2.3 Ingredients

#### Meat and Fat

From a product quality perspective, the type of meat used in FMPs is important. It is less important for the microbiological safety of the product, unless some types of meat are more highly contaminated with pathogens than others. The proportion of meat to fat, and the type of fat, is not important microbiologically except in the sense that the proportion of fat affects the amount of free water in the product. Of the lean muscle in the mix, about 70–75% by weight is water. It is the *concentration* of the additives in the aqueous (water) phase of the food that is important for understanding the microbiology of the product. More fat in the mixture means that there is less lean meat, which in turn means less water. As a guide, for a product containing 30% (by weight) fat, water makes up only about 53% of the weight of the batch. Thus, the effective concentration of any water-soluble additives is about twice that predicted simply on the basis of its weight compared to the overall weight of the batch. During maturing of FMPs, weight losses of 20–30% occur in “semi-dry” FMPs, and even more for “dry”-style products. This is due to loss of water only, and further increases the effective concentration of the water-soluble components, so that the final concentration can be up to four times the apparent level added to the mixture expressed on a weight-for-weight basis.

**Table A4.2** Typical physico-chemical properties of styles of finished FMPs products.

Category	Final pH	Lactic acid (%)	Moisture : protein ratio	Moisture loss	Moisture <sup>(1)</sup>	Comments
Dry sausages	5.0–5.3 (<5.3)	0.5–1.0	<2.3:1	25–50	<35	See Note (2)
Cervelat			1.9:1		32–38	Shelf-stable
Cappicola			1.3:1		23–29	Shelf-stable
German Dauerwurst	4.7–4.8		1.1:1		25–27	Shelf-stable
German salami	4.7–4.8		1.6:1	1	34–35	Shelf-stable
Peperoni	4.5–4.8	0.8–1.2	1.6:1	35	25–32	Shelf-stable
Italian salami, hard or dry			1.9:1	30	32–38	Shelf-stable
Genoa salami	4.9	0.79	2.3:1	28	33–39	Shelf-stable
Thüringer, dry	4.9	1.0	2.3:1	28	46–50	Shelf-stable
Semi-dry sausages	4.7–5.1 (<5.3)	0.5–1.3	>2.3<3.7:1	8–15	45–50	See Note (3)
Lebanon bologna	4.7	1.0–1.3	2.5:1	10–15	56–62	Refrigerate
Cervelat, soft			2.6:1	10–15		Refrigerate
Salami, soft			2.3–3.7:1	10–15	41–51	Refrigerate
Summer sausage	<5.0	1.0	3.1:1	10–15	41–52	Refrigerate
Thüringer, soft			3.7:1		46–50	Refrigerate
For comparison						
Dried beef			2.04:1	29		
Beef jerky			0.75:1	>50	28–30	
Air-dried sausage			2.1:1			

NOTES: (1) Water activity ranges for dry and semi-dry sausages are <0.85 to 0.91, and 0.90 to 0.94, respectively. European Council Directive 77/99/EEC (Health problems affecting the production and marketing of meat products and certain other products of animal origin) requests  $a_w$  of <0.91 or pH <4.5 for dry sausages to be shelf-stable, or a combined  $a_w$  and pH of 0.95 and <5.2, respectively. (2) Heat processed (optional, but see note (4)); dried or aged after fermentation for moisture loss; smoked. (3) Heat processed (but see Note (4)); typically smoked; packaged after processing and chilling. (4) USDA/FSIS Title 9 CFR may be amended to require specified time and temperature heating combinations after fermentation, or verification that processing conditions destroy all pathogenic microorganisms. SOURCES: Various authors, cited in Ricke and Keeton, 1997.

### **Salt**

Typically, 2.5–3.3% NaCl (w/w) is added to FMPs mixes. The water activity ( $a_w$ ) of the product decreases during processing as the product loses water. This leads to effective concentrations in the typical finished semi-dry product of about 7.5–12% salt, corresponding to water activities in the range 0.95–0.92. Lower water activities (~0.85) are achieved in southern European style dry sausages (Calicioglu et al., 1997; Ricke and Keeton, 1997). Water activity values can be translated to aqueous phase NaCl concentration by reference to calibration tables or curves, e.g. Chirife and Resnik (1984).

### **Sugar, pH and organic acids**

Sugars (0.4–0.8%) are added to the mixture as a carbon source for the fermentative bacteria. These bacteria, usually lactic acid bacteria, metabolize the sugars, producing lactic acid in the process, which is released into the FMPs. In a review of lactic acid bacterial fermentation and the principal antimicrobial factors produced by lactic acid bacteria, Adams and Nicolaidis (1997) concluded that the principal antimicrobial factor is the ability of all lactic acid bacteria to produce organic acids and decrease the pH of foods in which they grow.

The biochemistry of conversion of simple sugars (e.g. glucose) results in almost twice as much lactic acid being produced as the concentration of simple sugars added. For more complex sugars, a smaller ratio of lactic acid to sugar results due to incomplete utilization of the carbohydrate. Other organic acids are also produced, but at much lower levels. The presence of lactic acid reduces the pH of the product during fermentation, typically to the range 4.6–5.0. The range of lactic acid concentrations in the final product is shown in Table A4.2. The range corresponds to total effective lactic acid concentrations (i.e. in the water phase) of from about 100 (e.g. semi-dry) to 500 mM (pepperoni).

### **Other additives**

Other ingredients of FMPs may include a variety of spices, and nitrite or nitrate. Spices, including pepper, paprika, garlic, mace, pimento and cardamom, may be added, but their primary role is sensory. The redox potential ( $E_h$ ) of FMPs is low. After mixing, the unfermented product is stuffed into casings. This effectively removes some oxygen. The predominant spoilage organisms of raw, aerobically stored meat, will be included in the mix and quickly consume residual oxygen. The presence of ascorbate and sugars also contributes to the creation of a low redox potential in the sausage.

#### **A4.2.4 Production of “safe” FMPs**

Production of safe FMPs requires prevention of the growth of pathogens during the fermentation step and maximizing death of surviving pathogens during maturation and storage. Some processors (especially in North America) include a heating step after fermentation that is intended to inactivate pathogens, including *Salmonella*, pathogenic *E. coli* and *Trichinella spiralis*. The initial stages of the fermentation process can permit growth of enteric pathogens such as *Salmonella*, *E. coli* and *Staphylococcus aureus*. The rapid acidification of the medium by the starter culture is considered a critical control point for minimization or prevention of pathogen growth (Bacus, 1997).

### A4.3 PHYSICO-CHEMICAL PARAMETERS OF FMPs


See Table A4.3 and Figure A 4.1

**Table A4.3** Typical composition and processing parameters for various FMPs styles

Examples	Semi-dry sausage	Dry sausage	
		Northern European type	Southern European type
	Summer sausages German cervelat Bologna sausages	German salamis Danish salamis	Italian salamis (Milanese; Calabrese) Saucissons secs Spanish chorizos
<b>Raw mixture</b>			
Meat : fat	Pork or beef, lean and fat	Lean pork : lean beef : fat pork (1 : 1 : 1)	Lean pork : fat pork (2 : 1)
Sugars (e.g. glucose, lactose, sucrose)	0.3–1.5%	0.3–0.8%	0–0.4%
Nitrate	–	–	<300 ppm
Nitrite	0–150 ppm	20–200 ppm	20–200 ppm
NaCl	2–2.5%	2–2.5%	2–2.5%
Seasoning (e.g. pepper, garlic, cardamom)	++	++	++
Starter cultures (10 <sup>6</sup> CFU/g)	Yes	Yes	Yes
<i>Lactobacillus sakei</i> , <i>L. curvatus</i> , <i>L. plantarum</i>	+	++	+++
<i>Pediococcus acidilactici</i> , <i>P. pentosaceus</i>	+++	+++	+
<i>Staphylococcus carnosus</i> , <i>S. xylosus</i>	–	++ or –	++
<i>Kocuria varians</i>	–	++ or –	+
<i>Penicillium chrysogenum</i> , <i>P. nalfiogense</i>	+	–	++
<i>Debaryomyces hansenii</i> , <i>Candida lipolytica</i>	+	–	++
<b>Fermentation period</b> (time/temperature/ relative humidity)	15–20 h/27–41°C/90% (USA) 18–48 h/20–32°C/85–95% (Germany)	18–48 h/20–30°C/58–95%	Day 1 – 22–24°C/94–96% Day 2 – 20–22°C/90–92% Day 3 – 18–20°C/85–88% or 2–3 d at 22–25°C/90–95%
<b>Drying period</b> (time/temperature/relative humidity)	2–3 d/10°C/68–72% (USA) 10–25 days (Germany)	1–3 weeks/12–15°C/75–80%	4–6 weeks/12–15°C/75–78%, or 8–14 weeks (traditional)
<b>Method of production</b>			
Smoking	Yes	Yes	No
Cooking	Yes	No	No
<b>Product characteristics</b>			
Final pH	4.4–5	4.6–5.1	5.1–5.5
Final a <sub>w</sub>	0.93–0.98	0.92–0.94	0.85–0.86
Water content	40–50%	30–40%	20–30%
Moisture : protein ratio (w/w)	2.3–3.7	2–2.3	1.6–1.9

KEY: + = occasionally used; ++ = frequently used; +++ = regularly used; – = not used.

SOURCE: Reprinted from Montel, M.C. Fermented meat products. pp. 745–753, in: R.K. Robinson, C.A. Batt and P.D. Patel (eds). *Encyclopaedia of Food Microbiology*. Copyright (2000), with permission from Elsevier.



**Figure A4.1** Collation of final pH and water activity of fermented meat products available in North America (open diamonds) and Australia (closed diamonds). pH and water activity limits below which the product is considered safe, in the absence of other inhibitors of microbial growth, are also shown (dotted lines)

SOURCE: After Ross et al., in press.

#### A4.4 ESTIMATION OF LACTIC ACID CONCENTRATION IN FMPs


The primary determinant of pH in meat and fermented meat is lactic acid. Natural levels in post-mortem meat are up to 125 mM (Gill, 1982) for meat in the pH range 5.5–6.5.

pH values, lactic acid and moisture content levels presented in Table A4.1 were tabulated. The lactic acid level was converted to lactic acid (using the Henderson-Hasselbalch equation) in the aqueous phase and pH plotted against lactic acid concentration. Data for pH and lactic acid concentration in meat were also included in the tabulation, and the data plotted (*see* Figure A4.2).

The simple regression through the data is described by the line:

$$\text{lactic acid (ppm)} = 50 + (((6.6 - \text{pH}) / 2.3) \times 300)$$

The model was generated based on the assumption that the lactic acid concentration in meat at pH 6.6 (highest pH reported in salami in Figure A4.1) is 50 mM, and that the lactic acid concentration in the salami with the lowest pH (4.3) is ~350 mM (*see* Section A4.2). It was further assumed that pH was directly related to lactic acid concentration in the range pH 6.6–4.3.



**Figure A4.2** Relationship between the pH of FMPs and the aqueous phase lactic acid concentration.


#### **A4.5 DATA ILLUSTRATING THE REMAINING SHELF-LIFE OF AUSTRALIAN FMPs AT THE TIME OF PURCHASE BY DOMESTIC CONSUMERS**

An ad hoc survey was conducted of retail outlets in Hobart, Australia, including supermarkets and small stores and delicatessens, of nominal remaining shelf-life of fermented meats on retail display. The survey involved 28 samples of 13 different products from 3 Australian producers. The survey was part of a larger survey of all processed meats, involving >700 samples.

By comparing the “use-by” (or “expiration”) date with the survey data, it is possible to infer the nominal shelf-life remaining. From the “use-by” date and manufacture date it is possible to infer the total shelf-life. The survey revealed that the mean stated shelf-life of Australian FMPs is 140 ( $\pm 70$ ) days, but samples included only three producers.

It is assumed that the survey represents a snapshot of the remaining shelf-life that a purchaser would have available to them. Full details of the survey are presented in Ross et al. (in press).

A summary is shown in Figure A4.3.



**Figure A4.3** A histogram of nominal remaining shelf-life of Australian fermented meats on retail display (heavy lines) and the cumulative frequency curve derived from that data. The sample size was 24, comprising 13 different products from three producers.

#### A4.6 SERVING SIZE DISTRIBUTION


See Figure A4.4.

#### A4.7 NATIONAL CONSUMPTION OF FMPs

Differences in consumption between nations is expected and reported indirectly (Holdsworth et al., 2000), but practically no data quantifying national FMPs production were found. Typically, FMPs statistics are included in total processed meat statistics. However, some data were extracted from supermarket sales records and nutritional surveys, as described below.

##### United States of America

FDA/FSIS (2001) estimated an average of 6.41 servings of FMPs per annum for a population of 271 000 000 (NGS, 1999). This consumption is relatively low compared to other developed nations, and expert opinion (B. Tompkin, pers. comm., 2001) also suggests that this estimate of consumption is unrealistically low. The 50<sup>th</sup> percentile serving size is 46 g. This equates to 295 g/person-year, or a total national consumption of 82 265 tonne/year. It is noted that total consumption of “deli meats” in the United States of America is 12 times higher than FMPs consumption.



**Figure A4.4** Comparison of serving size distribution from United States of America data (FDA/FSIS, 2001) (top) and that generated by the model used in this study (bottom).

## Australia

National survey data (ABS, 1995) suggests that annual per capita consumption of processed meat is between 5 and 19 kg, but this could include sausages and other meat products intended to be cooked prior to consumption. From diverse Australian production and sales data reviewed by Ross et al. (in press), it was estimated that total production of FMPs in Australia was 7795–32 379 tonne/year, equivalent to an annual per capita consumption of 400–1680 g. It is noted that this also suggests that in Australia the total FMPs consumption is one tenth to one twelfth of total processed meats, as reported for the United States of America. The Australian population is about 19 800 000 (ABS, 2002).

## Canada

National consumption of FMPs in Canada is estimated as 912 g/person/year, based on 5% of consumers daily eating approximately 50 g. Canada has a population of about 31 000 000 (NGS, 1999).

## Germany

van Schothorst (1997) suggests that per capita annual average national consumption of processed meats in Germany is 28.5 kg. Assuming that 10% of this consumption is FMPs, (analogous to Australian and United States of America estimates), an estimate of 2.85 kg FMPs/person/year is made. However, German survey data from 1986 (G. Klein, pers. comm., 2000) indicate that per capita annual consumption of semi-dry fermented salami averaged only 723 g in West Germany, lower than that estimated by comparison with other nations. The reason for this large difference in estimates is currently unresolved. The population of Germany is about 81 000 000 (NGS, 1999).

## Finland

FFDIF (2000) reported that in 1998 and 1999 national annual per capita consumption of processed meat averaged 32 kg. National consumption included about 7000 tonnes dry sausage, 120 000 tonnes of other sausage, and 38 000 tonnes of hams and other processed meats. The population of Finland is approximately 5 170 000 (NGS, 1999). If it is assumed that dry sausage refers only to FMPs, then consumption is estimated at 1.35 kg/person/year. If the assumption from other nation's data is used, i.e. that 10% of processed meat consumption is FMPs, estimated annual per capita consumption is about 3.2 kg, and for consistency this was the estimated value used in the present risk assessment.

## A4.8 REFERENCES CITED IN APPENDIX 4

- ABS [Australian Bureau of Statistics]. 1995. Australian National Nutrition Survey 1995. Australian Government Publishing Service, Canberra, Australia.
- ABS. 2002. *Australia Now. Population. Population clock*. Downloaded 14/10/2002 from [www.abs.gov.au/Ausstats/ABS](http://www.abs.gov.au/Ausstats/ABS)
- Adams, M.R. & Nicolaides, L. 1997. Review of the sensitivity of different foodborne pathogens to fermentation. *Food Control*, **8**: 227–239.
- ASCA [Association for Science Cooperation in Asia]. 1986. A Concise Handbook of indigenous fermented foods in the ASCA Countries. Edited by S. Saono, R.R. Hull and B. Dhamcharee. Association for Science Cooperation in Asia, Australian Government Publishing Service, Canberra.
- Bacus, J. 1997. Processing procedures to control *Salmonella* and *E. coli* in fermented sausage products. *Food Australia*, **49**: 543–547.
- Breer, C. & Schopfer, K. 1989. Listerien in Nahrungsmitteln [*Listeria* in foodstuffs]. *Schweizerische Medizinische Wochenschrift*, **119**: 306–311.
- Breuer, V. J. & Prandl, O. 1988. Nachweis von Listerien und deren Vorkommen in Hackfleisch und Mettwurst in Österreich [The detection and incidence of *Listerias* in ground beef and mettwurst in Austria]. *Archiv für Lebensmittelhygiene*, **39**: 28–30.

- Buncic, S. 1991. The incidence of *Listeria monocytogenes* in slaughtered animals, in meat and in meat products in Yugoslavia. *International Journal of Food Microbiology*, **12**: 173–180.
- Calicioglu, M., Faith, N.G., Buege, D.R. & Luchansky, J.B. 1997. Viability of *Escherichia coli* O157:H7 in fermented semidry low-temperature-cooked beef summer sausage. *Journal of Food Protection*, **60**: 1158–1162.
- Chirife, J. & Resnik, S.L. 1984. Unsaturated solutions of sodium chloride as reference sources of water activity at various temperatures. *Journal of Food Science*, **49**: 1486–1488.
- Destro, M.T., Serrano, A.D. & Kabuki, D.Y. 1991. Isolation of *Listeria* species from some Brazilian meat and dairy products. *Food Control*, **2**: 110–112.
- Encinas, J.P., Sanz, J.J., Garcia-Lopez, M.-L. & Otero, A. 1999. Behaviour of *Listeria* spp. in naturally contaminated chorizo (Spanish fermented sausage). *International Journal of Food Microbiology*, **46**: 167–171.
- Farber, J.M., Sanders, G.W. & Johnston, M.A. 1989. A survey of various foods for the presence of *Listeria* species. *Journal of Food Protection*, **52**: 456–458.
- FDA/FSIS [U.S. Food and Drug Administration/Food Safety and Inspection Agency (USDA)]. 2001. Draft Assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. Center for Food Safety and Applied Nutrition (FDA) and Food Safety Inspection Service (USDA) (Available at: [www.foodsafety.gov/~dms/lmrisk.html](http://www.foodsafety.gov/~dms/lmrisk.html)). [Report published September 2003 as: Quantitative assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. Available at: [www.foodsafety.gov/~dms/lmr2-toc.html](http://www.foodsafety.gov/~dms/lmr2-toc.html)].
- FFDIF [Finnish Food and Drink Industries Federation]. 2000. Facts about Finnish Food Industry. Downloaded from: <http://www.etl.fi/english/stat/pdf/facts.pdf> (22 October 2002).
- Gilbert, R.J. 1991. Occurrence of *Listeria monocytogenes* in foods in the United Kingdom. pp. 82–88, in: Proceedings of the International Conference on Listeriosis and Food Safety. Laval, France, cited in J.M. Farber and P.I. Peterkin, 1999. Incidence and behaviour of *Listeria monocytogenes* in meat products. pp. 505–564, in: Ryser & Marth, 1999, q.v.
- Gilbert, R.J., Hall, S.M. & Taylor, A.G. 1989. Listeriosis update. *U.K. PHLS Microbiology Digest*, **6**: 33–37.
- Gill, C.O. 1982. Microbial interaction with meats. pp. 225–264, in: M.H. Brown (ed). *Meat Microbiology*. UK: Applied Science Publishers.
- Grau, F.H. & Vanderlinde, P.B. 1992. Occurrence, numbers, and growth of *Listeria monocytogenes* on some vacuum-packaged processed meats. *Journal of Food Protection*, **55**: 4–7.
- Holdsworth, M., Gerber, M., Haslam, C., Scali, J., Bearsdworth, A., Avallone, M.H. & Sherratt, E. 2000. A comparison of dietary behaviour in Central England and a French Mediterranean region. *European Journal of Clinical Nutrition*, **54**: 530–539.
- Karches, H. & Teufel, P. 1988. *Listeria monocytogenes*. Vorkommen in Hackfleisch bei Kuhl- und Gefrierlagerung. *Fleischwirtschaft*, **68**: 1388–1392.
- Kiss, R., Papp, N. E., Vamos, G. Y. & Rodler, M. 1996. *Listeria monocytogenes* isolation from food in Hungary. *Acta Alimentaria*, **25**: 83–91.
- Kovacs-Domjan, H. 1991. Occurrence of *Listeria* infection in meat industrial raw-materials and end products. *Magyar Allatorvosok Lapja*, **46**: 229–233.
- Leistner, L. 1995. Stable and safe fermented sausages world-wide. pp. 160–175, in: G. Campbell-Platt and P.E. Cook, (eds). *Fermented Meats*. UK: Blackie Academic and Professional.

- Leistner, L. & Schmidt, U. 1992. Aktuelles über Listerien in Fleisch. *Mitteilungsblatt der Bundesanst. für Fleischforschung, Kulmbach*, **31**: 197–206.
- Leistner, L., Schmidt, U. & Kaya, M. 1989. Listerien bei Fleisch und Fleischerzeugnissen. [Listeria in meat and meat products]. *Mitteilungsblatt der Bundesanst. für Fleischforschung, Kulmbach*, **104**: 192–199.
- Lücke, F.-K. 1985. Fermented sausages. pp. 41–83, in: B.J.B. Wood (ed). *Microbiology of Fermented Foods*. London: Elsevier Applied Science Publ.
- Lücke, F.-K. 1995. Fermented meats. pp. F-1–F-23, in: *LFRA Microbiology Handbook: Meat Products*. Leatherhead, Surrey, UK: Food Research Association.
- MacGowan, A.P., Bowker, K., McLauchlin, J., Bennett, P.M. & Reeves, D.S. 1994. The occurrence and seasonal changes in the isolation of *Listeria* spp. in shop bought foodstuffs, human faeces, sewage and soil from urban sources. *International Journal of Food Microbiology*, **21**: 325–334.
- McClain, D. & Lee, W.H. 1988. Development of USDA FSIS method for isolation of *Listeria monocytogenes* from raw meat and poultry. *Journal of the Association of Official Analytical Chemists*, **71**: 660–664.
- Montel, M.C. 2000. Fermented meat products. pp. 745–753, in: R.K. Robinson, C.A. Batt and P.D. Patel (eds). *Encyclopaedia of Food Microbiology*. San Diego CA: Academic Press.
- NGS [National Geographic Society]. 1999. The National Geographic desk reference. Edited by R.M. Downs, F.A. Day, P.L. Knox, P.H. Meserve and B. Warf. Washington DC: National Geographic Society. 700p.
- Nicolas, J.A. & Vidaud, N. 1985. Contamination des viandes et des produits de charcuterie par *Listeria monocytogenes* en Haute-Vienne. *Sciences des aliments*, **5**: 175–180.
- Noack, D.J. & Joeckel, J. 1993. *Listeria monocytogenes*, occurrence and significance in meat and meat products and experience with recommendations for its detection and assessment. *Fleischwirtschaft*, **73**: 581–584.
- Nørrung, B., Andersen, J.K. & Schlundt, J. 1999. Incidence and control of *Listeria monocytogenes* in foods in Denmark. *International Journal of Food Microbiology*, **53**: 195–203.
- Ozari, R. von & Stolle, F. A. 1990. Zum Vorkommen von *Listeria monocytogenes* in Fleisch und Fleischerzeugnissen einschliesslich Geflügelfleisch des Handels. *Archiv. für Lebensmittelhygiene*, **41**: 47–50.
- Pacini, M., Quagli, E., Galassi, R., Marinari, M. & Marzotto, G. 1995. Profilo microbiologico di alcuni prodotti alimentari di largo consumo. *Annali di Microbiologia ed Enzimologia*, **45**: 37–49.
- Petchsing, U. & Woodburn, M.J. 1990. *Staphylococcus aureus* and *Escherichia coli* in nham (Thai-style fermented pork sausage). *International Journal of Food Microbiology*, **10**: 183–192.
- Ricke, S.C. & Keeton, J.T. 1997. Fermented meat, poultry, and fish products. pp. 610–628, in: M.P. Doyle, L.R. Beuchat and T.J. Montville (eds). *Food Microbiology Fundamentals and Frontiers*. Washington DC: ASM Press.
- Rørvik, L.M. & Yndestad, M. 1991. *Listeria monocytogenes* in foods in Norway. *International Journal of Food Microbiology*, **13**: 97–104.
- Ross, T., Fazil, A., Paoli, G. & Sumner, J. In press. *Listeria monocytogenes* in Australian processed meat products: risks and their management. Meat and Livestock Australia, Sydney, NSW, Australia.

- Ryser, E.T., & Marth, E.H. (eds). 1999. *Listeria, Listeriosis, and Food Safety*. 2nd edition, revised and expanded. New York NY: Marcel Dekker. 738p.
- Schmidt, U., Seeliger, H.P.R., Glenn, E., Langer, B. & Leistner, L. 1988. Listerienfunde in rohen fleischerzeugnissen [Listeria findings in raw meat products]. *Fleischwirtschaft*, **68**: 1313–1316.
- Trüssel, M. 1989. The incidence of *Listeria* in the production of cured and air-dried beef, salami and mettwurst. *Schweizer Archiv für Tierheilkunde*, **131**: 409–421.
- Uyttendaele, M., de Troy, P. & Debevere, J. 1999. Incidence of *Salmonella*, *Campylobacter jejuni*, *Campylobacter coli* and *Listeria monocytogenes* in poultry carcasses and different types of poultry products for sale on the Belgian retail market. *Journal of Food Protection*, **62**: 735–740.
- van Schothorst, M. 1997. Practical approaches to risk assessment. *Journal of Food Protection*, **60**: 1439–1443.
- Varabioff, Y. 1992. Incidence of *Listeria* in smallgoods. *Letters in Applied Microbiology*, **14**: 167–169.
- Velani, S. & Gilbert, R.J. 1990. *Listeria monocytogenes* in prepacked ready-to-eat sliced meals. *UK PHLS Microbiology Digest*, **7**: 56.
- Vorster, S.M., Greebe, R.P. & Nortje, G.L. 1993. The incidence of *Listeria* in processed meats in South Africa. *Journal of Food Protection*, **56**: 169–172.
- WHO. 1988. Foodborne listeriosis. Report of the WHO Working Group. *Bulletin of the World Health Organization*, **66**: 421–428.
- Yu, C.F., & Chou, C.C. 1997. Fate of *Escherichia coli* O157:H7 in Chinese-style sausage during the drying step of the manufacturing process as affected by the drying condition and curing agent. *Journal of the Science of Food and Agriculture*, **74**: 551–556.

# Appendix 5.

## Background for the cold-smoked fish assessment

### A5.1 ESTIMATE OF GLOBAL PRODUCTION AND NATIONAL AND INDIVIDUAL CONSUMPTION OF COLD-SMOKED FISH

#### A5.1.1 Scope

The most abundant type of cold-smoked fish is cold-smoked salmon. Due to a paucity of data, cold-smoked fish consumption estimates were based on data describing global production of cold-smoked salmon.

#### A5.1.2 National and global consumption characteristics

National consumption was initially estimated from data in Globefish (1996)<sup>3</sup> detailing national production and imports/exports of cold-smoked salmon, as shown in Table A5.1. Various other sources of consumption estimates are also shown, and it is noted that the estimates from different sources are not completely consistent, which leads to uncertainty in the estimates.

National population data (NGS, 1999) were combined with the national consumption (calculated as “disappearance” data), to determine per capita consumption. Those data are also shown in Table A5.1.

In Germany and Denmark, hot-smoked product constitutes only a negligible or very small proportion of smoked salmon consumption (P.K. Ben Embarek, pers. comm., 2000; G. Klein, pers. comm., 2000). Similarly, in Australia, hot-smoked salmon products constitute ~10% of production and consumption (Walsh, 1999). Conversely, the contribution of other types of cold-smoked fish is not included in the estimates. Recognizing this limitation, the data are nonetheless used as proxy values for total cold-smoked fish consumption.

From that data, there are various approaches available to calculate the annual per-person consumption of cold-smoked fish and its variability and uncertainty. If the total population of the nations is considered against the total production, the average consumption is 90.2 g/person/year. Per capita consumption in individual nations appears to vary between 8 and 1000 g/person/year, with a median value of 138 g/consumer/year. The average of the estimates of national per-person annual consumption is, however, 231 g. This estimate

---

<sup>3</sup> Globefish have published an updated report on Salmon - A Study of Global Supply and Demand (Globefish, 2003). This provides more recent data on national production and imports/exports of cold-smoked salmon. However, due to limited time and resources it was not possible to incorporate the more recent data into this risk assessment.

differs from the global average obtained if each national data set has equal weight in the average global consumption estimate. If each national consumption estimate is weighted according to the population size, the global average is calculated to be 146 g. The difference between this and the original global estimates arises because data for Canada, Chile, Germany and West Germany, and Norway could not be used because one element of the needed data was missing; see Table A5.1. The median of the remaining national estimates is 144 g/person/year.

**Table A5.1** Production, import and export of cold-smoked salmon, and estimated per capita consumption.

	<b>Production (P)</b> (tonne)	<b>Import (I)</b> (tonne)	<b>Export (E)</b> (tonne)	<b>Consumption (inferred from P+I - E)</b> (tonne)	<b>Population</b> (million)	<b>Consumption per person-year</b> (gram)	<b>Note</b>
Australia	980	700	–	1 167	18.53	90.7	[1]
Austria	0	848	0	848	8.09	104.9	
Belgium	2 775	2 324	297	4 802	10.23	469.6	
Canada	501	–	501	0	30.59	–	
Chile	1 074	–	1 074	0	15.02	–	
Denmark	15 786	1 406	13 102	4 090	5.24	781.9	[2]
Denmark	–	–	–	–	–	202	[3]
Faeroe islands	407	–	407	0	–	–	
France	11 059	2 941	1 059	12 941	59.07	219.1	
Germany	5 063	7 279	693	11 649	81.95	142.1	
W. Germany	–	–	–	–	–	47.8	[4]
Germany	–	–	–	–	–	1 000	[5]
Ireland	357	–	357	0	3.73	–	
Italy	2 500	5 100	0	7 600	57.72	131.7	
Italy (1998)	–	–	–	9 000	57.72	145.5	[6]
Japan	7 853	765	0	8 618	126.75	68.0	
Netherlands	3 668	0	0	3 668	15.80	232.2	
Norway	2 446	0	2 446	0	4.46	–	
Others	3 954	5615	2 916	6 653	–	–	
Spain	0	313	0	313	39.42	7.9	
Sweden	–	–	–	2 400	8.86	271.0	[7]
Switzerland	0	656	0	656	7.12	92.1	
UK	11 000	146	3 247	7 899	59.36	133.1	
USA	5 116	1133	164	6 085	270.93	22.5	
<b>Total or average</b>	<b>73 562</b>	<b>28 526</b>	<b>26 263</b>	<b>79 389</b>	<b>880.56</b>	<b>90.2</b>	

SOURCES: Data from Globefish (1996) unless otherwise noted.

NOTES: [1] Estimated by Ross and Sanderson, 2000. [2] P.K. Ben Embarek, pers. comm., 2000. [3] Danish food Authority via P.K. Ben Embarek, pers. comm., 2000, [4] 1986 data, G. Klein, pers. comm., 2000, based on 1998 population, including East Germany. [5] Buchanan et al., 1997. [6] AC Nielsen data supplied to FAO, 2000. [7] Lindqvist and Westö, 2000.

### A5.1.3 Serving size estimates


#### *Individual consumption frequency by nation*

There is limited data available on the number of consumers who eat cold-smoked salmon products. West German data from 1986 (G. Klein, pers. comm., 2000,) reports that 311 of 23 131 interviewees (1.34%) consumed cold-smoked salmon on the survey day and that the mean serving size among eaters was 9.77 g/day, but with an upper 95<sup>th</sup> percentile of 28.60 g. From the same data source, differences among population sub-groups were revealed but are not used explicitly in this assessment. Using data for consumption of all smoked seafoods, there was no significant difference in serving size by geographical region (north or south Germany) or age group (more or less than 60 years).


In Australia, cold-smoked salmon is considered a luxury food. National consumption was estimated by Ross and Sanderson (2000) at approximately 0.15–0.20 kg/person/year, roughly equivalent to 1% of consumers per day eating a 60 g serving of cold-smoked salmon, or all members of the population eating 3 to 4 servings per year.

Canadian data (CFPNS, 1992–1995) shows that cold-smoked salmon is consumed infrequently in that country. Approximately 5% of consumers on the survey day ate cold-smoked fish products, which included kippered Atlantic herring; cold-smoked Chinook (spring) salmon; smoked haddock; Chinook (lox) salmon; and smoked cod. About half of these were smoked salmonid products. It should be noted that smoked cod is normally cooked before consumption, but represents 8% of eating occasions in the data.

Smoked fish meal size data were estimated by FDA/FSIS (2001) using data from CSFII and NHANES. The data were modified for this case study by removal of data (mostly for smoked oysters) that did not relate to smoked fish products. The average serving size based on age and gender is shown in Figure A5.1. It should be noted that there were few data available – the number of data represented by each bar in the figure varies from 1 to 8.



**Figure A5.1** Cold-smoked fish serving size as a function of age and gender based on United States of America data (FDA/FSIS, 2001).



**Figure A5.2** Modelled cumulative probability distribution of serving sizes.


Those data indicate that serving size for cold-smoked fish products varies between 1 and 357 g with a median value of ~50 g per serving, and an average value of 58 g/serving. Similarly, using Canadian data (CFPNS, 1992–1995), consumption amounts were aggregated over the smoked fish foods considered, up to the amount consumed on all occasions in a day. The observations are skewed, with median value at 61 g, and long upper tail extending to approximately 225 g, representing approximately the 97.5<sup>th</sup> percentile.

The above serving size data was used to estimate per capita frequency of consumption from the annual per person consumption estimates in Table A5.1. Those data were used as the basis for the meal size distribution that was used in the model, which is shown in Figure A5.2 and was described in the model using a CumDist function based on the values shown in Table A5.2. The median value of the distribution is 57 g, and the mean value is 63 g.

For the nations considered, the consumption frequency is in the range 0.15–18 servings per person per year, with most in the range 2–5 servings per person per year. The distribution of the number of meals per consumer per year in the model is described empirically by Beta(0.5, 2.5, 0, 18), as shown in Figure A5.3.

**Table A5.2** Values used to generate the distribution of serving sizes of cold-smoked fish used in the simulation model.

Serving size (g)	Cumulative probability
0	0.00
57	0.50
75	0.75
136	0.95
142	0.99
284	1.00



**Figure A5.3** Modelled cumulative probability distribution of number of cold-smoked fish servings compared to the observed data.

#### A5.1.4 Reality check

From the values described above, global total annual consumption in the nations considered can be estimated from the simulation model from the distribution of the product of serving frequency estimate and serving size estimate. This is a useful check on the performance of the simulation model. The median modelled global consumption is 62 300 tonnes and the mean modelled global consumption is 118 000 tonnes. The latter value is ~50% higher than the consumption of cold-smoked salmon estimated from the data in Table A5.1. The basis of this difference is not known with certainty, but may derive from the fact that serving size estimates in the model are derived from all types of smoked fish whereas consumption is based on smoked salmon data only.


### A5.2 DESCRIPTION OF STORAGE TEMPERATURES

The storage temperature distribution (Section 4.5.3.4) is derived from Audits International (2000) data for refrigerated cabinets used for storage of cold-smoked fish at retail.

The data used is tabulated below (Table A5.3) and was fitted empirically to a Beta distribution. Comparison of the original data and the fitted distribution is shown in Figure A5.4.

**Table A5.3** Data used to simulate storage temperature.

Temperature (°C)	Observed % Frequency	Cumulative % Frequency
-1.67	5	5
0.00	5	10
1.67	8	18
3.33	21	39
5.00	26	65
6.67	16	81
8.33	8	89
10.00	7	96
11.67	2	98
13.33	0.4	98.4
15.00	0.5	98.9
16.67	0.5	99.4
18.33	0.2	99.6
20.00	0.2	99.8
21.67	0.2	100

**Figure A5.4** Comparison of observed and fitted temperature distribution data.

## A5.3 DETAILS OF GROWTH MODELLING

### A5.3.1 Physico-chemical parameters

Ranges of physico-chemical parameters of cold-smoked fish that could affect growth of *L. monocytogenes* were presented in Section 4.5.3.5. In the simulation model, these features were described by the following distributions:

- water activity                      Normal(0.98, 0.0027)
- pH                                      Normal(6.2, 0.05)
- lactic acid concentration (mM) was estimated from pH by the empirical relationship:  
1105.0 -162.50 × pH.

Other growth inhibiting factors (e.g. phenol, spices, etc.) can be included in the model predictions by a simple multiplicative constant.

### A5.3.2 *L. monocytogenes* growth rate model

The *L. monocytogenes* growth rate model is derived from Tienungoon (1998). It is a square root type model. The model was further developed, described and evaluated against independent literature data by Ross (1999) and found to have  $B_f = 0.88$  and  $A_f = 1.94$  (measures of predictive model performance – see Ross, 1996, and Baranyi, Pin and Ross, 1999), which was as good as or better than other published models for *L. monocytogenes* growth rate. In the modelling, the growth rate prediction of the model was multiplied by 0.9 to compensate for the bias of the model. This correction was implemented using the “Other growth inhibiting factors” input in the simulation model.

To calculate growth, physico-chemical parameters sampled from the distributions described above are first “filtered” through the growth/no-growth model of Tienungoon et al. (2000) to determine whether the scenario sampled represents a product that will allow growth of *L. monocytogenes*. If growth is predicted to be possible, the extent of growth is modelled using the sampled storage time and the growth rate model, including a correction for lag time.


The prediction of *L. monocytogenes* growth is further filtered by applying an upper limit to the population density (CFU/g product) predicted to be achievable. Including the effects of lactic acid bacteria in the model is expected to preclude this being necessary in most model iterations, but in those (rare) scenarios where *L. monocytogenes* growth is modelled not to be constrained by any other factor, it will eventually limit its own growth, i.e. achieve its maximum population density (MPD). FDA/FSIS (2001) reviewed the available literature and noted that *L. monocytogenes* rarely achieves levels in cold-smoked salmon as high as it does in pure culture in laboratory broth. It is probable, however, that many of those observations are due to the effects of other bacteria in the foods, which are modelled in this assessment. MPD was therefore set at  $3 \times 10^9$  CFU/g, a level representative of otherwise ideal conditions for those scenarios in which no other factors constrain *L. monocytogenes* growth.

### A5.3.3 Lag time

Lag time data specific for *L. monocytogenes* in cold-smoked fish were not found in the literature. Ross (1999) collated data for lag times from the published literature and expressed

these as relative lag times (see Section 3.5.3.3). *L. monocytogenes* relative lag times in foods were in the range 0–40, with a peak value near 2.5. Lag times in laboratory broths had a similar range, but the peak value was nearer to 4.5. Figure A5.5 presents this data.

Dalgaard and Jørgensen (1998) state that *L. monocytogenes* cells that contaminate cold-smoked fish are likely to be damaged due to the effects of processing. Other studies (Rørvik and Yndestad, 1991; Rørvik et al., 1997, 2000), however, suggest that most contamination of cold-smoked fish arises after smoking, from contamination sources in processing plants. Because of this ambiguity, two distributions were assessed to gauge the importance of assumptions about lag time distributions. These were termed “short” (Beta(3, 30, 0, 35)) and “long” (Beta(6, 35, 0, 35)) relative lag times, and produce the distributions shown in Figure A5.6. The overall growth model and the interaction of factors governing the extent of growth are depicted in Figure A5.7.



**Figure A5.5** Relative lag time data for *Listeria monocytogenes* reported in published literature. SOURCE: After Ross, 1999.




Figure A5.6. Outputs from the simulation model showing the two relative lag time distributions used to model the effects of lag time on risk of listeriosis from cold-smoked fish.





Figure A5.7 Influence diagram taken from the simulation model used in this risk assessment and showing the interaction of factors used to estimate growth of *Listeria monocytogenes* during storage of cold-smoked fish.


### A5.3.4 Effect of lactic acid bacteria on shelf life and *L. monocytogenes* growth potential

The nominal shelf lives for vacuum-packed cold-smoked fish are in the range of 3 to 6 weeks at a storage temperature of 4–5°C. Several studies have assessed the sensory acceptability of cold-smoked salmon (Truelstrup Hansen, Drewes Røntved and Huss, 1998; Jørgensen, Dalgaard and Huss, 2000; Leroi et al., 2001) and found that at 5°C the sensory shelf-life of cold-smoked salmon is highly variable (3–9 weeks) and that there is no single indicator of the onset of spoilage. Attributes of cold-smoked salmon associated with spoilage are a softening of the texture and development of “stickiness” or “pastiness”, and the presence of “sour”, “bitter”, “rancid”, “ammoniacal”, “cabbage” and faecal odours. The lack of a clear relationship between microorganisms present and spoilage is illustrated by data in Figures A5.8 and A5.9, derived from two independent research groups, one working with a Danish product, the other with a French product. Thus, spoilage per se is difficult to model mechanistically.

In raw and processed meats and fish chilled and stored under vacuum, lactic acid bacteria become the dominant population and preserve the product with a “hidden” fermentation (Stiles, 1996). Thus, of particular note in Figures A5.8 and A5.9 is the cessation of growth of any component of the population when the total psychrotrophic count appears to achieve a stationary phase at a level of  $10^7$ – $10^8$  CFU/g. This behaviour is consistent with the Jameson effect (Stephens et al., 1997; see also Section 3.5.3.1).



**Figure A5.8** Microbiological changes in Danish-produced vacuum-packed cold-smoked salmon (4.6% water phase salt (WPS)) during storage at 5°C. Total count (○); total psychrotrophic count (▼); lactic acid bacteria (◇) and Enterobacteriaceae (■). Arrow indicates the time of sensory rejection. (Data of Truelstrup Hansen, Gill and Huss, 1995, reproduced from Gram and Huss, 1996).



**Figure A5.9** Microbiological changes in French-produced vacuum-packed cold-smoked salmon during storage at 5°C. Total psychrotrophic count (●); total lactic acid bacteria (▲); lactobacilli (△); Enterobacteriaceae (■); *Brocothrix thermosphacta* (X) and yeast (⋈). The arrow indicates the time of sensory rejection (Reproduced from Leroi et al., 2001).

The Jameson effect can be likened to a race to reach stationary phase. The winner is that sub-group within the total microbial population that first achieves stationary phase.

When that happens, the race is over and all other contestants finish the race (i.e. they also enter stationary phase) at that point in time, although unpublished data (L.A. Mellefont, B. Davidson and T. Ross, Univ. of Tasmania, pers. comm., 2002) indicate that in some cases growth is not completely inhibited, but is nevertheless greatly reduced.

The Jameson effect has relevance for estimation of the risk from microbiological hazards in cold-smoked fish products. As Figures A5.8 and A5.9 show, and as has been reported also for vacuum-packed meats (Mol et al., 1971; Egan, Ford and Shay, 1980; Korkeala et al., 1989), spoilage of vacuum-packed meat and fish does not usually occur until well after the total count has reached stationary phase. In cold-smoked salmon, that occurs after one to two weeks under recommended storage conditions. Thus, growth of pathogens in the product may only be possible for 25–50% of the full shelf-life (use-by period) of the product.

### A5.3.5 Modelling the effect of lactic acid bacteria

The mechanism of the Jameson effect is not yet fully understood. It may be due to competition for nutrients, production of toxic end products, or production of specific antibiotics against other bacteria. Under some circumstances, a pathogen may be numerically dominant at the time of production and, under improper storage, may grow fast enough to

reach stationary phase before any other element of the population on the food (e.g. see Nilsson, Huss and Gram, 1997). This possibility is explicitly recognized in the Seafood HACCP Alliance's recommendations (SHA, 1997):

“In cold-smoked fish, it is important that the product does not receive so much heat that the number of spoilage organisms are significantly reduced. This is true because spoilage organisms must be present to inhibit the growth and toxin formation of *C. botulinum* type E and nonproteolytic types B and F.”


Based on Figures A5.8 and A5.9, time to stationary phase of the lactic acid bacteria (i.e. the onset of the Jameson effect) at 5°C was described by Normal(14, 2) days. The variation in the time to reach stationary phase is assumed to be due to initial numbers of bacteria and specific product composition.

Because the time to stationary phase is expected to depend strongly on the temperature of storage, it is adjusted according to the temperatures selected using a relative rate function with a  $T_{min}$  of 0°C. This value was used as a first approximation, which was adjusted based on temperature–growth rate responses of lactic acid bacteria associated with vacuum-packed processed meat products, which have a similar microbial ecology (Mol et al., 1971).

The time to reach the stationary phase is deducted from the total storage time sampled in an iteration to determine the duration of the second, constrained, phase of growth. During the first phase of *L. monocytogenes* growth, the growth rate is predicted to be unconstrained and predicted by the growth rate model for the temperature and product characteristics sampled during that iteration. After that time, however, growth is predicted to be reduced by some factor. It could be complete inhibition (as described in Figures A5.8 and A5.9), but other data (L.A. Mellefont, B. Davidson and T. Ross, Univ. of Tasmania, pers. comm., 2002, unpublished data) that suggest that *L. monocytogenes* might continue to grow slowly. Thus, the specific growth behaviour is uncertain and has been left as an assumption whose influence can be tested (see Section 4.5.5).

Nominal storage life also has to be adjusted for storage conditions because higher temperatures will cause premature spoilage. Conversely, there are reports (Ben Embarek, pers. comm., 2001) that in some countries 2–3-month shelf lives are realized, apparently without consumer rejection. Accordingly, several scenarios have been modelled. However, in each case a filter is applied so that if the storage time at 5°C exceeds 10 weeks (or its equivalent calculated at other temperatures) the product is considered completely spoiled and no further growth occurs. This is achieved by truncating the predictions of growth based on shelf-life at the growth levels that could have occurred at the equivalent of 70 days at 5°C. These predictions are not, however, removed from the simulation.

These interactions are shown as an influence diagram in Figure A5.10.



**Figure A5.10** Influence diagram derived from the simulation model, showing the interaction of nominal storage life and spoilage due to bacterial growth, as well as the estimation of the time required for lactic acid bacteria to reach the stationary phase and for the Jameson effect to constrain *Listeria monocytogenes* growth.

#### A5.4 REFERENCES CITED IN APPENDIX 5

- Audits International. 2000. 1999 U.S. Food Temperature Evaluation. Design and Summary Pages. Audits International and U.S. Food and Drug Administration. 13p.
- Baranyi, J., Pin, C. & Ross, T. 1999. Validating and comparing predictive models. *International Journal of Food Microbiology*, **48**: 159–166.
- Buchanan, R.L., Damert, W.G., Whiting, R.C. & van Schothorst, M. 1997. Use of epidemiologic and food survey data to estimate a purposefully conservative dose-response relationship for *Listeria monocytogenes* levels and incidence of listeriosis. *Journal of Food Protection*. **60**: 918–922.
- CFPNS [Canadian Federal-Provincial Nutrition Surveys]. 1992–1995. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada. See also: Karpinski & Nargundkar, 1992; Junkins & Karpinski, 1994; Junkins, 1994; Junkins & Laffey, 2000; Junkins, Laffey & Weston, 2001.
- Dalgaard, P. & Jørgensen, L.V. 1998. Predicted and observed growth of *Listeria monocytogenes* in seafood challenge tests and in naturally contaminated cold-smoked salmon. *International Journal of Food Microbiology*, **40**: 105–115.
- Egan, A.F., Ford, A.L. & Shay, B.J. 1980. A comparison of *Mocrobacterium thermosphactum* and lactobacilli as spoilage organisms of vacuum-packaged sliced luncheon meats. *Journal of Food Science*, **45**: 1745–1748.

- FDA/FSIS [U.S. Food and Drug Administration/USDA Food Safety and Inspection Agency]. 2001. Draft Assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. Center for Food Safety and Applied Nutrition (FDA) and Food Safety Inspection Service (USDA) (Available at: [www.foodsafety.gov/~dms/lmrisk.html](http://www.foodsafety.gov/~dms/lmrisk.html)). [Report published September 2003 as: Quantitative assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods. Available at: [www.foodsafety.gov/~dms/lmr2-toc.html](http://www.foodsafety.gov/~dms/lmr2-toc.html)].
- Globefish. 1996. The World Market for Salmon. FAO Fishery Industries Division, Rome, Italy.
- Globefish. 2003. Salmon- A Study of Global Supply and Demand. FAO Fishery Industries Division, Rome, Italy.
- Gram, L. & Huss, H.H. 1996. Microbial spoilage of fish and fish products. *International Journal of Food Microbiology*, **33**: 121–138.
- Jørgensen, L.V., Dalgaard, P. & Huss, H.H. 2000. Multiple compound quality index for cold-smoked salmon (*Salmo salar*) developed by multivariate regression of biogenic amines and pH. *Journal of Agriculture and Food Chemistry*, **48**: 2448–2453.
- Junkins, E. 1994. Saskatchewan Nutrition Survey 1993/94. Methodology for estimating usual intake. BCCA Technical Report E451311-005. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada.
- Junkins, E. & Karpinski, K. 1994. Enquête québécoise sur la nutrition. Méthodologie pour estimer l'apport habituel, les statistiques sommaires et les erreurs-types. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada.
- Junkins, E. & Laffey, P. 2000. Alberta Nutrition Survey 1994. Methodology for estimating usual intake. BCCA Technical Report E451311-006. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada.
- Junkins, E., Laffey, P. & Weston, T. 2001. Prince Edward Island Nutrition Survey 1995. Methodology for estimating usual intake. BCCA Technical Report E451311-007. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada.
- Karpinski, K. & Nargundkar, M. 1992. Nova Scotia Nutrition Survey. Methodology Report. BCCA Technical Report E451311-001. Bureau of Biostatistics and Computer Applications, Food Directorate, Health Canada.
- Korkeala, H., Alanko, T., Makela, P. & Lindroth, S. 1989. Shelf-life of vacuum-packed cooked ring sausages at different chill temperatures. *International Journal of Food Microbiology*, **9**: 237–347.
- Leroi, F., Joffraud, J.J., Chevalier F. & Cardinal, M. 2001. Research of quality indices for cold-smoked salmon using a stepwise multiple regression of microbiological counts and physico-chemical parameters. *Journal of Applied Microbiology*, **90**: 578–587.
- Lindqvist, R. & Westöö, A. 2000. Quantitative risk assessment for *Listeria monocytogenes* in smoked or gravad salmon/rainbow trout in Sweden. *International Journal of Food Microbiology*, **58**: 181–196.
- Mol, J.H.H., Hietbrink, J.E.A., Mollen, H.W.M. & van Tinteren, J. 1971. Observations on the microflora of vacuum-packed sliced cooked meat products. *Journal of Applied Bacteriology*, **34**: 377–397.

- NGS [National Geographic Society]. 1999. The National Geographic desk reference. Edited by R.M. Downs, F.A Day, P.L. Knox, P.H. Meserve and B. Warf. Washington DC: National Geographic Society. 700p.
- Nilsson, L., Huss, H.H. & Gram, L. 1997. Inhibition of *Listeria monocytogenes* on cold-smoked salmon by nisin and carbon dioxide atmosphere. *International Journal of Food Microbiology*, **38**: 217–228.
- Rørvik, L.M. & Yndestad, M. 1991. *Listeria monocytogenes* in foods in Norway. *International Journal of Food Microbiology*, **13**: 97–104.
- Rørvik, L.M., Aase, B., Alvestad, T. & Caugant, D.A. 2000. Molecular epidemiological survey of *Listeria monocytogenes* in seafoods and seafood-processing plants. *Applied and Environmental Microbiology*, **66**: 4779–4784.
- Rørvik, L.M., Skjerve, E., Knudsen, B.R. & Yndestad, M. 1997. Risk factors for contamination of smoked salmon with *Listeria monocytogenes* during processing. *International Journal of Food Microbiology*, **37**: 215–219.
- Ross, T. 1996. Indices for performance evaluation of predictive models in food microbiology. *Journal of Applied Bacteriology*, **81**: 501–508.
- Ross, T. 1999. Predictive food microbiology models in the meat industry. Meat and Livestock Australia, Sydney, Australia. 196p.
- Ross, T. & Sanderson, K. 2000. A risk assessment of selected seafoods in New South Wales. SafeFood NSW, Sydney, Australia. 275p.
- SHA [Seafood HACCP Alliance]. 1997. Compendium of fish and fishery product processes, hazards and controls. National Seafood HACCP Alliance for Training and Education. Available at <http://seafood.ucdavis.edu/haccp/compendium/compend.htm>
- Stephens, P.J., Joynson, J.A., Davies, K.W., Holbrook, R., Lappin-Scott, H.M. & Humphrey, T.J. 1997. The use of an automated growth analyser to measure recovery times of single heat injured *Salmonella* cells. *Journal of Applied Microbiology*, **83**: 445–455.
- Stiles, M.E. 1996. Biopreservation by lactic acid bacteria. *Antonie van Leeuwenhoek*, **70**: 331–345.
- Tienungoon, S. 1998. Some aspects of the ecology of *Listeria monocytogenes* in salmonid aquaculture. Ph.D. Thesis, University of Tasmania, Hobart, Tasmania, Australia.
- Tienungoon, S., Ratkowsky, D.A., McMeekin, T.A. & Ross, T. 2000. Growth limits of *Listeria monocytogenes* as a function of temperature, pH, NaCl and lactic acid. *Applied and Environmental Microbiology*, **66**: 4979–4987.
- Truelstrup Hansen, L., Drewes Røntved, S. & Huss, H.H. 1998. Microbiological quality and shelf life of cold-smoked salmon from three different processing plants. *Food Microbiology*, **15**: 137–150.
- Truelstrup Hansen, L., Gill, T. & Huss, H.H. 1995. Effects of salt and storage temperature on chemical, microbiological and sensory changes in cold-smoked salmon. *Food Research International*, **28**: 123–130.
- Walsh, P. 1999. Consultancy for Researching the Business Profile of the NSW Seafood Industry and Food Safety Hazards of Seafood in NSW. Safe Food Production NSW, Sydney, NSW. 151p.



## FAO/WHO MICROBIOLOGICAL RISK ASSESSMENT SERIES

- 1 Risk assessments of *Salmonella* in eggs and broiler chickens: Interpretative Summary. 2002
- 2 Risk assessments of *Salmonella* in eggs and broiler chickens. 2002
- 3 Hazard characterization for pathogens in food and water: Guidelines. 2003
- 4 Risk assessment of *Listeria monocytogenes* in ready-to-eat foods: Interpretative Summary. 2004
- 5 Risk assessment of *Listeria monocytogenes* in ready-to-eat foods: Technical Report. 2004
- 6 *Enterobacter sakazakii* and micro-organisms in powdered infant formula: Meeting Report. 2004