

Basic statistics

Indicators	Statistics	Year
Population (thousands)	253	2013
Population aged under 15 (%)	37	2013
Population aged over 60 (%)	6	2013
Median age (years)	22	2013
Population living in urban areas (%)	26	2013
Total fertility rate (per woman)	3.4	2013
Number of live births (thousands)	6.7	2013
Number of deaths (thousands)	1.1	2013
Birth registration coverage (%)	43	2013
Cause-of-death registration coverage (%)	...	
Gross national income per capita (PPP int \$)	2840	2013
WHO region	Western Pacific	2013
World Bank income classification	Lower middle	2013

... Data from 2007 onwards not available.

Source:
Country statistics and global health estimates
by WHO and UN partners

For more information visit the Global Health Observatory
(<http://www.who.int/gho/en/>)


Last updated: January 2015

Life expectancy (years), 2012

		Country	WHO region	World Bank income group
Life expectancy	At birth	72	76	66
	At age 60	18	21	17
Healthy life expectancy	At birth	62	68	57

Life expectancy at birth for both sexes increased by 3 year(s) over the period of 2000–2012; the WHO region average increased by 4 year(s) in the same period.


In 2012, healthy expectancy in both sexes was 10 year(s) lower than overall life expectancy at birth. This lost healthy life expectancy represents 10 equivalent year(s) of full health lost through years lived with morbidity and disability.



■ WHO regional life expectancy at birth
■ Healthy life expectancy at birth
■ Lost healthy life expectancy

Millennium Development Goals (MDGs)


Indicators	Statistics	
	Baseline*	Latest**
Under-five mortality rate (per 1000 live births)	33	17
Maternal mortality ratio (per 100 000 live births)	170	86
Deaths due to tuberculosis among HIV-negative people (per 100 000 population)	17	6.3




*1990 for under-five mortality and maternal mortality; 2000 for other indicators
**2012 for deaths due to HIV/AIDS and malaria ; 2013 for other indicators

■ Country
■ WHO region


Distribution of causes of deaths in children under-5, 2013



DTP3 immunization among 1-year-olds



Children aged under-5 stunted




Country
WHO region


Source: Country statistics and global health estimates by WHO and UN partners
For more information visit the Global Health Observatory (<http://www.who.int/gho/en/>)
Last updated: January 2015

Utilisation of health services*

*Data refer to the latest year available from 2007.




Per capita total expenditure on health




... Data not available or applicable.

Adult risk factors



Population using improved water and sanitation



+ Improved drinking-water sources
X Improved sanitation facilities