



### Control programs for multiresistant *Staphylococcus aureus* (MRSA)

	Able to calculate the proportion of MRSA among all <i>S aureus</i> isolates
Western Europe	25/43 (58%)
Eastern Europe	13/27 (48%)
Africa	1/6 (17%)
USA	1/5 (20%)
South America	4/6 (67%)

Richet et al. Infect Control Hospital Epi 2003; 24: 334-341

### The important role of sentinel hospitals

- Centralization of available laboratory resources in a few selected centers
- Monitoring and reporting of AB susceptibility data (*WHOnet*)
- Adapt empiric treatment regimens


Archibald LK & Reller LB. *Clinical Microbiology in Developing Countries.* Emerg Infect Dis 2001; 7: 302-305  
Harbarth, SPCI-HUG

### Improve antibiotic use

- Monitor and provide feedback on occurrence and impact of AMR
- Optimize choice and duration of empiric antimicrobial therapy

Harbarth, SPCI-HUG

### Survival Among 401 Patients with Nosocomial Pneumonia Assigned to Short (8 d) or Long (15 d) Antimicrobial Treatment




JAMA 2003; 290: 2588-98

Harbarth, SPCI-HUG

Courtesy: J. Chastre, Paris

### Emergence of multiresistant pathogens for patients who had pulmonary infection recurrence



JAMA 2003; 290: 2588-98

Harbarth, SPCI-HUG

Courtesy: J. Chastre, Paris

### BMJ Three day versus five day treatment with amoxicillin for non-severe pneumonia in young children: a multicentre randomised controlled trial

BMJ 2004; 328:791; originally published online 16 Mar 2004; doi:10.1136/bmj.38049.490255.DE

ARTICLES

© Clinical efficacy of 3 days versus 5 days of oral amoxicillin for treatment of childhood pneumonia: a multicentre double-blind trial

Pakistan Multicentre Amoxicillin Short Course Therapy (MASCOT) pneumonia study group\*

Short-course versus long-course antibiotic therapy for non-severe community-acquired pneumonia in children aged 2 months to 59 months (Review)

Haider BA, Saeed MA, Bhutta ZA

### Non-severe CAP in children - Conclusions of recent review -

- Most episodes of pneumonia can be treated for a short duration
- Ambulatory non-severe pneumonia can be treated with 3 d of oral antibiotics
- Shorter course results in lower prevalence of resistant organisms

Grant GB et al. Lancet Infect Dis. 2009 Mar;9(3):185-96  
Harbarth, SPCI-HUG

### Reduced antimicrobial R

Carriage of cotrimoxazole non-susceptible organisms 2-4 weeks after antibiotic therapy

- 5 vs 10 days (Schrage et al JAMA 2001) → *S. pneumoniae* 34% vs 44%
- 3 vs 5 days (ISCAP BMJ 2004) → *H. influenzae* 57% vs 61%  
*S. pneumoniae* 67% vs 78%
- 3 vs 5 days (SCC WHO report 2003) → *H. influenzae* 54% vs 62%  
*S. pneumoniae* 62% vs 64%

Harbarth, SPCI-HUG

### Improve antibiotic use

- Monitor and provide feedback on occurrence of AMR
- Optimize choice and duration of empiric antimicrobial therapy
- Optimize perioperative antimicrobial prophylaxis

Harbarth, SPCI-HUG

### Antibiotic Prophylaxis and the Risk of Surgical Site Infections following Total Hip Arthroplasty: Timely Administration Is the Most Important Factor

M.E.E. van Kasteren, J. Manniën, A. Ott, B.J. Kullberg, A.S. de Boer, I.C. Gyssens. Clin Infect Dis 2007;44(7):921-7

Harbarth, SPCI-HUG

### Common Misconceptions in Surgical Prophylaxis

- Broad-spectrum is better
- Longer antibiotic prophylaxis is better
- Prophylaxis should be continued until all "tubes" are out

Harbarth, SPCI-HUG

### Misuse of prophylactic antibiotics in a university hospital, China

80% of prophylactic antibiotics (191/239) were started after the end of the operation

Stapin H et al. J Infect 2003; 46:161-63

### Overuse of prophylactic antibiotics in a community hospital, Saudi Arabia -- representative cases --

Procedure	Prophylactic antibiotics administered
Delivery	Ampicillin, amikacin, cefotaxime
Urinary cath	Amoxicillin, metronidazole
C-section	Cephadrine, ceftriaxone, gentamicin, metronidazole
Appendectomy	Cephadrine, ceftoxitin, amikacin, metronidazole, TMP-SMX
Cystoscopy	Amikacin, tetracycline, ceftazidime, amoxicillin-clav
Cholecystectomy	Cephadrine, cefuroxime, gentamicin
Incision	Ampicillin, amikacin, amoxicillin-clav, cephradine
Episiotomy	Amoxicillin, gentamicin, cephradine, metronidazole
Delivery	Amoxicillin, gentamicin, cephradine, metronidazole

Al-Ghamdi S et al. J Hosp Infect 2002; 50:115-21

### Duration of surgical prophylaxis and selection of resistance

**Cardiovascular surgery**  
n= 2'641, multivariate analysis



	< 48 h prophylaxis	> 48 h prophylaxis
	OR (95%CI)	P
	> 48 h prophylaxis	
<b>SSI</b>	<b>1.0 (0.8-1.3)</b>	<b>ns</b>
<b>Resistant Enterobacteriaceae/enterococci</b>	<b>1.7 (1.1-2.7)</b>	<b>0.027</b>

Harbarth et al. Prolonged antibiotic prophylaxis after cardiovascular surgery and its effect on surgical site infections and antimicrobial resistance. Circulation 2000;101:2916 - 2921  
Harbarth, SPCI-HUG


### Improve perioperative antibiotic prophylaxis (ABP)

Process	Problem area	System changes
ABP choice appropriate	- Suboptimal - Coverage too large	- Guidelines - Adequate supply
Duration adequate	Too long (>1 dose)	- Standard order form
Timing correct	Too early or too late	- Administer in preoperative area - Designate responsible person

Harbarth, SPCI-HUG  
Huskins et al. Infect Control Hosp Epidemiol 1998;19:125-35







- ### Improve antibiotic use (2)
- **Decrease diagnostic uncertainty:**
    - Improve diagnostic tools
    - Promote use of clinical algorithms
- Harbarth, SPCI-HUG




### Use of clinical algorithms

- Prediction of bacteremia & mortality in hospitalized Malawian children
  - Association with lethargy, oral thrush, chronic cough and malnutrition
- The WHO Young Infants Study Group
  - Clinical prediction rule to identify and treat serious bacterial infection

Norton EB et al. Pediatr Infect Dis J 2004; 23: 145-51  
WHO study group. Pediatr Infect Dis J 1999; 18: S23-31



**Use of Procalcitonin to Shorten Antibiotic Exposure in ICU Patients : The ProRata Trial**



Harbarth, Bouadma et al. Lancet 2010

**Improve antibiotic use (2)**

- **Decrease diagnostic uncertainty**
- **Implement formulary restrictions for important types of antimicrobial use**

Harbarth, SPCI-HUG

**Impact of an antibiotic restriction policy on hospital expenditures and bacterial susceptibilities: a lesson from a pediatric institution in a developing country**

Variable	Pre-intervention period (1995/96)	Intervention period (1997/98)
Vials (#)	199,427	132,496
Total costs (\$)	699,543	347,261

-- Stable or decreasing resistance rates --

Saez-Llorens et al. Ped Infect Dis J 2000; 19: 200-6

**Does restriction always work?**

Formulary restriction at Mass Gen Hosp, Boston (USA) :

“ Imipenem, tic/clav, aztreonam, cefta, cipro, pip/tazo require prior approval by infectious diseases “

Gilbert et al. Am J Med; 1998; 104: 17-27

The reality at the same hospital ....

35-y old woman with severe sepsis:  
 “ Ampicillin-sulb, clindamycin, penicillin, gentamicin, vancomycin were infused intravenously “

Case report 28-2002 of the MGH, NEJM Sept 12, 2002, p.831-37  
 Harbarth, SPCI-HUG

**Improve antibiotic use (2)**

- **Improve diagnostic tools**
- **Implement formulary restrictions for important types of antimicrobial use**
- **Improve antimicrobial prescribing:**
  - **Education (pre- and postgraduate)**
  - **Practice guidelines**
  - **Administrative means (antibiotic order forms)**
  - **Feedback to prescribers**

Harbarth, SPCI-HUG

**Implementing practice guidelines for appropriate AB use: Systematic review**

- 40 studies (in- and outpatient areas)
- Multifaceted implementation methods were most successful
- Most useful implementation methods:
  - Locally adapted guidelines (drug committee)
  - Small-group interactive sessions
  - Academic detailing
  - Participation of opinion leaders
  - Feedback to prescribers

Gross PA et al. Med Care 2001; 39: Suppl 55-69  
 Harbarth, SPCI-HUG

### Impact of an educational program on antibiotic use in a tertiary care hospital in Thailand

---

#### Appropriate antibiotic use (in-patients, %)

Variable	Preintervention period (n = 4305)	Postintervention period (n = 2830)	P
Inappropriate antibiotic use	1808 (42)	566 (20)	<.001
Reason for inappropriateness <sup>a</sup>			
Inappropriate surgical prophylaxis <sup>b</sup>	452 (25)	115 (20)	.02
Use of antibiotic without any evidence of infection	723 (40)	200 (35)	.04
Redundant spectrum	217 (12)	50 (9)	.03
Bacterial resistance <sup>c</sup>	235 (13)	91 (16)	.07
Narrow spectrum was available <sup>d</sup>	181 (10)	41 (7)	.04
Department <sup>e</sup>			
Surgery	633 (35)	170 (30)	.01
Obstetrics and gynecology	452 (25)	125 (22)	.17
Internal medicine	416 (23)	113 (20)	.14
Other <sup>f</sup>	307 (17)	113 (20)	.12

Apisarnkarn et al. Clin Infect Dis 2006; 42: 768

### Impact of an educational program on antibiotic use in a tertiary care hospital in Thailand

---

#### Antibiotic resistance


Microorganism	Resistance rate, % <sup>a</sup>	
	Preintervention period	Postintervention period
Methicillin-resistant <i>Staphylococcus aureus</i>	48	33.5
ESBL-producing <i>Escherichia coli</i>	...	...
ESBL-producing <i>Klebsiella pneumoniae</i>	33	21
Third-generation cephalosporin-resistant <i>Acinetobacter baumannii</i>	30	20
Impipenem-resistant <i>Pseudomonas aeruginosa</i>	27	19
Multidrug-resistant <i>Acinetobacter baumannii</i>	5	4
	4	5


Apisarnkarn et al. Clin Infect Dis 2006; 42: 768

- ### Interventions to improve quality of antibiotic prescribing for hospital inpatients (review)
- Davey P, Brown E, Fenelon L, et al.  
Cochrane Database of Systematic Reviews 2005; Issue 4. Art.No CD003543.
- 51/66 studies showed a significant improvement in at least one outcome
    - Reduction of costs, AMR or HCAI
  - Interventions to improve antibiotic prescribing in inpatients likely to be successful
  - Absence of good evidence which interventions are most cost-effective in reducing AMR
- Davey P, et al. Cochrane Database of Systematic Reviews 2005; Issue 4. Art.No CD003543.

## Outpatient setting


Macro-level determinants





Social Science & Medicine 57 (2003) 733–744

www.elsevier.com/locate/socscimed





### Improving antibiotic use in low-income countries: an overview of evidence on determinants

Aryanti Radyowijati, Hilbrand Haak\*

Consultants for Health and Development, Skeddoornuin 7, 2317 MV Leiden, The Netherlands



Harbarth, SPIC-HUG




### Available in a pharmacy in Delhi – over-the-counter without prescription!

Harbarth, SPCT-HUG

### Country examples: Possible interventions






### Action plan

- In 1999, Chile decided an intervention to:
  - educate physicians & public
  - regulate the consumption of antibiotics
  - restrict over-the-counter antibiotic sales


Harbarth, SPCI-HUG



### Sales of oral antibiotics in US\$ (Million)

Year	1996	1997	1998	1999	2000	2001	2002
USD	37,6	45,8	45,8	38,9	32,1	29,4	26,1
				- 15%	- 30%	- 36%	- 43%

Harbarth, SPCI-HUG








## Korea- Government Policy

- A new Korean government policy announced in 2000 prohibited doctors from dispensing and pharmacists from prescribing drugs by law.



Harbarth, SPCI-HUG



## Policy priorities: AB use

---

**Local level**

- Improve perioperative prophylaxis
- Promote short-course, high-dose AB therapy
- Decrease diagnostic uncertainty by any type of diagnostic tools or decision support
- Promote local guidelines and drug committees

Harbarth, SPCI-HUG

## Policy priorities: AB use

---

**National level**

- Create sentinel laboratories for surveillance of antibiotic resistance
- Change consumer expectations
- Implement healthcare regulation for the prudent use of antibiotics
- Control marketing activities of industry

Harbarth, SPCI-HUG

## Summary of Measures

---

**Measures**

<ul style="list-style-type: none"> <li>■ Improve antibiotic prophylaxis</li> <li>■ Clinical algorithms</li> </ul>	<ul style="list-style-type: none"> <li>■ Surveillance</li> <li>■ Microbiologic support</li> <li>■ Restriction &amp; education</li> </ul>	<ul style="list-style-type: none"> <li>■ Decision support systems</li> <li>■ New diagnostic markers</li> <li>■ Academic detailing</li> </ul>
---	--	--

Harbarth, SPCI-HUG

“The development of new antibiotics without having mechanisms to insure their appropriate use is much like supplying your alcoholic patients with a finer brandy.”

Dennis Maki 1998

Harbarth, SPCI-HUG

Third International Conference for Improving Use of Medicines  
 Bibliotheca Alexandrina, Alexandria, Egypt • April 10-14, 2011  
*Informed Strategies, Effective Policies, Lasting Solutions*

**ICIUM 2011**

PLACE IT ON YOUR CALENDARS & IN YOUR DIARIES NOW!

ICIUM2011 will be as unique as its two predecessors in 1997 and 2004. [www.icium.org](http://www.icium.org)

**Participants:** will help to shape evidence-based policy recommendations, implementation strategies and a future research agenda.

**Sessions:** will be organized around different levels of the health care system, including: global, national, institutional, health care providers and patients.

**Topic tracks:** Access, policy, regulation, governance, economics, financing, insurance systems, maternal and child health, chronic care, HIV/AIDS, tuberculosis, malaria, drug resistance\*.

Abstracts will need to be submitted before Dec 7<sup>th</sup> 2010