



## Highlights: Week 5 (30-5 Feb 2012)

- 285 typhoid cases reported this week
- Increase in malaria cases at national level

### Contents


- A. General context
- B. Epidemic prone diseases
- C. Events of public health importance in the region
- D. Completeness and timeliness of the national data
- E. Acknowledgments
- F. Annexes
  1. Classification of events that may constitute a Public Event of International Concern
  2. Standard case definitions
  3. Alert/action epidemic thresholds

**Table 1:** Distribution of typhoid cases in Harare, 30-Jan - 5 Feb 2012

Suburbs	Frequency n= 285 (%)
Kuwadzana	158 (55.4)
Dzivarasekwa	20 (7)
Crowborough North	10 (3.5)
Glen View	8 (2.8)
Other areas	87 (31.2)

This week other cases have been reported from other areas outside Harare namely Chitungwiza (2); Bindura (1) in Mash Central Province; Chivero (3), Grenara (5), Denview (3) and Snake Park (4) in Mash West Province.

**Figure 1:** Harare Typhoid Epidemic Curve, 10 Oct 2011 - 29 Jan 2012



## A. General Context

On-going typhoid outbreak reported in Harare City since 10 October 2011. Other cases have since been reported in other parts of the country.

No report of outbreaks cholera and measles this week countrywide.

No report of Pandemic Influenza A cases this week.

## B. Epidemic prone diseases

### Typhoid outbreak in Harare City

Since 10 October 2011 Harare City has been experiencing an outbreak of Typhoid Fever. As of 5 February 2012, a total of 2697 cases have been reported of which 60 were confirmed by laboratory diagnosis. From 1 January 2012 to 9 February 2012, a total 1605 cases were reported. The most affected suburbs were Kuwadzana and Dzivarasekwa. The distribution of cumulative cases to date by place of residence is shown in the map in Figure 2.


### Response activities

- Coordination meetings with the thematic committees and Partners
- Social mobilization activities
- Investigation and contact tracing
- Water quality monitoring
- Case management
  - BRIDH
  - Kuwadzana polyclinic
  - Dzivarasekwa Rujeko polyclinic

# Zimbabwe Weekly Epidemiological Bulletin

- Mobile clinic in Granary (Zvimba District in Mashonaland West Province)

Figure 2: Distribution of typhoid cases by place of residence, Harare City, 10 Oct 2011- 5 Feb 2012




## Anthrax

No case of anthrax reported this week from Gokwe North. The national cumulative cases as of 29 January were six.

## Malaria

This week, 9733 cases and 9 deaths were reported. The most affected areas were Makoni, Mutasa 1321 (15%), Mutare 902 (11%), Chimanimani 843(10%), Mutoko 575 (7%) and Makonde 520(6%). Since the beginning of 2012, a total of 33532 cases and 29 deaths (CFR= 0.09%) were reported.

Figure 3: Comparison of national malaria cases 2012 vs 2011 and epidemic threshold.



## C. Completeness and timeliness of the National data

National data completeness and timeliness reported this week n° 5 increased from 70% to 73% and from 50% to 55% respectively

## D. Events of public health importance within SADC

### Cholera

In Angola, a total of 400 case of which 26 deaths were reported (CFR=0.065%).

### Anthrax

No update was received on the anthrax situation.

## E. Acknowledgements

All health workers, operating at different levels of the health system, providing information are greatly acknowledged. In addition, special thanks to the members of Health and WASH clusters for sharing their data with our team.

MOHCW is grateful to all Partners including UN family and NGOs for their support.

Information on events of public health importance occurring within SADC is consolidated from the WHO daily summary of health events.

# Zimbabwe Weekly Epidemiological Bulletin

## **Annex 1: Classification of Events that may constitute a Public Health Emergency of International Concern**

There are three groups of events if detected by the national surveillance system should trigger the use of the IHR (2005) Decision Instrument to be notified as they may constitute Public Health Emergencies of International Concern. These are:

1. A case of unusual or unexpected diseases which may have serious public health impact: smallpox, poliomyelitis due to wild-type poliovirus, human influenza caused by a new subtype and SARS.
2. Any event of potential international public health concern including events of unknown causes or sources and those involving other events or diseases:
  - environmental health emergencies (natural events, chemical and radio-nuclear events, technological incidents, complex emergencies and deliberate events)
  - Food borne diseases
  - Zoonotic diseases or other infectious diseases.
3. Any of following diseases that have demonstrated the ability to cause serious public health impact and spread rapidly and internationally: Cholera, pneumonic plague, yellow fever, viral haemorrhagic fevers, West Nile Fever, other diseases that are of special national or regional concern e.g. dengue, RVF and meningococcal disease.

# Zimbabwe Weekly Epidemiological Bulletin

## Annex 2: Standard Case Definitions

Diseases	Standard Case Definitions
Cholera	<p><b>Suspected case</b></p> <ul style="list-style-type: none"> <li>In an area where there is no cholera outbreak, any person aged five years or more, presenting with severe dehydration or death from acute watery diarrhoea</li> <li>In an area where there is a cholera outbreak, any person aged two years or more presenting with acute watery diarrhoea, with or without vomiting</li> </ul>
	<p><b>Confirmed case</b> A suspected case in which <i>Vibrio cholerae</i> sero-groups O1 or O139 has been isolated in the stool.</p>
	<p><b>Note</b></p> <ul style="list-style-type: none"> <li>All suspected cases under the age of two years must be confirmed.</li> <li>The inclusion of all ages in the case definition somewhat reduces specificity, that is, inclusion of more non-cholera childhood diarrhoea cases (mainly those below 5years). It does not impede meaningful interpretation of trends. Teams should monitor any shift in the age distribution of cases, which might indicate a changing proportion of non-cholera cases among patients seen.</li> </ul>
Malaria	<p><b>Suspected uncomplicated malaria</b> Any person living in a malaria area or history of travelling in a malaria area within the last 6 weeks, presenting with fever, malaise, chills, and rigors, without signs of severe disease such as vital organ dysfunction</p>
	<p><b>Confirmed uncomplicated malaria</b> Is suspected uncomplicated malaria with laboratory diagnosis by malaria blood slide or RDT for malaria parasites</p>
	<p><b>Confirmed severe malaria</b> A patient hospitalized with <i>P. falciparum</i> asexual parasitaemia as confirmed by laboratory tests with accompanying symptoms of severe disease (vital organ dysfunction)</p>
Typhoid	<p><b>Suspected case</b> Any person with gradual onset of steadily increasing and then persistently high fever, chills, malaise, headache, sore throat, cough, and, sometimes, abdominal pain and constipation or diarrhoea</p>
	<p><b>Confirmed case</b> A suspected case confirmed by isolation of <i>Salmonella typhi</i> from blood, bone marrow, bowel fluid or stool</p>
Diarrhoea	<p><b>Suspected case</b> Passage of 3 or more loose or watery stools in the past 24 hours with</p> <ul style="list-style-type: none"> <li>or without dehydration <b>or</b></li> <li>some dehydration and two or more of the following signs: restlessness, irritability, sunken eyes, thirsty, skin pinch goes back slowly, <b>or</b></li> <li>severe dehydration and two or more of the following signs: lethargy or unconsciousness; sunken eyes; not able to drink or drinking poorly; skin pinch goes back very slowly</li> </ul>
	<p><b>Confirmed case</b> Suspected case confirmed with stool culture for a known enteric pathogen.</p>
	<p><b>Note</b> <i>Laboratory confirmation of specific agent causing outbreak is not routinely recommended for surveillance purposes.</i></p>

# Zimbabwe Weekly Epidemiological Bulletin

## Annex 3: Alert/Action Epidemic Thresholds for selected epidemic prone diseases and other diseases of public health importance in Zimbabwe

Disease or condition	Alert Threshold	Action Threshold
Measles	5 suspected cases within a district in a month	1 measles IgM confirmed case
		<i>Note: This also applies to closed settings like Refugee camps, schools, or health facilities</i>
Meningococcal meningitis	1 suspected case	1 confirmed case
Plague	1 suspected case	1 confirmed case
Rabies (suspected rabid bites)	1 case of a bite from suspected rabid animal	1 case of a bite from suspected rabid animal
Trypanosomiasis	1 suspected case	<ul style="list-style-type: none"> <li>1 case in an area that is not endemic <b>or</b></li> <li>For endemic areas 3 cases per 100,000</li> </ul>
Typhoid fever	1 case	<ul style="list-style-type: none"> <li>5 suspected cases per 50,000 population <b>or</b></li> <li>20 suspected cases per District's catchment area <b>or</b></li> <li>any 1 confirmed case by blood culture</li> </ul>
Viral Haemorrhagic Fever	1 suspected case	1 confirmed case
Outbreak of unknown cause	3-5 cases or deaths with similar symptoms that don't fit most case definitions	Any cluster of cases or deaths that had similar symptoms over a short period of time and fail to respond to treatment for the usual causes of the symptoms
Acute Flaccid paralysis (AFP) / Polio	1 AFP case	1 confirmed case of polio (virus isolated).
Dysentery	5 cases or more per reporting site per week	<ul style="list-style-type: none"> <li>A 2-fold increase in the number of cases compared to an expected number usually seen in previous season – specific time period</li> <li>Any increase in number of deaths due to bloody diarrhoea</li> </ul>
Cholera	1 suspected case	1 confirmed case (where it has not been reported before)
Diarrhoea under five	Increasing number of cases in a short time	Doubling of no of cases as compared to the same time period of a previous year.
Malaria	Increasing cases above the median	<ul style="list-style-type: none"> <li>N<sup>o</sup> of cases that exceed those in the 3<sup>rd</sup> quartile (the upper limit) of the expected number of cases <b>or</b></li> <li>N<sup>o</sup> of cases that exceed the mean plus 1.5 x Standard Deviations (Mean + 1.5 SD).</li> </ul>
Neonatal Tetanus (NNT)	1 suspected case	1 confirmed case
Human influenza caused by a new Subtype	1 suspected case	1 confirmed case
Severe Acute Respiratory Syndrome (SARS)	1 suspected case	1 confirmed case
Adverse Events Following Immunisation (AEFI)	1 suspected case	1 confirmed case
Acute Viral Hepatitis	1 suspected case	1 confirmed case
Anthrax	1 suspected case	1 confirmed case

### Notes

# Zimbabwe Weekly Epidemiological Bulletin

An *alert threshold* suggests to health workers that further investigation is needed. Health workers respond to an alert threshold by:

- Reporting the suspected problem to the next level
- Reviewing data from the past
- Requesting laboratory confirmation to see if the problem is one that fits a case definition
- Being more alert to new data and the resulting trends in the disease or condition
- Investigating the case or condition
- Alerting the appropriate disease-specific programme manager and district epidemic response team to a potential problem.

An *epidemic/action threshold* triggers a definite response. Possible actions include communicating laboratory confirmation to affected health centres, implementing an emergency response, community awareness campaign, or improved infection control practices in the health care setting.

## Reporting

- T1 for notification of an infectious notifiable disease (used for up to five cases after which line lists must be filled)
- Weekly Rapid Disease Notification Form
- Reporting is to the next level (health facility to district to province to national level)