

Measles vaccines: WHO position paper - 28 August 2009

Grading of scientific evidence in support of key recommendations

Table II. Duration of protection following measles immunization

Question: Is there a substantial decrease in protection against clinical measles with increasing time following immunization?

Settings: Global

Conclusion: The protective immunity against clinical measles does not seem to be substantially decreased with increasing time following immunization (Low level of scientific evidence)

Quality assessment						Summary of Findings	Importance
No of studies	Design	Limitations	Inconsistency	Indirectness	Imprecision	Quality	
Evidence of vaccine-induced protection against measles following exposure to the disease							
11	Observational	No serious	No serious ¹	No serious	No serious	Low	High

¹Except for the reports by Hutchins SS et al and Yuan L the majority of these studies show consistent results

Long-term protection against clinical measles

Outbreak studies from many countries have failed to identify declining immunity as an important risk factor. Investigation of measles attack rates in 72 RMI households during a major measles outbreak in the Marshall Islands in 2003 showed similar and high vaccine-induced protection rates among children from 6 months to 14 years of age (*Marin M et al 2006*). Similarly, among 78 measles contacts in the islands of Palau, where no measles had occurred for 27 years, time since vaccination was not a significant risk factor for developing measles: RR1.6; (95% CI 0.3-9.4) for vaccination >15 versus <5 years ago (*Güriş D et al, 1996*). The investigation of a major measles epidemic in Romania 1996-98 concluded that vaccine induced protection was similarly high among 2,561 children vaccinated 6-8, 9-11, or 12-14 years previously (*Hennessey KA et al 1999*).

A follow-up of 4,500 trial participants in England and Wales 21 years after receiving one dose of measles vaccine showed continuing high level of protection compared with unvaccinated controls; there was no indication of waning immunity in the vaccinated group (*Miller C, 1987*). Subsequent follow-up of this trial population suggests that the protection from live measles vaccine persists unchanged for up to 27 years after vaccination (*Ramsay ME, 1994*).

A number of studies of measles outbreaks in teaching institutions have concluded that the risk of clinical measles was not associated with length of time since vaccination: *Sutcliffe PA et al, 1996*, investigated a measles outbreak among 1135 students 14-21 years of age in Toronto; *Nkowane BM et al, 1987*, analysed the attack rate in relation to number of years after vaccination in 197 individuals during a measles outbreak in a highly vaccinated high-school population in Massachusetts; *Faust HS et al, 1983*, studied a measles outbreak involving 250 school-aged individuals in Michigan; and *Hersh BS et al 1991* studied an outbreak involving 86 cases at a college in Colorado.

On the other hand, a retrospective cohort study of single-dose vaccinees in one school in New Mexico found that persons vaccinated ≥ 10 years before the outbreak were at increased risk, independently of age at vaccination (*Hutchins SS et al 1990*). Also, Yuan L, 1994, who investigated a measles outbreak involving 87 children in 31 Canadian schools concluded that subjects vaccinated before 1980 were at greater risk of contracting the disease than those vaccinated after 1980 (adjusted OR 14.5, 95% CI 1.5 to 135.6).

References

- Anders JF, Jacobson RM, Poland GA, Jacobsen SJ, Wollan PC. Secondary failure rates of measles vaccines: a metaanalysis of published studies. [Pediatr Infect Dis J](#). 1996 Jan;15(1):62-6.
- Faust HS, Thompson FE. Age at and time since vaccination during a measles outbreak in a rural community. [Am J Dis Child](#). 1983 Oct;137(10):977-80.
- Güriş D, McCready J, Watson JC, Atkinson WL, Heath JL, Bellini WJ, Polloi A. Measles vaccine effectiveness and duration of vaccine-induced immunity in the absence of boosting from exposure to measles virus. [Pediatr Infect Dis J](#). 1996 Dec;15(12):1082-6
- Hennessey KA, Ion-Nedelcu N, Craciun MD, Toma F, Wattigney W, Strebel PM. Measles epidemic in Romania, 1996-1998: assessment of vaccine effectiveness by case-control and cohort studies. [Am J Epidemiol](#). 1999 Dec 1;150(11):1250-7.
- Hersh BS, Markowitz LE, Hoffman RE, Hoff DR, Doran MJ, Fleishman JC, Preblud SR, Orenstein WA. A measles outbreak at a college with a prematriculation immunization requirement. [Am J Public Health](#). 1991 Mar;81(3):360-4.
- Hutchins SS, Markowitz LE, Mead P, Mixon D, Sheline J, Greenberg N, Preblud SR, Orenstein WA, Hull HF. A school-based measles outbreak: the effect of a selective revaccination policy and risk factors for vaccine failure. [Am J Epidemiol](#). 1990 Jul;132(1):157-68
- Nkowane BM, Bart SW, Orenstein WA, Baltier M. Measles outbreak in a vaccinated school population: epidemiology, chains of transmission and the role of vaccine failures. [Am J Public Health](#). 1987 Apr;77(4):434-8.
- Marin M, Nguyen HQ, Langidrik JR, Edwards R, Briand K, Papania MJ, Seward JF, LeBaron CW. Measles transmission and vaccine effectiveness during a large outbreak on a densely populated island: implications for vaccination policy. [Clin Infect Dis](#). 2006 Feb 1;42(3):315-9. Miller C. Live measles vaccine: a 21 year follow up. [Br Med J \(Clin Res Ed\)](#). 1987 Jul 4;295(6589):22-4.
- Ramsay ME, Moffatt D, O'Connor M. Measles vaccine: a 27-year follow-up. [Epidemiol Infect](#). 1994 Apr;112(2):409-12.
- Sutcliffe PA, Rea E. Outbreak of measles in a highly vaccinated secondary school population. [CMAJ](#). 1996 Nov 15;155(10):1407-13.
- Yuan L. Measles outbreak in 31 schools: risk factors for vaccine failure and evaluation of a selective revaccination strategy. [CMAJ](#). 1994 Apr 1;150(7):1093-8.