

## PREVENTING DISEASE THROUGH HEALTHY ENVIRONMENTS

### EXPOSURE TO LEAD: A MAJOR PUBLIC HEALTH CONCERN

Lead is a toxic metal whose widespread use has caused extensive environmental contamination and health problems in many parts of the world. It is a cumulative toxicant that affects multiple body systems, including the neurological, haematological, gastrointestinal, cardiovascular and renal systems. Children are particularly vulnerable to the neurotoxic effects of lead, and even relatively low levels of exposure can cause serious and, in some cases, irreversible neurological damage.<sup>1,2</sup> Lead exposure is estimated to account for 0.6% of the global burden of disease, with the highest burden in developing regions.<sup>3</sup> Recent reductions in the use of lead in petrol (gasoline), paint, plumbing and solder have resulted in substantial reductions in lead levels in the blood.<sup>1</sup> However, significant sources of exposure to lead still remain, particularly in developing countries. Further efforts are required to continue to reduce the use and releases of lead and to reduce environmental and occupational exposures, particularly for children and women of child-bearing age.

#### Sources of exposure to lead

Lead is found at low levels in Earth's crust, mainly as lead sulfide.<sup>4</sup> However, the widespread occurrence of lead in the environment is largely the result of human activity, such as mining, smelting, refining and informal recycling of lead; use of leaded petrol (gasoline); production of lead-acid batteries and paints; jewellery making, soldering, ceramics and leaded glass manufacture in informal and cottage (home-based) industries; electronic waste<sup>5</sup>; and use in water pipes and solder. Other sources of lead in the environment include natural activities, such as volcanic activity, geochemical weathering and sea spray emissions, and remobilization of historic sources, such as lead in soil, sediment and water from mining areas.

As lead is an element, once it is released into the environment, it persists.<sup>2</sup> Because of lead's persistence and potential for global atmospheric transport, atmospheric emissions affect even the most remote regions of the world.<sup>6</sup>

#### ***Industrial processes***<sup>1,2</sup>

Lead is used mainly in the production of lead-acid batteries, plumbing materials and alloys. Other uses are in cable sheathing, paints, glazes and ammunition. Human occupational exposure can also take place during the application and removal of protective lead-containing paints, during the grinding, welding and cutting of materials painted with lead-containing paints, such as in shipbuilding, construction, demolition industries, and fabrication of heavy lead glass and crystal, and in crystal carving. Mining, smelting, and informal processing and recycling of electric and electronic waste can also be significant sources of exposure. Lead has been used widely in the form of tetraethyl and tetramethyl lead as antiknock and lubricating agents in petrol, although the majority of lead is emitted from vehicles in the form of inorganic particles. This use has been phased out in most countries, which has resulted in a significant reduction of human exposure and mean blood lead levels. In the few parts of the world where leaded petrol is still in use, however, it continues to be a major source of exposure. Old industrial hotspots that have not been cleaned up can also represent a hazard

even years after contamination has stopped, particularly to children who might ingest contaminated soil or dust as a result of their hand-to-mouth behaviour.

### ***Food and smoking***

For the non-smoking general population, the largest contribution to the daily intake of lead is derived from the ingestion of food, dirt and dust. The amount of lead in food plants depends on soil concentrations and is highest around mines and smelters. Cereals can contain high levels of lead, and spices may be contaminated with lead. The use of lead-soldered food and beverage cans (which is now diminishing) may considerably increase the lead content of the food or beverage, especially in the case of acidic foods or drinks. As alcoholic drinks tend to be acidic, the use of any lead-containing products in their manufacture, distribution or storage will raise lead levels. Migration of lead into food from lead-glazed ceramic or pottery dinnerware is also a source of exposure. Smoking tobacco increases lead intake.<sup>2</sup>

### ***Drinking-water***

Lead present in tap water is rarely the result of its dissolution from natural sources but is mainly due to household plumbing systems containing lead pipes, solders and fittings. Water that has been in contact with lead in this way for an extended period (e.g. overnight) will have a higher concentration. Thus, lead concentrations can vary over the day, and flushing of the taps before use is a control mechanism. Soft acidic water dissolves the most lead.<sup>7</sup>

### ***Domestic sources***

Contaminated dust may be the main source of exposure for infants in countries that no longer use leaded petrol. The weathering, peeling or chipping of lead-based paints, mainly found in older houses, plays a role in children's exposure, especially as some young children eat the fragments or lick dust-laden fingers. Lead-containing dust may be brought into the home on the clothes of those who work in industries where such dust is generated. Some toys either are made from lead or contain lead (e.g. some plastics or paints).<sup>8</sup> Some traditional medicines and makeup (e.g. kohl) contain lead.<sup>2</sup>

## **World Health Organization (WHO) lead guidelines**

### **Tolerable intake level**

In a review of the latest scientific evidence, conducted in 2010, the Joint Food and Agriculture Organization of the United Nations (FAO)/WHO Expert Committee on Food Additives (JECFA) estimated that the previously established provisional tolerable weekly intake (PTWI) of 25 µg/kg body weight per week could no longer be considered health protective and withdrew it. As the dose–response analyses did not provide any indication of a threshold for the key adverse effects of lead, the Committee concluded that it was not possible to establish a new PTWI that would be health protective. The dose–response analyses conducted by the Committee should be used as guidance to identify the magnitude of effect associated with identified levels of dietary lead exposure in different populations.<sup>9</sup>

### **Drinking-water**

10 µg/l<sup>10</sup>

### **Air**

0.5 µg/m<sup>3</sup> (annual average)<sup>11</sup>

## Health effects

- It has been estimated that lead exposure was responsible, in 2004, for 143 000 deaths and 0.6% of the global burden of disease (expressed in disability-adjusted life years, or DALYs<sup>\*</sup>), taking into account mild mental retardation and cardiovascular outcomes resulting from exposure to lead.<sup>3</sup>
- Lead in the body is distributed to the brain, liver, kidney and bones. It is stored in the teeth and bones, where it accumulates over time. Human exposure can be assessed directly through measurement of lead in blood, teeth or bones (bone and tooth lead reflect cumulative exposure).<sup>14</sup>

### **Effects in children and pregnant women**

- Young children absorb 4–5 times as much lead as adults (apart from pregnant women). Infants, young children (especially those less than 5 years of age) and pregnant women are most susceptible to the adverse effects of lead.
- The potential for adverse effects of lead exposure is greater for children than for adults, because in children 1) the intake of lead per unit body weight is higher, 2) more dust may be ingested, 3) lead absorption in the gastrointestinal tract is higher, 4) the blood–brain barrier is not yet fully developed and 5) neurological effects occur at lower levels than in adults.<sup>15</sup>
- The most critical effect of lead in young children is that on the developing nervous system. Subtle effects on intelligence quotient (IQ) are expected from blood lead levels at least as low as 5 µg/dl (50 µg/l), and the effects gradually increase with increasing levels of lead in blood.<sup>1</sup> A 2010 review of the latest scientific evidence indicating effects at lower levels did not provide any indication of a threshold for the key adverse effects of lead.<sup>9</sup> Lead exposure has also been linked epidemiologically to attention deficit disorder and aggression.<sup>4</sup>
- Exposure of pregnant women to high levels of lead can cause miscarriage, stillbirth, premature birth and low birth weight, as well as minor malformations.<sup>2</sup>

### **Acute effects**

- Lead is classically a chronic or cumulative toxin; hence, acute adverse effects are usually observed only following short-term exposures to high concentrations. Acute exposures to lead may cause gastrointestinal disturbances (anorexia, nausea, vomiting, abdominal pain), hepatic and renal damage, hypertension and neurological effects (malaise, drowsiness, encephalopathy) that may lead to convulsions and death.<sup>2</sup>

### **Effects following chronic exposure**

- Chronic lead exposure commonly causes haematological effects, such as anaemia, or neurological disturbances, including headache, irritability, lethargy, convulsions, muscle weakness, ataxia, tremors and paralysis.<sup>9</sup>
- There is some evidence that long-term occupational exposure to lead may contribute to the development of cancer. The International Agency for Research on Cancer (IARC) has classified inorganic lead compounds as *probably carcinogenic to humans*

---

\* The DALY combines the burden due to death and disability in a single index. Use of such an index permits the comparison of the burden due to various environmental risk factors with those from other risk factors or diseases. One DALY can be thought of as 1 lost year of healthy life.<sup>12,13</sup>

(Group 2A), meaning that there is limited evidence for carcinogenicity in humans and sufficient evidence of carcinogenicity in experimental animals. According to IARC, organic lead compounds are *not classifiable as to their carcinogenicity to humans* (Group 3), meaning that there is inadequate evidence for their carcinogenicity to humans.<sup>4</sup>

- In men, the reproductive effects of lead include decreased sperm count and increased number of abnormal sperm.

## Risk mitigation recommendations

Primary prevention (i.e. the elimination of exposure to lead at its source) is the single most effective intervention against childhood lead poisoning. The following actions are needed:

### ***Eliminating use***

- Phase out the use of lead additives in motor fuels in countries where this has not yet been done.
- Phase out the use of lead in paints on a worldwide basis.
- Eliminate the use of leaded solder in food and drink cans, as well as in water pipes.
- Eliminate the use of lead in homes, schools, school materials and children's toys.
- Eliminate the use of lead glazing for pottery intended for cooking, eating or drinking.
- Encourage the removal of plumbing and fittings containing lead (as this is costly, other measures, such as corrosion control and minimizing the dissolving of lead in water systems, should be implemented in the meantime).
- Identify and eliminate lead use in traditional medicines and cosmetics.

### ***Preventing exposure***

- Prevent exposure to lead from electric and electronic waste (e.g. lead-acid batteries, computers), particularly for children.
- Ensure that the recycling of lead-containing waste is undertaken only in the presence of appropriate industrial hygiene measures and that informal recycling and use of lead-containing waste are discouraged.
- Identify contaminated sites, and take necessary action to prevent human exposure to lead from these areas.

### ***Monitoring***

- Monitor blood lead levels during occupational exposure (preferably supplemented by bioanalytical measures of exposure, such as zinc protoporphyrin levels and 5-aminolevulinate dehydratase activities).
- Extend the monitoring of blood lead levels in children and women of child-bearing age by use of sensitive analytical methods.
- Enhance the collection of data on lead in foodstuffs, and make this information publicly available so that appropriate action can be taken; identify foodstuffs with high lead content and the sources of lead contamination, and use the information to support appropriate action.

## Education

- Educate the public regarding the dangers of misusing lead-containing products.
- Promote preventive and educational measures to protect young children from lead in their environment.

## References

1. Fewtrell L, Kaufmann R, Prüss-Üstün A (2003). *Lead: Assessing the environmental burden of disease at national and local levels*. Geneva, World Health Organization (Environmental Burden of Disease Series, No. 2; [http://www.who.int/quantifying\\_ehimpacts/publications/en/leadebd2.pdf](http://www.who.int/quantifying_ehimpacts/publications/en/leadebd2.pdf)).
2. IPCS (1995). *Inorganic lead*. Geneva, World Health Organization, International Programme on Chemical Safety (Environmental Health Criteria 165; <http://www.inchem.org/documents/ehc/ehc/ehc165.htm>).
3. WHO (2009). *Global health risks: Mortality and burden of disease attributable to selected major risks*. Geneva, World Health Organization ([http://www.who.int/healthinfo/global\\_burden\\_disease/GlobalHealthRisks\\_report\\_full.pdf](http://www.who.int/healthinfo/global_burden_disease/GlobalHealthRisks_report_full.pdf)).
4. IARC (2006). *Summaries & evaluations: Inorganic and organic lead compounds*. Lyon, International Agency for Research on Cancer (IARC Monographs for the Evaluation of Carcinogenic Risks to Humans, Vol. 87; <http://www.inchem.org/documents/iarc/vol87/volume87.pdf>).
5. UNEP (2008). *Draft final review of scientific information on lead*. Geneva, United Nations Environment Programme, Chemicals Branch ([http://www.chem.unep.ch/Pb\\_and\\_Cd/SR/Draft\\_final\\_reviews/Pb\\_Review/Final\\_UNEP\\_Lead\\_review\\_Nov\\_2008.pdf](http://www.chem.unep.ch/Pb_and_Cd/SR/Draft_final_reviews/Pb_Review/Final_UNEP_Lead_review_Nov_2008.pdf)).
6. WHO (2007). *Health risks of heavy metals from long-range transboundary air pollution*. Copenhagen, World Health Organization Regional Office for Europe (<http://www.euro.who.int/document/E91044.pdf>).
7. WHO (2003). *Lead in drinking-water. Background document for development of WHO Guidelines for Drinking-Water Quality*. Geneva, World Health Organization (WHO/SDE/WSH/03.04/09; [http://www.who.int/water\\_sanitation\\_health/dwq/chemicals/lead.pdf](http://www.who.int/water_sanitation_health/dwq/chemicals/lead.pdf)).
8. WHO (2007). *Lead exposure in children*. Geneva, World Health Organization ([http://www.who.int/phe/news/Lead\\_in\\_Toys\\_note\\_060807.pdf](http://www.who.int/phe/news/Lead_in_Toys_note_060807.pdf)).
9. WHO (in preparation). *Safety evaluation of certain food additives and contaminants in food*. Geneva, World Health Organization (WHO Food Additives Series, No. 64; <http://www.who.int/ipcs/publications/jecfa/monographs/en/index.html>) [summary in FAO/WHO (2010). *Summary and conclusions of the seventy-third meeting of the Joint FAO/WHO Expert Committee on Food Additives, Geneva, 8–17 June 2010*. Rome, Food and Agriculture Organization of the United Nations; Geneva, World Health Organization (JECFA/73/SC; <http://www.who.int/entity/foodsafety/publications/chem/summary73.pdf>)].
10. WHO (2008). *Guidelines for drinking-water quality, 3rd edition incorporating 1st and 2nd addenda. Vol. 1. Recommendations*. Geneva, World Health Organization, pp. 392–394 ([http://www.who.int/water\\_sanitation\\_health/dwq/GDW12rev1and2.pdf](http://www.who.int/water_sanitation_health/dwq/GDW12rev1and2.pdf)).
11. WHO (2001). Lead. In: *Air quality guidelines for Europe, 2nd ed.* Copenhagen, World Health Organization Regional Office for Europe ([http://www.euro.who.int/\\_data/assets/pdf\\_file/0005/74732/E71922.pdf](http://www.euro.who.int/_data/assets/pdf_file/0005/74732/E71922.pdf)).
12. WHO (2010). *Quantifying environmental health impacts*. Geneva, World Health Organization ([http://www.who.int/quantifying\\_ehimpacts/en/](http://www.who.int/quantifying_ehimpacts/en/)).

13. WHO (2009). *Quantification of the disease burden attributable to environmental risk factors*. Geneva, World Health Organization, Department of Public Health and Environment ([http://www.who.int/quantifying\\_ehimpacts/summaryEBD\\_updated.pdf](http://www.who.int/quantifying_ehimpacts/summaryEBD_updated.pdf)).
14. WHO (2004). Lead exposure. In: *Comparative quantification of health risks*. Geneva, World Health Organization, pp. 1495–1542 (<http://www.who.int/publications/cra/chapters/volume2/1495-1542.pdf>).
15. WHO (2007). *Blood lead levels in children*. Copenhagen, World Health Organization Regional Office for Europe, European Environment and Health Information System (Fact Sheet No. 4.5; [http://www.enhis.org/object\\_document/o4738n27387.html](http://www.enhis.org/object_document/o4738n27387.html)).

© World Health Organization 2010

All rights reserved.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damage arising from its use.

WHO gratefully acknowledges the financial support provided by the Swiss Federal Office for the Environment.

Printed by the WHO Document Production Services, Geneva, Switzerland

**Public Health and Environment  
World Health Organization  
20 Avenue Appia, 1211 Geneva 27, Switzerland**