





Cameroon

African Region



I. Epidemiological profile

Population (UN)	2015	%	Parasites and vectors
High transmission (> 1 case per 1000 population)	16,600,000	71	Plasmodium species: <i>P. falciparum</i> (100%), <i>P. vivax</i> (0%)
Low transmission (0-1 cases per 1000 population)	6,770,000	29	Major anopheles species: <i>An. gambiae</i> , <i>An. arabiensis</i> , <i>An. funestus</i> , <i>An. moucheti</i> , <i>An. nili</i>
Malaria-free (0 cases)	0	-	Reported confirmed cases (health facility): 1,162,784 Estimated cases: 5,300,000 [3,500,000 ; 7,700,000]
Total	23,340,000	-	Confirmed cases at community level: 30,497 Reported deaths: 3,440 Estimated deaths: 9,200 [4,900 ; 13,000]

II. Intervention policies and strategies

Intervention	Policies/strategies	Yes/No	Adopted	Antimalarial treatment policy	Medicine	Year adopted
ITN	ITNs/ LLINs distributed free of charge	Yes	2004	First-line treatment of unconfirmed malaria	AS+AQ	2004
	ITNs/ LLINs distributed to all age groups	Yes	2011	First-line treatment of <i>P. falciparum</i>	AS+AQ	2004
IRS	IRS is recommended	Yes	2007	Treatment failure of <i>P. falciparum</i>	QN	2004
	DDT is authorized for IRS	No	-	Treatment of severe malaria	AS, AM;QN	2004
Larval control	Use of larval control recommended	No	-	Treatment of <i>P. vivax</i>	-	-
IPT	IPT used to prevent malaria during pregnancy	Yes	2004	Dosage of Primaquine for radical treatment of <i>P. vivax</i>	-	-
Diagnosis	Patients of all ages should receive diagnostic test	Yes	2011	Type pf RDT used	-	-
	Malaria diagnosis is free of charge in the public sector	No	-			
Treatment	ACT is free of charge for all ages in public sector	No	-			
	The sale of oral artemisinin-based monotherapies (oAMTs)	Is banned	-			
	Single dose of primaquine is used as gametocidal medicine for <i>P. falciparum</i>	No	-			
	Primaquine is used for radical treatment of <i>P. vivax</i>	No	-			
	G6PD test is a requirement before treatment with primaquine	No	-			
	Directly observed treatment with primaquine is undertaken	No	-			
	System for monitoring adverse reactions to antimalarials exists	No	2004			
Surveillance	ACD for case investigation (reactive)	-	-			
	ACD of febrile cases at community level (pro-active)	No	-			
	Mass screening is undertaken	No	-			
	Uncomplicated <i>P. falciparum</i> cases routinely admitted	No	-			
	Uncomplicated <i>P. vivax</i> cases routinely admitted	No	-			
	Foci and case investigation undertaken	-	-			
	Case reporting from private sector is mandatory	-	-			

Therapeutic efficacy tests (clinical and parasitological failure, %)


Medicine	Year	Min	Median	Max	Follow-up	No of studies	Species
AL	2006-2013	0	1.9	5	28 days	12	<i>P. falciparum</i>
AS+AQ	2005-2013	0	3.7	8.7	28 days	15	<i>P. falciparum</i>

Insecticide susceptibility bioassays (reported resistance to at least one insecticide for any vector at any locality)

Year	Pyrethroid	DDT	Carbamate	Organophosphate	Species/complex tested
2010-2014	Yes	Yes	Yes	No	<i>An. gambiae</i> s.s.

III. Financing


Sources of financing




Government expenditure by intervention in 2015

IV. Coverage

Coverage of ITN and IRS




Cases tested and treated in public sector




V. Impact


Cases treated



Test positivity



Confirmed malaria cases per 1000 and ABER



Malaria admissions and deaths (per 100 000)

