





United Republic of Tanzania (Mainland)

African Region



Based on 2015 reported data


I. Epidemiological profile

Population (UN)	2016	%	Parasites and vectors
High transmission (> 1 case per 1000 population)	53,600,000	100	Plasmodium species: <i>P. falciparum</i> (100%)0%
Low transmission (0-1 cases per 1000 population)	-	-	Major anopheles species: <i>An. gambiae</i> , <i>An. arabiensis</i> , <i>An. funestus</i>
Malaria-free (0 cases)	-	-	Reported confirmed cases (health facility): 5,188,505 Estimated cases: -
Total	53,590,000	-	Confirmed cases at community level: - Reported deaths: 5,045 Estimated deaths: -

II. Intervention policies and strategies


Intervention	Policies/strategies	Yes/No	Adopted	Antimalarial treatment policy	Medicine	Year adopted					
ITN	ITNs/ LLINs distributed free of charge	No	2008	First-line treatment of unconfirmed malaria	AL	2004					
	ITNs/ LLINs distributed to all age groups	No	2009	First-line treatment of <i>P. falciparum</i>	AL	2004					
IRS	IRS is recommended	Yes	2006	Treatment failure of <i>P. falciparum</i>	QN	2004					
	DDT is authorized for IRS	No	-	Treatment of severe malaria	AS, AM; QN	2004					
Larval control	Use of larval control recommended	Yes	-	Treatment of <i>P. vivax</i>	-	-					
IPT	IPT used to prevent malaria during pregnancy	Yes	2001	Dosage of Primaquine for radical treatment of <i>P. vivax</i>	-	-					
Diagnosis	Patients of all ages should receive diagnostic test	Yes	2009	Type of RDT used	-	-					
	Malaria diagnosis is free of charge in the public sector	No	1997								
Treatment	ACT is free of charge for all ages in public sector	Yes	2006	Therapeutic efficacy tests (clinical and parasitological failure, %)							
	The sale of oral artemisinin-based monotherapies (oAMTs)	Is banned	2006	Medicine	Year(s)	Min	Median	Max	Follow-up	No. of studies	Species
	Single dose of primaquine is used as gametocidal medicine for <i>P. falciparum</i>	No	-	Organophosphates	2011-2016	0.8	1	1	39	<i>An. arabiensis</i> , <i>An. gambiae</i> s.l.	
	Primaquine is used for radical treatment of <i>P. vivax</i>	No	-	Carbamates	2010-2016	0.2	0.9	1	50	<i>An. gambiae</i> s.l.	
	G6PD test is a requirement before treatment with primaquine	No	-	Organochlorines	2010-2015	0.1	0.9	1	50	<i>An. arabiensis</i> , <i>An. gambiae</i> s.l.	
	Directly observed treatment with primaquine is undertaken	No	-	Pyrethroids	2010-2016	0	0.8	1	68	<i>An. arabiensis</i> , <i>An. gambiae</i> s.l.	
System for monitoring adverse reactions to antimalarials exists	Yes	2001									
Surveillance	ACD for case investigation (reactive)	No	-	Insecticide resistance tests (mosquito mortality, %)							
	ACD of febrile cases at community level (pro-active)	No	-	Insecticide class	Years	Min	Mean	Max	No. of sites	Species	
	Mass screening is undertaken	No	-	Organophosphates	2011-2016	0.8	1	1	39	<i>An. arabiensis</i> , <i>An. gambiae</i> s.l.	
	Uncomplicated <i>P. falciparum</i> cases routinely admitted	No	-	Carbamates	2010-2016	0.2	0.9	1	50	<i>An. gambiae</i> s.l.	
	Uncomplicated <i>P. vivax</i> cases routinely admitted	No	-	Organochlorines	2010-2015	0.1	0.9	1	50	<i>An. arabiensis</i> , <i>An. gambiae</i> s.l.	
	Foci and case investigation undertaken	-	-	Pyrethroids	2010-2016	0	0.8	1	68	<i>An. arabiensis</i> , <i>An. gambiae</i> s.l.	
Case reporting from private sector is mandatory	Yes	-									

III. Financing




Government expenditure by intervention in 2016


IV. Coverage




Cases tested and treated in public sector




V. Impact



Test positivity



Confirmed malaria cases per 1000 and ABER



Malaria admissions and deaths (per 100 000)

