


Bulgaria


Total population: 7 278 000
Income Group: Upper middle

Percentage of population living in urban areas: 73.1%
Population proportion between ages 30 and 70 years: 55.8%

Age-standardized death rates*





Proportional mortality (% of total deaths, all ages, both sexes)*



Total deaths: 106,000
NCDs are estimated to account for 94% of total deaths.

Premature mortality due to NCDs*

The probability of dying between ages 30 and 70 years from the 4 main NCDs is 24%.



Adult risk factors

	males	females	total
Current tobacco smoking (2011)	48%	31%	39%
Total alcohol per capita consumption, in litres of pure alcohol (2010)	17.9	5.3	11.4
Raised blood pressure (2008)	44.3%	39.5%	41.8%
Obesity (2008)	23.1%	24.3%	23.7%

National systems response to NCDs

Has an operational NCD unit/branch or department within the Ministry of Health, or equivalent	DK
Has an operational multisectoral national policy, strategy or action plan that integrates several NCDs and shared risk factors	No
Has an operational policy, strategy or action plan to reduce the harmful use of alcohol	No
Has an operational policy, strategy or action plan to reduce physical inactivity and/or promote physical activity	No
Has an operational policy, strategy or action plan to reduce the burden of tobacco use	No
Has an operational policy, strategy or action plan to reduce unhealthy diet and/or promote healthy diets	No
Has evidence-based national guidelines/protocols/standards for the management of major NCDs through a primary care approach	Yes
Has an NCD surveillance and monitoring system in place to enable reporting against the nine global NCD targets	No
Has a national, population-based cancer registry	Yes

* The mortality estimates for this country have a high degree of uncertainty because they are not based on any national NCD mortality data (see Explanatory Notes).
DK = Country responded "don't know"