

Equatorial Guinea

Population	504 000
GNI/capita	21 230 US\$
% urbanization	39%
% people living in cities greater than 100 000 inhabitants	25%
Population below the poverty line (national)	NA
Population below the poverty line (international, <\$1/day)	NA
Under age 5 mortality rate	206/1000 live births (2006)
Life expectancy	46 years (2006)

Environmental burden of disease for selected risk factors, per year

Estimates based on national exposure and WHO country health statistics 2004















Risk factor	Exposure	Deaths /year	DALYs/ 1000 cap /year
Water, sanitation and hygiene	Improved water:	43%	700
	Improved sanitation:	53%	44
Indoor air	SFU% households:	NA	NA
Outdoor air	Mean urban PM10: 12 ug/m3	-	-
Main malaria vectors	<i>A. gambiae</i> , <i>A. funestus</i> , <i>A. melas</i> , <i>A. moucheiti</i> , <i>A. carnevalei</i>		
Main other vectors	<i>Culex quinquefasciatus</i> , anopheline spp		

Environmental burden of disease (preliminary), per year

Estimates based on Comparative Risk Assessment, evidence synthesis and expert evaluation for regional exposure and WHO country health statistics 2004

DALYs/1000 cap	(World - lowest: 13, highest: 289)	169
Deaths		2 500
% of total burden		30%

Environmental burden by disease category [DALYs/1000 capita], per year

Disease group	World's lowest country rate	Country rate	World's highest country rate
Diarrhoea	0.2		47
Respiratory infections	0.1		28
Malaria	0.0		29
Other vector-borne diseases	0.0		2.0
Lung cancer	0.0		0.1
Other cancers	0.3		1.8
Neuropsychiatric disorders	1.4		1.8
Cardiovascular disease	1.4		3.9
COPD	0.0		1.4
Asthma	0.3		2.3
Musculoskeletal diseases	0.5		0.7
Road traffic injuries	0.3		5.2
Other unintentional injuries	0.6		9.7
Intentional injuries	0.0		1.9

Other indicators

Use of leaded gasoline	No	(2008)
Overcrowding	14%	(1993)
Malnutrition (% stunting)	35%	(2004)