

Ethiopia

Population	77.4 mio
GNI/capita	780 US\$
% urbanization	16%
% people living in cities greater than 100 000 inhabitants	7%
Population below the poverty line (national)	44% (1999-2000)
Population below the poverty line (international, <\$1/day)	23% (1999-2000)
Under age 5 mortality rate	123/1000 live births (2006)
Life expectancy	56 years (2006)

Environmental burden of disease for selected risk factors, per year

Estimates based on national exposure and WHO country health statistics 2004

Risk factor	Exposure	Deaths /year	DALYs/ 1000 cap /year
Water, sanitation and hygiene	Improved water: 22%	112 100	47
	Improved sanitation: 13%		
Indoor air	SFU% households: >95%	72 400	31
Outdoor air	Mean urban PM10: 88 ug/m3	2 500	0.5
Main malaria vectors	<i>A. arabiensis, A. nlli</i>		
Main other vectors	<i>Phlebotomus longipes, P. pedifer</i>		

Environmental burden of disease (preliminary), per year

Estimates based on Comparative Risk Assessment, evidence synthesis and expert evaluation for regional exposure and WHO country health statistics 2004

DALYs/1000 cap	(World - lowest: 13, highest: 289)	146
Deaths		336 800
% of total burden		31%

Environmental burden by disease category [DALYs/1000 capita], per year

Disease group	World's lowest country rate	Country rate	World's highest country rate
Diarrhoea	0.2	49	107
Respiratory infections	0.1	28	71
Malaria	0.0	9.8	34
Other vector-borne diseases	0.0	1.3	4.9
Lung cancer	0.0	0.1	2.6
Other cancers	0.3	1.1	4.1
Neuropsychiatric disorders	1.4	1.7	3.0
Cardiovascular disease	1.4	2.6	14
COPD	0.0	0.8	4.6
Asthma	0.3	1.2	2.8
Musculoskeletal diseases	0.5	0.6	1.5
Road traffic injuries	0.3	3.2	15
Other unintentional injuries	0.6	6.1	30
Intentional injuries	0.0	1.3	7.5

Other indicators

Use of leaded gasoline	No	(2008)
Overcrowding	NA	
Malnutrition (% stunting)	51%	(2005)