



Levelling up (part 1): **a discussion paper on concepts and principles** **for tackling social inequities in health**

Margaret Whitehead
Göran Dahlgren

WHO Collaborating Centre for
Policy Research on Social Determinants of Health
University of Liverpool



Levelling up (part 1):
a discussion paper on concepts
and principles for tackling social
inequities in health

Margaret Whitehead
Göran Dahlgren

The WHO Regional Office for Europe is one of six regional offices throughout the world, each with its own programme geared to the particular health problems of the countries it serves. The European Region embraces some 870 million people living in an area stretching from Greenland in the north and the Mediterranean in the south to the Pacific shores of the Russian Federation. The European programme of WHO therefore concentrates both on the problems associated with industrial and post-industrial society and on those faced by the emerging democracies of central and eastern Europe and the former USSR.

To ensure the widest possible availability of authoritative information and guidance on health matters, WHO secures broad international distribution of its publications and encourages their translation and adaptation. By helping to promote and protect health and prevent and control disease, WHO's books contribute to achieving the Organization's principal objective – the attainment by all people of the highest possible level of health.

WHO European Office for Investment for Health and Development

The WHO European Office for Investment for Health and Development, which coordinated the activities leading to this publication, was set up by the WHO Regional Office for Europe, with cooperation and support from the Ministry of Health and the Veneto Region of Italy. One of its key responsibilities is to provide evidence on and act upon the social and economic determinants of health. The Office systematically reviews what is involved in drawing together the concepts, scientific evidence, technology and policy action necessary to achieve effective investment for the promotion of health and synergy between social, economic and health development. The Office fulfils two interrelated main functions:

- to monitor, review and systematize the policy implications of the social and economic determinants of population health;
and
- to provide services to help Member States in the WHO European Region increase their capacity to invest in health by addressing these policy implications and integrating them into the agenda for development.



EUROPE

Levelling up (part 1):
a discussion paper on concepts
and principles for tackling social
inequities in health

Margaret Whitehead
Göran Dahlgren

Address requests about publications of the WHO Regional Office for Europe to:

Publications
WHO Regional Office for Europe
Scherfigsvej 8
DK-2100 Copenhagen Ø, Denmark

Alternatively, complete an online request form for documentation, health information, or for permission to quote or translate, on the WHO/Europe web site at <http://www.euro.who.int/pubrequest>.

© World Health Organization 2006

All rights reserved. The Regional Office for Europe of the World Health Organization welcomes requests for permission to reproduce or translate its publications, in part or in full.

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Where the designation “country or area” appears in the headings of tables, it covers countries, territories, cities, or areas. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers’ products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

The World Health Organization does not warrant that the information contained in this publication is complete and correct and shall not be liable for any damages incurred as a result of its use. The views expressed by authors or editors do not necessarily represent the decisions or the stated policy of the World Health Organization.

Keywords

HEALTH SERVICES ACCESSIBILITY SOCIOECONOMIC FACTORS - DELIVERY OF HEALTH CARE - ECONOMICS, MEDICAL - HEALTH POLICY - EUROPE

CONTENTS

	Acknowledgements	VII
	<i>Foreword</i>	VIII
Part I.	Concepts	1
	What is the difference between variations in health and social inequities in health?	2
	Fairness and human rights	3
	Inequality and inequity are synonymous	4
	So what is equity in health?	4
	Widespread inequities throughout Europe	5
	A widening health divide	5
	The phenomenon of the social gradient	6
	Social inequities in access to health services	7
	What does equitable health care look like?	10
	Different goals for equity in health and in health care	11
Part II.	Ten principles for policy action	13
	1. Policies should strive to level up, not level down	14
	2. The three main approaches to reducing social inequities in health are interdependent and should build on one another	14
	3. Population health policies should have the dual purpose of promoting health gains in the population as a whole and reducing health inequities	16
	4. Actions should be concerned with tackling the social determinants of health inequities	16
	5. Stated policy intentions are not enough: the possibility of actions doing harm must be monitored	17
	6. Select appropriate tools to measure the extent of inequities and the progress towards goals	17
	7. Make concerted efforts to give a voice to the voiceless	18
	8. Wherever possible, social inequities in health should be described and analysed separately for men and women	18
	9. Relate differences in health by ethnic background or geography to socioeconomic background	19
	10. Health systems should be built on equity principles	20
	Summary	20
	References	22
Annex.	Illustration of the difference between relative and absolute measures of social inequities in health.	29



Acknowledgements

This report was prepared in close cooperation with senior staff of the WHO European Office for Investment for Health and Development in Venice and with advice from their Expert Group on Poverty, Health Inequalities and Related Social and Economic Determinants of Health. Drafts of this paper, and its companion paper on strategies, have been presented at various meetings of WHO staff and at European consultations held in the Venice Office for advice. These drafts have also been subjected to anonymous peer reviews. Revisions have been made according to the comments gratefully received, but the authors alone are responsible for the final text and any errors it contains.

Text editing: Jerome M. Rosen

Graphic design: Marta Pasqualato

Administrative support: Antonella Biasiotto

Foreword

Why this topic and why now?

In 1990, the WHO Regional Office for Europe published two technical documents: The concepts and principles of equity and health (Whitehead, 1990), followed by a companion paper on policies and strategies (Dahlgren & Whitehead, 1992). Both documents tried to distil the elusive concept of equity, in the context of health, and to clarify what the implications would be for policy development in this field. These documents were intended for a non technical audience of policy-makers and practitioners. Their aim was to raise awareness and stimulate debate in a wider general audience, and they certainly did that: the documents were taken up across Europe and North America, translated into 20 languages, and incorporated into training materials for a wide spectrum of disciplines.

Why is a similar exercise necessary now? Surely, fundamental concepts and principles do not change that much. In response to the question, several key points need to be made. First, the European context has changed, with social divisions across and within European countries widening in many respects. These divisions have challenged policy-makers to the limit. In terms of actions needed to tackle health inequities, new concerns, new ways of addressing issues and collective actions may now be possible, and these require reformulation of principles to match present-day needs. Of particular relevance is the new body of evidence that indicates the social cause of many health inequities. Second, new debates about how to measure differences in health have appeared in the literature. However, without a parallel conceptual debate about the assumptions built into these measures, the technical developments risk providing little support to policy-makers. Also, they may not measure what they intended to or may distort objectives to match what can be measured. Third, concepts, principles and measurements of social inequity in health need urgently to be related to equity-oriented strategies and actions, both within and without the health policy domain.

The WHO Regional Office for Europe has evolved and re-emphasized its key values (WHO Regional Office for Europe, 2005). There has, for example, been a renaissance of the Declaration of Alma-Ata (WHO, 1978). This landmark declaration by the health ministers of the world, in 1978, identified the importance of equity in health, the need to recognize the social determinants


of health and, hence, to take intersectoral action. The health ministers also recognized the importance of participation by communities in decisions about their health. These principles reflect strongly the needs of the current situation, not least in Europe (WHO Regional Office for Europe, 2005). The recently deceased Director-General of WHO, Dr Lee Jong-wook, made equity and social justice the underlying theme of his office, emphasizing the need to strengthen health systems to combat poverty and ill health. Before his death, he initiated the formation of the global Commission on Social Determinants of Health (Lee, 2004).

Over the past 20 years, the European Region of WHO has been at the forefront of advocating policies that promote equity. In 2002, it reaffirmed this commitment by setting up the WHO European Office for Investment for Health and Development (the WHO Venice Office), which focuses specifically on the social determinants of health and what health systems can do to confront poverty – with intensified work directed at the eastern parts of the Region.

*As part of an effort to develop useful tools and guidance for countries on the issue of equity in health, the WHO Venice Office invited Margaret Whitehead and Göran Dahlgren to prepare this paper on the concepts and principles needed to tackle social inequities in health. The present paper is the result of a wide range of consultations. Earlier versions of this paper have been discussed in several meetings organized by WHO, including one with the European Committee for Health Promotion and Development, in 2005. This final version is also the result of inputs from the WHO Regional Office for Europe and WHO headquarters technical units and experts in the field in various countries. Together with the forthcoming *Levelling up (part 1): a discussion paper on European strategies for tackling social inequities in health (Dahlgren and Whitehead, 2006)*, this discussion paper marks the first official contribution of the new WHO Collaborating Centre for Policy Research on Social Determinants of Health to the development, analysis and dissemination of knowledge on the social pathways to and from social inequities. It is hoped that this work will help policy-makers in their efforts to address socially caused inequities in health in a Europe that is rapidly changing.*

Erio Ziglio
Head,

WHO European Office for Investment for Health and Development



Part I.

Concepts

Part I. Concepts

What is the difference between variations in health and social inequities in health?

Social inequities in health are the central focus of this paper. Such inequities concern systematic differences in health status between different socioeconomic groups. But what exactly does that entail? Within any country, differences in health can be observed across the population. Genetic and constitutional variations ensure that the health of individuals varies, as it does for any other physical characteristic. The prevalence of ill health also differs between different age groups, with older people tending to be sicker than younger people, because of the natural ageing process. Biologically, women in older industrialized countries exhibit an advantage in survival over men at every stage of life. Chance also plays a role in everyone's life, with luck deciding which individuals avoid a particular infectious disease or hazard and which succumb.

Three distinguishing features, when combined, turn mere variations or differences in health into a social inequity in health. They are systematic, socially produced (and therefore modifiable) and unfair.

The first feature is the systematic pattern of the differences in health. These differences are not distributed randomly, but show a consistent pattern across the population. One of the most striking examples is the systematic differences in health between different socioeconomic groups. Mortality and morbidity increase with declining social position, as illustrated in subsequent sections. This social pattern of disease is universal, though its magnitude and extent vary among countries.

The second feature is the social processes that produce health differences, rather than these differences being determined biologically. No Law of Nature, for instance, decrees that the children of poor families should die at twice the rate as that of children born into rich families (Blane et al., 1993), so this health inequity is not fixed or inevitable. Theoretically, at least, if social processes generate these differences in a country, then these differences should be amenable to alteration by a concerted effort by that country.

The third feature is that social inequities are differences widely considered to be unfair, because they are generated and maintained by what Evans & Peters

(2001) have termed “unjust social arrangements” that offend common notions of fairness. Of course, this depends on the meaning attributed by different people to the idea of unfair. Although ideas about what is unfair may differ to a certain degree from place to place, there is much common ground. For example, most (if not all) people in European countries share the view that all children, regardless of social group, should have the same chance of survival. It would be widely considered unfair if the chance of survival was much poorer for the children of some socioeconomic groups, compared with that of others. This is but one illustration of an all-embracing concern across Europe for linking fairness to human rights.

Fairness and human rights

The bias and discrimination that lead to differences in access to the resources and opportunities for health between social groups is unfair. This touches on the special place that health holds in human rights: everyone has the right to enjoy the highest attainable standard of health in their society (WHO, 1946). Health is also a unique resource for achieving other objectives in life, such as better education and employment. Health is therefore a way of promoting the freedom of individuals and societies (Sen, 2000).

It is therefore important for a society to organize its health resources equitably, so that access to those resources are open to everybody. The existence of clear social differentials in health and in their determinants (illustrated in subsequent sections) goes against accepted values of fairness and justice (Daniels, Kennedy & Kawachi, 2000).

In today’s Europe, working out what social differences in health are fair and unfair is unnecessary. Essentially, all systematic differences in health between different socioeconomic groups within a country can be considered unfair and, therefore, classed as health inequities. There is no biological reason for their existence, and it is clear that even systematic differences in lifestyles between socioeconomic groups are to a large extent shaped by structural factors. Summing up briefly, social inequities in health are directly or indirectly generated by social, economic and environmental factors and structurally influenced lifestyles. These determinants of social inequities are all amenable to change.

In the International Covenant on Economic, Social and Cultural Rights, the

wording deliberately sets health in the context in which people live (Kälin et al., 2004). For the purpose of taking action, the health status of groups of people who are better off can be used as a practical indicator of the standard of health attainable in any given society and as the standard to which policies that address inequities in health should strive.

Inequality and inequity are synonymous

Earlier papers by the present authors, adopted the phrase inequities in health throughout, while explaining that in some countries, notably the United Kingdom, the phrase inequalities in health was used and had the same meaning. In the intervening years, more European countries have adopted the British terminology, as illustrated by the title of the 2005 EU Summit on Tackling Inequalities in Health. For consistency with other WHO documents, however, the phrase social inequities in health has been retained in this paper. The authors would still like to emphasize, though, that in the public health community the phrase social inequalities in health carries the same connotation of health differences that are unfair and unjust. Indeed, as some European languages have only one word for the two terms, there is no distinction between the two when they are translated.

So what is equity in health?

If inequity in health is unfair and unjust, what then is the converse: equity in health? The concept is related intimately to the central human rights thread that has run right through the key articles of WHO, from its inception in the 1940s to the resolutions of the 21st century. The WHO Constitution (WHO, 1946) asserted back in 1946 that “the highest standards of health should be within reach of all, without distinction of race, religion, political belief, economic or social condition”.

Echoing these sentiments nearly 60 years later, equity in health implies that ideally everyone could attain their full health potential and that no one should be disadvantaged from achieving this potential because of their social position or other socially determined circumstance.

This refers to everyone and not just to a particularly disadvantaged segment of the population. Efforts to promote social equity in health are therefore aimed at

creating opportunities and removing barriers to achieving the health potential of all people. It involves the fair distribution of resources needed for health, fair access to the opportunities available, and fairness in the support offered to people when ill.

The outcome of these efforts would be a gradual reduction of all systematic differences in health between different socioeconomic groups. The ultimate vision is the elimination of such inequities, by levelling up to the health of the most advantaged.

Widespread inequities throughout Europe

A widening health divide

Within the countries of Europe, there are many examples of systematic differences in health between different social groups. In all European countries, most disadvantaged groups have worse health and higher mortality. This manifests itself in large differences in life expectancy between the extremes of the social scale. In Scotland, for instance, a baby born in the most disadvantaged neighbourhood in Glasgow can expect to live 10 fewer years than a baby living in the most affluent neighbourhood (Acheson et al., 1998). In Estonia, the gap in mortality between the groups with the highest and lowest levels of education increased tremendously from 1989–2000, the transition period after the cessation of Soviet rule. By 2000, a male graduate 25 years of age could expect to live 13 years longer than men the same age with the lowest level of education (Leinsalu, Vagero & Kunst, 2003).

Examples from the northern and southern parts of the Region are also evident. In the Netherlands, for example, there is a 5-year gap in life expectancy, and a 13-year gap in disability-free life expectancy, between men from groups with low and high levels of education (van de Water, Boshuizen & Perenboom, 1996). Also, women in Finland from the group with the lowest level of education have eight fewer years of disability-free life expectancy than women from the group with the highest level of education (Valkonen et al., 1994). Moreover, in the south, studies in Turin, Italy, have shown that although mortality decreased for all socioeconomic groups over the past 30 years, the decrease was less pronounced among men and women with a lower level of education and among men with poorer housing conditions (Marinacci et al., 2004).

In France, two examples are striking. Between a 35-year-old unskilled manual worker and a white-collar (*cadre supérieur*) worker of the same age, the difference in life expectancy is about nine years. When looking at rates of premature death (before 54 years of age), four times more unskilled manual workers die prematurely of diabetes than do white collar workers, and the difference is tenfold when it comes to cirrhosis and alcohol psychosis (Mesrine, 1999; Jouglu et al., 2000).

The health situation in central and eastern European countries is of grave concern. The Russian Federation had a drastic, indeed unprecedented, deterioration in life expectancy after the economic and political shocks of the early 1990s. This deterioration did not affect all segments of the population equally. The hardest hit were the least educated groups. In fact, between 1990 and 1994, the chances of survival for men from groups with lower levels of education were poorer than in the 1970s and 1980s. Mortality rates among people with a university education, however, were closer to mortality rates for general western populations (Shkolnikov, Field & Andreev, 2001).

The phenomenon of the social gradient

Typically, a stepwise, or linear, decrease in health – and not just an extreme group in poor health and the rest in reasonably good health – is seen with decreasing social position and is referred to as the social gradient (Marmot, 2004). For example, the health of the populations of Florence, Leghorn and Turin has been followed over a number of years, and the findings clearly show that mortality increases linearly with increasing degree of social disadvantage. This is so both when social disadvantage is measured by the characteristics of individuals, such as education, employment or social class, and when it is measured by deprivation of the area in which people live (Costa, Spadea & Dirindin, 2002).

The impact of the social gradient on health is sometimes expressed as a shortfall in health – that is, the number of lives that would have been saved if all groups in society had the same high level of health as the most advantaged group. As an example of a shortfall, it has been estimated that if all working men 20–64 years of age in England and Wales had the same mortality rates as men of the same age in professional and managerial positions, then over 17 000 fewer deaths would have occurred each year in the early 1990s (Acheson et al., 1998). Excess mortality in the more deprived areas of Spain, compared

with the most affluent, has been estimated at 35 000 deaths a year (Benach & Yasui, 1999). A similar shortfall calculation for the Netherlands estimated that average morbidity and mortality in the Dutch population would be reduced by 25–50% if men with lower levels of education had the morbidity and mortality levels of men with a university education (Mackenbach, 1994).

The existence of these social inequities in almost all countries poses the most serious challenge to improving the health of the Region's population, in general. Advances in life expectancy for a country as a whole, for example, are held back by the magnitude of the inequity experienced within that country.

Inequity does, however, indicate the scale of the improvement possible. It demonstrates, for example, what groups with the greatest advantages have already achieved and, therefore, what is feasible for others to attain in a particular country at a particular time. Social inequities in health among babies and children provide the starkest examples of injustice. Evidence of large differences in infant mortality between rich and poor babies in Stockholm as far back as the 1930s was a source of political concern, which led to a programme of maternal health care reforms. When combined with more general social security and housing reforms, to improve the living conditions of families in the city, these programmes and reforms brought about a reduction in the social inequity in infant mortality (Burström, 2004). Today, in Sweden, the inequity in infant mortality is almost eliminated, which is not always the case in other high-income countries.

Social inequities in access to health services

Inadequate access to essential health services is one of several determinants of social inequities in health. It may not be the major determinant, but it is an important one for the health sector to tackle directly – to put its own house in order. Furthermore, the burden of payment for health services is a growing cause of poverty, particularly for socially vulnerable groups, and is one for which the health sector has special responsibility (and opportunity) to address effectively (Ziglio et al., 2003).

Taking the human rights approach, a very basic right for people is that of having access to effective health care that ameliorates their suffering when they become sick, that protects them and prevents them from developing disease

in the first place, and that helps them maintain their own health when well. Without the benefits that access to health services can bring, by improving health and providing freedom from pain and suffering, all other human activities are compromised. But for many millions of people around the globe, even across a relatively rich region like Europe, access is inadequate for the needs of some groups or may even be unattainable for all practical purposes. Moreover, those in the greatest need often have the poorest access to care – a striking example of unfairness.

The idea of inequity in access to care was famously captured in the Inverse Care Law (Hart, 1971), which states, “The availability of good medical care tends to vary inversely with the need for it in the population served.” This so-called law came to Tudor Hart when he surveyed the situation in the coal-mining valleys of Wales. The location of health facilities and their quality were much worse in the poorer towns where the miners lived and suffered from occupational disabilities, compared with the facilities available in the more prosperous towns where morbidity rates were lower.

The Welsh example above relates to geographic access – that is, the location and, therefore, physical availability of health services in different parts of a country. Studies from around Europe have found large differences in geographic access for different population groups. In northern Europe, for instance, recent studies in Sweden have shown a clustering of specialists serving the more affluent neighbourhoods in Stockholm and a scarcity in the low-income areas of the same city (Dahlgren, 1994).

Two other types of access, however, are also important: economic access and cultural access. What use is there in having health services on your doorstep if you cannot afford to use them? This is the situation in which many urban dwellers in some countries find themselves. The starkest example of the problem of economic access is when people in need of emergency care are turned away from a clinic or hospital and left to die, because they cannot afford to pay. This is very rare in Europe, but there are an increasing number of instances of patients delaying seeking non-urgent care for financial reasons. In Belgium, for example, recent surveys found that patients with chronic illnesses spent an average of 23% of their disposable income on care. Also, about a third of the Belgian population reported that they experienced difficulty in paying for medical care, and 8% of families postponed seeking medical care because of the cost (Louckx, 2002).

In France, likewise, a study in 1997 found that 600 000 people did not have access to social security to cover medical care costs and that 16% of the population did not have supplementary coverage. Many of them delayed seeking treatment because of the cost. In addition, the emergency safety net system operated in such a way that, although the very poor could request free medical aid from the local government department each time they needed treatment, the procedure for doing so was stigmatizing. This French system was reformed in 2000, to ensure that previously excluded segments of the population could have improved economic access to care in a less stigmatizing way (Ziglio et al., 2003).

Another aspect and illustration of the problem of economic access is when people do muster the money to pay out-of-pocket fees (including informal payments), but are then pushed into debt as a result. This we have termed the medical poverty trap – that is, impoverishment caused by paying for medical care (Whitehead, Dahlgren & Evans, 2001). The most striking examples come from the developing world, but the medical poverty trap is prevalent in the developed world, most notably in the United States and more recently in eastern Europe. The burden of payment for essential health services and drugs should therefore always be analysed, as it might be substantial, even if it does not drive people into poverty (see companion paper: Dahlgren & Whitehead, 2006).

The other type of access, cultural access, relates to acceptability and respect. Do some groups experience cultural barriers to available services, rendering them unacceptable? Are there unacceptable shortcomings in the respect and dignity afforded to marginalized groups, such as the homeless or impoverished patients, by health workers? Language barriers and cultural practices, for example, may prevent minority ethnic groups or recent immigrants from accepting preventive care and benefiting from psychiatric care, even when free. Aside from language barriers, another major barrier is that between professional health workers and less educated patients, where health service providers lack an awareness and understanding of the day-to-day restrictions in the lives of patients living in hardship. Also, there may be differences in the attention that patients from less advantaged backgrounds receive, which leads to differences in the quality of care and the respect they are afforded.

Also, differences in treatment outcome between different socioeconomic groups may be observed – for example, when recovery after an operation is slower for poorer patients, or when drug treatment to control a chronic disease is less successful for people living in inadequate housing. Such differences may occur even when the treatment has been provided in an equitable way, because of the socioeconomic conditions under which patients live. These differences in treatment outcome are the result of inequities in the wider social determinants of health, over which the health services may have little or no control. For this reason, striving for equity in treatment outcome is not a feasible option for a health service.

What does equitable health care look like?

In consideration of these potential inequities in access, the converse – equity in health care – can be seen as being multifaceted and incorporating ideas about fair arrangements that allow equal geographic, economic and cultural access to available services for all in equal need of care.


The idea of allowing for differences in need for health services across different socioeconomic groups is of critical importance in definitions of equity in health care. This is most easily illustrated by the use of health services by different socioeconomic groups, where service use is taken as an indirect indicator of the wider issues of access. If utilization rates were found to be similar for each socioeconomic group, it would not necessarily signify equitable service. On the contrary, it would most probably indicate an inequitable situation. This is because, given the social gradient in health status, the underlying need for care tends to be greater among lower socioeconomic groups and, therefore, correspondingly greater use of services by them than by more advantaged groups would be expected if their increased need was being met.

A pragmatic goal for equity in health care in the case just mentioned would include striving for equal use for equal need. The need for care, however, has to be taken into account when assessing progress towards this goal. A segment of the population with full eligibility to use a service may exercise their right not to use it, resulting in lower utilization rates for this group than the level of need indicates (Whitehead, 1990). Preventive screening and health checks provided nationally are cases in point, where some people may be reluctant to take up the service offered. In this situation, pressuring people to use a service against

their will would be inequitable. Where further investigation reveals, however, that the reason for not using the services is poor geographic access or social or economic barriers, so that people who wished to use the service could not, this would be considered unacceptable and unfair.

Different goals for equity in health and in health care

In practical terms, it is important to understand that the goals for equity in health and equity in health care are very different. With equity in health, the ultimate goal would be the elimination of all systematic differences in health status between socioeconomic groups, as stated in the section on “So what is equity in health?” The end goal of equity in health care, however, would be to closely match services to the level of need, which may very well result in large differences in access and use of services between different socioeconomic groups, favouring the more disadvantaged groups in greatest need.



Part II.

Ten principles for policy action

Part II. Ten principles for policy action

The concepts of health equity imply certain pragmatic principles of action when striving to reduce social inequities in health status. These are set out below as 10 principles for general guidance. A framework for analysing causes of social inequities in health, and for highlighting policy options and strategies for reducing them, is contained in the companion paper (Dahlgren & Whitehead, 2006).

1. Polices should strive to level up, not level down

Nobody would seriously suggest trying to close the health gap by bringing healthier people down to the level of the least healthy. A worsening in the infant mortality rate of the babies of rich parents, for example, with no change in the mortality rate of poor babies, would not be seen as a success, but would rather be seen as a tragedy – even if it led to a narrowing of the differences between the two groups, purely in terms of measurement. Yet, opponents of an equity policy have warned of this danger. Therefore, to make it absolutely clear, the principle set out in this paper emphasizes that the only way to narrow the health gap in an equitable way is to bring up the level of health of the groups of people who are worse off to that of the groups who are better off. Levelling-down is not an option.

2. The three main approaches to reducing social inequities in health are interdependent and should build on one another

The relative merits of different ways of addressing social inequities in health have been debated recently (Mackenbach et al., 2002; Graham, 2004a). Essentially, the three main approaches being applied to measure and tackle social inequities in health are: focusing on people in poverty only, narrowing the health divide and reducing social inequities throughout the whole population.

- a** Focusing on people in poverty only. This is a so-called targeting approach, which measures progress in terms of an improvement in health for the targeted group only, without any reference to improvements in health taking place in the population as a whole or among the most privileged group. From this perspective, any improvement in the health status of disadvantaged groups can

be considered a success, even if the health divide between rich and poor is increasing.

Narrowing the health divide. This approach takes as its starting point the health of disadvantaged groups relative to the rest of the population. The focus of action in this category is to reduce the gap between the worst off in society and the best off – the disparity in health status between the extremes of the social scale. **b**

Reducing social inequities throughout the whole population. This approach recognizes that morbidity and premature mortality tend to increase with declining socioeconomic status and that they are not just an issue of a gap in health between rich and poor. This approach therefore takes in the whole population, including middle-income groups, and seeks to reduce the differences in health between high-, middle- and low-income groups, by equalizing health opportunities across the socioeconomic spectrum. **c**

The definition of equity in health adopted in this paper encourages seeing these three approaches as not only complementary to one another, but also seeing them as interdependent. They must build on one another. The logical sequence is therefore to improve health among disadvantaged groups as a first step. This targeted approach can be assessed as successful when the health of the poor is improved even marginally, even if inequities in health overall are increasing. The second step, narrowing the health divide, has the more ambitious aim of improving the health of people in poverty at a faster rate than that among the rich. The third step is to reduce health inequities between all groups, not just between the extremes of the social scale. The third approach, however, cannot be isolated from the other two, as reduced differences can technically be achieved by reducing the health divide between middle and higher socioeconomic groups, while neglecting people in poverty and leaving their health even further behind. The only valid indicator of reduced social inequities throughout the whole population is when the health of the most disadvantaged groups has improved faster than that of the middle- and high-income groups, as explained in Principle 1 above.

3. Population health policies should have the dual purpose of promoting health gains in the population as a whole and reducing health inequities

Some portray these twin goals as conflicting, presenting a trade-off between improved health for the population as a whole and even faster improvement in health among the worse off in society – that is, between overall gains in population health and reducing social inequities in health. This is a false trade-off. The objective of reducing health inequities constitutes an integral part of a comprehensive strategy for health development, not an alternative option. In reality, no national strategy in Europe, or elsewhere, abandons attempts to improve the health of the population as a whole in favour of concentrating solely on reducing health inequities. Also, it is increasingly recognized that national health targets for the population as a whole stand little chance of being met without attention to the health of the worst off in society. The two goals typically go hand in hand.

4. Actions should be concerned with tackling the social determinants of health inequities

This principle focuses not only on the social determinants of health in general (the social conditions that can affect people's health), but also focuses on the main determinants of the systematic differences in opportunities, living standards and lifestyles associated with different positions in society (Graham, 2004b). Working conditions are a good illustration of this point. In post-industrial Europe, exposure to poor working conditions has ceased to be a major determinant of ill health in the population overall. However, a study in Sweden found that differences in exposure to poor working conditions across the social spectrum explained a considerable proportion of the observed inequities in health between socioeconomic groups in the country (Lundberg, 1991). Tackling such social determinants, therefore requires a greater understanding of the processes that generate and maintain social inequities and then intervening in these processes at the most effective points (see companion paper: Dahlgren & Whitehead, 2006).

Stated policy intentions are not enough: the possibility of actions doing harm must be monitored

5,

This principle requires an assessment of differential impacts, not just average effects. The classic example in the field of health equity is that of the adverse effects of some health sector reforms of recent decades, which have created a medical poverty trap (Whitehead, Dahlgren & Evans, 2001), where the increasing necessity to pay for care when sick pushes more people into poverty. In this case, the welfare system, which originally intended to support the sick, is turned into a poverty-generating system. This trap developed at a pace that required vigilance to catch it in time. Similarly, interventions designed to help people in poverty may be implemented in such a way as to stigmatize the very people the programme was designed to help and, in so doing, push them to avoid the help on offer.

This principle of carrying out health inequity impact assessments applies to a variety of policies outside and inside the health sector. Indeed, the greatest danger may lie in wider macro-policies that hide the negative health impacts, because they are not seen as health related (see companion paper: Dahlgren & Whitehead, 2006).

Select appropriate tools to measure the extent of inequities and the progress towards goals

6.

This principle may seem obvious, but as principles 1–5 illustrate, interventions intended to reduce inequities can be focused on one of several distinct goals or targets, and each one may require a separate indicator. Measures that only monitor changes in health of the poor, for example, will not be able to contribute anything about how poorer groups are faring relative to more advantaged groups. This requires measuring progress at both the top and bottom of the social scale and then comparing these two measurements. This comparison between the extremes of the social scale will not be able to assess the impact across the whole of society. This may require indicators of the so-called shortfall – that is, the cumulative difference between the most advantaged group and each successive social group for each specific factor.

It is important to monitor both relative and absolute changes in social inequities in health, because they give different information about the magnitude and direction of change. An example of a relative measure is the ratio of the mortality rate of the most disadvantaged group to the mortality rate of the most privileged group. An absolute measure in this case would be the difference between mortality rates of the disadvantaged and privileged groups. Table 1 in the Appendix gives a numerical example, using English data on trends in circulatory diseases by deprivation category. Using the absolute measure in this example, inequities in mortality show a decrease, while using the relative measure they show an increase.

7. Make concerted efforts to give a voice to the voiceless

This principle entails, for example, seeking the views of marginalized groups and increasing their genuine participation (as opposed to token consultation). The more articulate members of the population and those with the most powerful representation tend to have more influence than those in a weaker position. To address this, administrators and professionals need to make a determined effort to provide administrative systems and information to make it easier for lay people to participate in decisions that affect their health.

8. Wherever possible, social inequities in health should be described and analysed separately for men and women

This separate description and analysis is needed because both the magnitude and the causes of observed social inequities in health are sometimes different for the two sexes. It is therefore of critical importance that these differences are known and taken into consideration when developing strategies to combat inequities in health.

The value of combining gender-specific and socioeconomic analyses has been clearly illustrated recently by the Swedish accident prevention programme. Data on accidents for the population as a whole indicated relatively good progress on injury prevention. When the data were separated by sex and social status, however, very different rates and types of accidents were observed for girls compared with boys, men compared with women, and low-income groups

compared with high-income groups. The combined gender and socioeconomic analysis also revealed that working class women, in particular, had very high levels of risk in the home and in the workplace. Once these differences were recognized, the need to tailor prevention strategies by gender and socioeconomic condition became clear (La Flamme, 1998). The same is increasingly true for tobacco control policies (Kunst, Giskes & Mackenbach, 2004).

Another reason for ensuring that systematic differences in health by gender are analysed by socioeconomic background is that the causes may differ by social position. Poor women, for example, may be discriminated against both for being women and for being poor. Tackling the differences in health generated by this so-called double burden may need different strategies from those designed to tackle the gender effects on health experienced by more affluent women. The present trend to neglect gender in analyses of social inequities in health and, conversely, to neglect social position in gender-specific analyses should therefore be replaced by a combined approach that considers both social position and gender.

Relate differences in health by ethnic background or geography to socioeconomic background

9.

Analyses of systematic differences in health by ethnic background should, whenever possible, be related to socioeconomic background, as the magnitude and causes of the ethnic differences observed tend to differ by social position. Likewise, ethnic background needs to be included in analyses of social inequities in health in countries with marked ethnic discrimination.

Differences in health between geographical areas should also be analysed, with due consideration to differences in the social structure. Age-adjusted health status in areas with a fairly homogenous population from a socioeconomic perspective can then be used as a proxy for assessing socioeconomic inequities in health, when measures of individual socioeconomic status, such as a person's occupation or income, are not available routinely.

10. Health systems should be built on equity principles

Equity principles include the following.

- a** Public health services should not be driven by profit, and patients should never be exploited for profit.
- b** Services should be provided according to need, not ability to pay. This requires a system of health care financing that pools risks across the population, so that those at high risk are subsidized by those at low risk at any given time.
- c** The same high standard of care should be offered to everyone, without discrimination with respect to social, ethnic, gender or age profile.
- d** The underlying values and equity objectives of a health system should be explicitly identified, and the monitoring carried out to ensure these objectives are approached in the most efficient way possible.


Summary

Social inequities in health are systematic differences in health status between different socioeconomic groups. These inequities are socially produced (and therefore modifiable) and unfair. In practice, all systematic differences in health between socioeconomic groups in European countries could be regarded as unfair and avoidable, and therefore regarded as inequities. This judgement about unfairness is based on universal human rights principles.

The evidence points to the existence of extensive (and widening) social inequities in health in Europe today. The need to take action to reduce these inequities and their root causes is becoming ever more pressing as a major public health challenge. This calls for a new way of thinking about the direction of policy and also calls for renewed vigilance in monitoring impacts, to make sure that no segment of the population is excluded or loses out.

Increasing numbers of countries across Europe have been striving to face the challenge of social inequities in health and are working out what practical action can be taken in their own country to improve the situation. The aim of

this paper is to help promote a common understanding of the concepts and principles on which actions for tackling health inequities can be based.



References

References

Acheson D et al. (1998). *Independent inquiry into inequalities in health report*. London, The Stationery Office.

Benach J, Yasui Y (1999). Geographical patterns of excess mortality in Spain explained by two indices of deprivation. *Journal of Epidemiology and Community Health*, 53:423–431.

Blane D, Davey Smith G, Bartley M (1993). Social selection: what does it contribute to social class differences in health? *Sociology of Health and Illness*, 15:1–15.

British Department of Health (2005). *Tackling health inequalities: status report on the programme for action*. London, Department of Health (<http://www.dh.gov.uk/assetRoot/04/11/76/98/04117698.pdf>, accessed 15 May 2006).

Burström B (2004). Social differentials in the decline of infant mortality in Sweden in the twentieth century: the impact of policies and policy. In: Navarro V, ed. *The political and social contexts of health*. New York, Baywood Publishing Company Inc.

Costa G, Spadea T, Dirindin N (2002). Italy. In: Mackenbach J, Bakker M, eds. *Reducing inequalities in health: a European perspective*. London: Routledge: 229–239.

Dahlgren G (1994). *Framtidens sjukvårdsmarknader – vinnare och förlorare?* [Health care markets of the future – winners and losers?]. Stockholm, Bokförlaget Natur och Kultur (in Swedish).

Dahlgren G, Whitehead, M (1992) *Policies and strategies to promote equity in health*. Copenhagen, WHO Regional Office for Europe (document number: EUR/ICP/RPD 414 (2); [http://whqlibdoc.who.int/euro/-1993/EUR_ICP_RPD414\(2\).pdf](http://whqlibdoc.who.int/euro/-1993/EUR_ICP_RPD414(2).pdf), accessed 15 May 2006).

Dahlgren G, Whitehead M (2006). *Levelling up (part 2): a discussion paper on European strategies for tackling social inequities in health*. Copenhagen, WHO Regional Office for Europe (Studies on social and economic determinants of population health, No.3).

Daniels N, Kennedy B, Kawachi I (2000). Justice is good for our health. In: Cohen J, Rogers J, eds. *Is inequality bad for our health?* Boston, Beacon Press.

Evans T, Peters F (2001). Ethical dimensions of health equity. In: Evans T et al., eds. *Challenging inequities in health: from ethics to action*. New York, Oxford University Press.

Graham H (2004a). Tackling inequalities in health in England: remedying health disadvantages, narrowing health gaps or reducing health gradients? *Journal of Social Policy*, 33(1):115–131.

Graham H (2004b). Social determinants and their unequal distribution: clarifying policy understandings. *The Milbank Quarterly*, 82:101–124.

Hart JT (1971). The inverse care law. *Lancet*, 1(7696): 405–412.

Jougla E et al. (2000). Chapitre de :Les inégalités sociales de santé.[Social inequalities in health] In: de Leclerc A et al., eds. *La découverte*. Paris, Inserm (http://www.inserm.fr/fr/questionsdesante/dossiers/sante_environnement/att00000579/12septembre2000.pdf, accessed 15 May 2006) (in French).

Källin W, Müller L, Wyttenbach J (2004). *The face of human rights*. Baden, Switzerland, Lars Müller Publishers.

Kunst A, Giskes K, Mackenbach J on behalf of the EU Network on Interventions to Reduce Socio-economic Inequalities in Health (2004). *Socio-economic inequalities in smoking in the European Union. Applying an equity lens to tobacco control policies*. Rotterdam, Department of Public Health, Erasmus Medical Centre.

Laflamme L (1998). *Social inequality in injury risks: knowledge accumulated and plans for the future*. Stockholm, Karolinska Institute and National Institute of Public Health (Report 1998: 33).

Lee Jong-wook. (2004). *Address by the Director-General to the Fifty-seventh World Health Assembly, Geneva, Tuesday, 18 May 2004*. Geneva, World Health Organization (document number: A57/3).

Leinsalu M, Vagero D, Kunst A (2003). Estonia 1989–2000: enormous increase in mortality differences by education. *International Journal of Epidemiology*, 32(6):1081–1087.

Louckx F (2002). Patient cost sharing and access to care. In: Mackenbach J, Bakker M, eds. *Reducing inequalities in health: a European perspective*. London, Routledge: 188–198.

Lundberg O (1991). Causal explanations for class inequality in health – an empirical analysis. *Social Science & Medicine*, 32(4):385–393.

Mackenbach J (1994). Socioeconomic inequalities in health in the Netherlands: impact of a five year research programme. *BMJ*, 309(6967):1487–1491.

Mackenbach JP et al. (2002). Strategies to reduce socioeconomic inequalities in health. In: Mackenbach J, Bakker M, eds. *Reducing inequalities in health: a European perspective*. London: Routledge: 25–50.

Marinacci C et al. (2004). The role of individual and contextual socioeconomic circumstances on mortality: analysis of time variations in a city of north west Italy. *Journal of Epidemiology and Community Health*, 58(3):199–207.

Marmot M (2004). *The status syndrome: how social standing affects our health and longevity*. London, Bloomsbury Publishing Plc.

Mesrine A. (1999). Les différences de mortalité par milieu social restent fortes [Differences in mortality according to social class are still important]. *Données Sociales*, 228–235.

Sen A (2000). *Development as freedom*. New York, Random House, Inc.

Shkolnikov VM, Field MG, Andreev EM (2001). Russia: dimensions of the gender gap in mortality. In: Evans T et al., eds. *Challenging inequities in health: from ethics to action*. New York, Oxford University Press: 138–155.

Valkonen T, Sihvonen A-P, Lahelma E (1994). Disability-free life expectancy by education in Finland. In: Mathers CD, McCallum J, Robine J-M, eds. *Advances in health expectancies. Proceedings of the 7th meeting of the International Network*

on Health Expectancy (REVES), Canberra, February 1994. Canberra, Australian Institute of Health and Welfare: 160–168.

Van de Water HPA, Boshuizen HC, Perenboom RJM (1996). Health expectancy in the Netherlands 1983–1990. *European Journal of Public Health*, 6:21–28.

Whitehead M (1990). *The concepts and principles of equity and health*. Copenhagen, WHO Regional Office for Europe (document number: EUR/ICP/RPD 414; http://whqlibdoc.who.int/euro/-1993/EUR_ICP_RPD_414.pdf, accessed 16 May 2006).

Whitehead M, Dahlgren G, Evans T (2001). Equity and health sector reforms: can low-income countries escape the medical poverty trap? *Lancet*, 358(9284):833–836.


Whitehead M, Dahlgren G, Gilson L (2001). Developing the policy response to inequities in health: a global perspective. In: Evans T et al., eds. *Challenging inequities in health: from ethics to action*. New York, Oxford University Press: 308–324.

WHO (1946). *Constitution of the World Health Organization*. London, World Health Organization (http://whqlibdoc.who.int/hist/official_records/constitution.pdf, accessed 16 May 2006).

WHO (1978). *Primary health care: report of the International Conference on Primary Health Care, Alma-Ata, USSR, 6–12 September 1978*. Geneva, World Health Organization (Health for all series, No. 1; <http://whqlibdoc.who.int/publications/9241800011.pdf>, accessed 16 May 2006).

WHO Regional Office for Europe (2005). *The Health for All policy framework for the WHO European Region: 2005 update*. Copenhagen, WHO Regional Office for Europe (European health for all series, No. 7; <http://www.euro.who.int/document/e87861.pdf>, accessed 16 May 2006).

Ziglio E et al., eds (2003). *Health systems confront poverty*. Copenhagen, WHO Regional Office for Europe (Public Health Case Studies Series, No. 1; <http://www.euro.who.int/document/e80225.pdf>, accessed 16 May 2006).



Annex.

Illustration of the difference between relative and absolute measures of social inequities in health

Annex. Illustration of the difference between relative and absolute measures of social inequities in health

Absolute and relative changes in age-standardized death rates (per 100 000 population) for circulatory diseases in people less than 75 years of age, by deprivation area, in England, 1995–1997 and 2001–2003

Table I

Rates and gaps	1995–1997	2001–2003	Change in inequity
Death rate for the 20% most deprived local authorities (in deaths per 100 000 population)	173	129	--
Death rate for England as a whole (in deaths per 100 000 population)	141	103	--
Absolute gap (difference) between disadvantaged and England as a whole (in deaths per 100 000 population)	$173-141 = 32$	$129-103 = 26$	Reduction
Relative gap (ratio) between disadvantaged and England as a whole	$173/141 = 1.22$	$129/103 = 1.25$	Increase

Source: Adapted from British Department of Health (2005): 32-33



EUROPE

World Health Organization
Regional Office for Europe
Scherfigsvej 8, DK-2100 Copenhagen Ø, Denmark

Tel.: + 45 39 17 17 17.

Fax: + 45 39 17 18 18.

E-mail: postmaster@euro.who.int

Web site: www.euro.who.int

