

# **Rolling Revision of the WHO Guidelines for Drinking-Water Quality**

**Draft for review and comments  
(Not for citation)**

## **Chloral hydrate in drinking-water Summary statement**



**World Health Organization  
July 2004**

© World Health Organization 2004

Requests for permission to reproduce or translate WHO publications - whether for sale or for non-commercial distribution - should be addressed to Publications (Fax: +41 22 791 4806; e-mail: [permissions@who.int](mailto:permissions@who.int)).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned.

Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

The World Health Organization does not warrant that the information contained in this publication is complete and correct and shall not be liable for any damage incurred as a result of its use.

## **Chloral hydrate (trichloroacetaldehyde)**

Chloral hydrate can be formed as a by-product of the chlorination of water containing organic precursor material, such as fulvic and humic acids. It has been found in drinking-water at concentrations of up to 100 µg/litre, but concentrations are usually below 10 µg/litre. Concentrations are generally higher in surface water than in groundwater, and concentrations appear to increase across the distribution system.

Chloral hydrate is used as an intermediate in the production of insecticides, herbicides and hypnotic drugs. It has also been widely used as a sedative or hypnotic drug in humans at oral doses of up to about 750–1000 mg/day. Clinical exposure is considerably greater than exposure from drinking-water, but of shorter-term duration.

No epidemiological or carcinogenic studies were found in humans associating chloral hydrate with cancer, despite the fact that chloral hydrate has been used for many decades (and still is used) as a sedative and hypnotic drug in adults and children (specifically for dental procedures). IARC classified chloral hydrate as not classifiable as to its carcinogenicity to humans (Group 3), based on inadequate evidence in humans and limited evidence in experimental animals. There is equivocal evidence of genotoxicity for chloral hydrate.

A health-based value of 0.1 mg/litre (rounded figure) can be calculated on the basis of a conservative TDI of 0.0045 mg/kg of body weight per day derived based on an increased incidence of liver histopathology observed in B6C3F<sub>1</sub> mice in a 2-year drinking-water study, allocating 80% of the TDI to drinking-water (because most exposure to chloral hydrate is from drinking-water) and assuming a 60-kg adult consuming 2 litres of water per day. However, because chloral hydrate usually occurs in drinking-water at concentrations well below those at which toxic effects are observed, it is not considered necessary to derive a health-based guideline value.

Chloral hydrate levels in drinking-water can be controlled by enhanced coagulation and softening, which will remove DBP precursors; moving the point of disinfection to reduce the reaction between chlorine and DBP precursors; and using chloramines for residual disinfection instead of chlorine.

### ***History of guideline development***

The 1958, 1963 and 1971 WHO *International Standards for Drinking-water* and the first edition of the *Guidelines for Drinking-water Quality*, published in 1984, did not refer to chloral hydrate. The 1993 Guidelines established a provisional health-based guideline value of 0.01 mg/litre for chloral hydrate in drinking-water. The guideline value was designated as provisional because of the limitations of the available database, necessitating the use of an uncertainty factor of 10 000. This guideline value was brought forward to the third edition of the Guidelines.

***Assessment date***

The risk assessment was conducted in 2004.

***Principal references***

IPCS (2000) *Chloral hydrate*. Geneva, World Health Organization, International Programme on Chemical Safety (Concise International Chemical Assessment Document 25).

IPCS (2000) *Disinfectants and disinfectant by-products*. Geneva, World Health Organization, International Programme on Chemical Safety (Environmental Health Criteria 216).

WHO (2004) *Chloral hydrate (trichloroacetaldehyde) in drinking-water. Background document for preparation of WHO Guidelines for drinking-water quality*. Geneva, World Health Organization (WHO/SDE/WSH/04.0x/49).