

13.

NUTRIENT MINERALS IN DRINKING WATER: IMPLICATIONS FOR THE NUTRITION OF INFANTS AND YOUNG CHILDREN

Erika Sievers

Institute of Public Health
North Rhine Westphalia
Munster, Germany

I. INTRODUCTION

The WHO Global Strategy on Infant and Young Child Feeding emphasizes the importance of infant feeding and promotes exclusive breastfeeding in the first six months of life. In infants who cannot be breast-fed or should not receive breast milk, substitutes are required. These should be a formula that complies with the appropriate Codex Alimentarius Standards or, alternatively, a home-prepared formula with micronutrient supplements (1). Drinking water is indispensable for the reconstitution of powdered infant formulae and needed for the preparation of other breast-milk substitutes. As a result of the long-term intake of a considerable volume in relation to body weight, the concentrations of nutrient minerals in drinking water may contribute significantly to the total trace element and mineral intake of infants and young children. This is especially applicable to formula-fed infants during the first months of life, who may be the most vulnerable group affected by excessive concentrations of nutrients or contaminants in drinking water.

Defining essential requirements of the composition of infant formulae, the importance of the quality of the water used for their reconstitution has been acknowledged by the Scientific Committee on Food, SCF, of the European Commission (2). Although it was noted that the mineral content of water may vary widely depending upon its source, the optimal composition remained undefined. Recommendations for the composition of infant formulae refer to total nutrient content as prepared ready for consumption according to manufacturer's instructions. However, these usually inform the consumer about the quantity of powder and drinking water required as well as other technical advice for the reconstitution of the formula, but do not refer to the mineral concentration in drinking water itself.

The use of desalination to provide drinking water may continue to increase all over the world. Remineralization of mineral content in waters that have been treated by demineralization may provide an opportunity to improve compositional choices. Specific aspects of vulnerable groups have to be considered, but the needs of infants and young children, especially if nourished with infant formula instead of human milk, may be of special concern. The question arises whether special nutrient mineral properties of these waters should be considered for use for infants and young children.

II. ASSESSMENT OF MINERAL INTAKE IN INFANT NUTRITION

Public health decisions about the composition of infant foods and the use of drinking water in infancy require awareness of the approaches used in the study methodology applied to nutrition trials in infancy and early childhood. These approaches may be of crucial importance in the interpretation of the results and potential limitations of their significance for the issues raised.

Analytical results of infant formulae may only refer to analyses of the powder itself or formulae prepared under standardized laboratory conditions with defined water (3). Metabolic studies investigating mineral bioavailability have attempted to exclude or minimize the potential influences of different water supplies, e.g. (4). They therefore have been performed either in the clinic, on metabolic wards, or at home using a standardized water supply or ready-to-feed liquid infant formula. On the other hand, recently published epidemiological trials concerning mineral supplementation in infancy neither assess representative mineral concentrations in the drinking water at a household level or the quantitative water intake (5-7). For example, investigations concerning the effects of copper concentrations in drinking water, analyzed neither the respective nutrition nor representative specimens (8). Finally, studies addressing both drinking water and powdered infant formula referred to the concentrations given by the manufacturer, but analysed the reconstituted formula, too (9).

III. THE QUANTITATIVE INTAKE OF DRINKING WATER IN INFANCY AND EARLY CHILDHOOD

Health and well-being in early childhood are dependent on an optimal supply of essential nutrients. Nutrient requirements for growth necessitate intake and positive retention of essential trace elements and minerals. The immaturity of homeostatic mechanisms in gastrointestinal and renal functions of resorption and excretion also raises a concern over the susceptibility to an excess or a deficiency of nutrients as well as to toxic substances during this crucial period for mental and motor development. A major factor in the nutrient mineral intake from drinking water is the quantity of water consumed.

1. The Choice of Nutrition – a significant Factor of Drinking Water Intake in Infancy

In considering the consumption of drinking water by vulnerable populations, a figure of 0.75 litres per day has been used for a 5kg child and a figure of one litre per day for a 10kg child (10). Although these figures may be applicable for standard calculations, the range of quantitative water intake observed in populations at that age might be considerable according to the Food and Nutrition Board of the Institute of Medicine (11). The choices made in terms of how an infant is fed, especially in the first six months of life, are significant factors in determining the level of nutrient mineral intake from drinking water:

- *Healthy, exclusively breast-fed infants* nourished according to present recommendations (1) will not directly consume drinking water in the first six months of life. They might eventually be indirectly affected by a potential passage of (trace) minerals into human milk. Unfortunately, sometimes the early introduction of supplementary feedings of water, tea or other nutrients is practised. After six months, the introduction of complementary foods and breastfeeding for up to two years of age or beyond is recommended. During this period, household drinking water intake will depend on the extent of breastfeeding and on whether complementary foods are prepared at home or bought commercially.
- *Healthy infants fed powdered infant formula* consume drinking water from the day they are born owing to the fact that it is used to reconstitute the product. Both may contribute to the mineral intake up until complementary foods are introduced. Different sources of water may be used, including tap water, well water, and natural bottled mineral water labelled as suitable for infant nutrition. Specific compositional characteristics of water used in a household or the use of stagnant water are very likely to have an effect throughout the entire formula feeding period if parents are not aware of the potential consequences. Infants are likely to be fed formula prepared at home, while older children or adults may consume considerable amounts

of drinking water from sources outside the household. Powdered formulae do not necessarily originate from the region in which they are consumed and may be subject to specific directives established by the place of origin. Therefore, different ecological and legal factors may independently affect the two components making up the nutrient mineral composition of infant formula.

- *Home-prepared formulas* are more likely to be based on local produce and are affected by local factors such as soil composition, the use of fertilizers or pollution. These may influence, too, the composition of the drinking water in the area. Home-prepared formulae may be prepared with a 1:1 dilution of cow's milk to drinking water with the addition of other components (12). Other concepts are based on a 2:1 dilution of cow's milk (13), with a resulting lower quantitative intake of drinking water. After the age of six months, drinking water intake will also be dependent on the choice of complementary foods and beverages and how they are prepared.
- *The choice of complementary food introduced after six months* may vary widely and it influences the evaluation of the nutrient mineral content in drinking water considerably. Inappropriate increases of the mineral intake may be of concern. Use of foods that are naturally rich in trace elements like zinc and iron (e.g. meat) or minerals like calcium (e.g. dairy products) is recommended, but may not be practised because these foods are not available, affordable or acceptable for the family. This may lead to reliance on predominantly plant-based or vegetarian diets of poor nutritional quality (12, 14). In these situations, and depending on its composition, drinking water may contribute a considerable part of the dietary mineral intake.

Assumptions about the mineral intake in infants may be influenced by factors not directly linked to the actual product composition. The "standard intake" of drinking water calculated by WHO in early infancy is 750 ml (10); the same figure is given by the SCF (2) for infant formula. Most standard instructions for infant formulae require 90g water to 100g formula. Therefore, the basic intake assumptions for the quantitative intake of drinking water in formula-fed infants could differ by approximately 10%. Recent recommendations supported a reduction of the upper margins for caloric density in infant formulae and follow-on formulae. As nutrient requirements for powdered infant formulae are given per 100 kcal intake (e.g. (2)), this results in a reduction of the recommended upper levels of mineral concentrations (-7% or -12%). These examples may only refer to differences of approximately 10%, but they emphasize the need for transparency regarding the factors that influence the final composition of infant formulae.

2. Specific aspects of quantitative intake of drinking water in infancy and childhood

In exclusively breast-fed infants, the mean intake increases from 699g up to 854g at the age of six months (15). The quantitative milk consumption of formula-fed infants exceeds the intake of breast-fed infants at the age of 2-4 months (16). The potential implications for the quantitative intake of household drinking water (Figure 1) have been calculated by re-evaluating data assessed in a longitudinal trial on nutrient mineral supply in breast-fed and formula-fed infants (17, 18):

Exclusive breast-feeding had been recommended for a minimum of four months in infants receiving human milk, the other families were provided with liquid infant-milk formulae. For infants from the age of five months, powdered follow-on formulae, age-specific complementary feeds, powdered cereals and a standard mineral water suitable for infant feeding were supplied. Weighing each meal over a period of 72 hours enabled its nutrient intake to be assessed. Liquid infant formulae were recalculated based on the assumption that the powdered formula and drinking water used corresponded to an equivalent product marketed in Germany. This precept


enabled the differences in drinking-water intake between full-term, breast-fed infants and premature or full-term formula-fed infants to be assessed (Figure 1, Table 1). Up until the age of four months, only two breast-fed infants consumed small amounts of drinking water. At the same time, the daily drinking water intake needed for the re-constitution of powdered infant formula decreased from 158 to 140g/kg in premature infants and from 153 to 122.9g/kg in full-term infants. With all families in the study being provided with standardized complementary feeds, it was not until infants reached the age of 32 weeks that comparable intakes of drinking water in all groups (57% of total water intake, Table 1) were attained.

Table 1. Drinking water intake in infancy (median, range)

Age \pm 2 weeks	4	8	16	24	32	42	52
Infants studied	Total water intake, g x kg⁻¹ x d¹						
FF-Pt	158.3 (128.1-201.9)	148.0 (123.2-254.2)	140.1 (102.2-172.4)	90.2 (52.9-130.6)	78.5 (58.0-116.2)	77.9 (65.1-110.9)	81.3 (56.9-116.6)
FF-T	152.7 (111.8-224.1)	142.6 (99.6-240.0)	122.9 (106.5-157.8)	94.5 (81.9-117.8)	87.9 (69.3-139.6)	83.7 (53.8-135.3)	77.7 (63.4-149.9)
BF-T	132.7 (81.7-147.7)	111.4 (75.6-153.0)	97.8 (50.1-144.2)	98.8 (66.9-132.2)	84.9 (71.4-140.3)	82.9 (60.0-165.3)	93.7 (49.2-192.2)
	Drinking water intake, g x kg⁻¹ x d¹						
FF-PT	158.3 (122.8-201.9)	148.0 (112.3-254.2)	140.1 (102.2-172.4)	70.3 (48.6-130.6)	43.1 (31.7-99.9)	42.9 (25.9-76.6)	43.9 (21.3-67.9)
FF-T	152.7 (111.8-224.1)	142.6 (99.6-240.0)	122.9 (106.5-157.8)	76.5 (63.4-100.6)	47.8 (38.0-97.8)	48.7 (33.3-89.9)	50.0 (33.2-104.6)
BF-T	0.0 (0.0-4.7)	0.0 (0.0-14.0)	0.0	18.6 (0.0-83.9)	46.6 (3.8-96.5)	49.5 (9.8-118.1)	48.9 (0.8-148.8)
	Drinking water intake, % of total water intake						
FF-PT	100.0 (95.8-100.0)	100.0 (91.1-100.0)	100.0 (98.9-100.0)	88.2 (65.4-100.0)	57.7 (41.4-85.9)	53.3 (39.6-73.5)	55.2 (37.4-71.9)
FF-T	100.0	100.0 (99.9-100.2)	100.0	81.7 (67.8-100.0)	57.7 (48.1-70.1)	58.2 (54.8-75.1)	58.9 (44.8-70.4)
BF-T	0.0 (0.0. to 3.6)	0.0 (0.0-11.4)	-	16.8 (0.0-84.3)	57.0 (4.6-71.6)	63.8 (12.2-77.3)	66.3 (0.9-81.3)

FF-PT: Premature, formula-fed infants, investigated in parallel with term infant groups, gestational age at birth 29 (25 - 32) weeks; fed cow's-milk-protein-based infant formula at least up until the age of 16 weeks, corrected for gestational age; n= 16 (14-15) infants. FF-T: Term, formula-fed infants, fed cow's-milk-protein-based infant formula at least up until the age of 16 weeks; n=15 (11-14) Infants. BF-T: Term, exclusively or predominantly breast-fed infants up until the age of 16 weeks; n = 20 (14-16) infants.

Figure 1. Household drinking water intake of breast-fed term infants and formula-fed term and preterm infants (based on mineral nutrition studies: 17, 18).



These drinking water intakes exceed those assessed in the DONALD (Dortmund Nutritional and Anthropometrical Longitudinally Designed) Study (19). The authors reported that in formula-fed infants, the daily intake decreased longitudinally from 107 ± 28 g/kg at three months to 33 ± 25 g/kg at twelve months of age. Comparing specifically the tap-water intake of formula- and breast-fed infants, the mean total daily tap-water intake in breast-fed infants observed throughout the first year was 15 ± 23 g/kg and, of formula-fed infants, 49 ± 33 g/kg. Between the ages of one and three years, tap-water intake averaged 15 ± 20 g/kg. The differences may be explained by several factors: the DONALD study omitted early infancy up until 3 months of age, where higher intakes are observed. Lower intakes of tap drinking water may be indicative of a higher proportion of home-prepared meals. The DONALD study also reported on water intake in school children (20) and assessed the hydration status of children aged 4 to 10.9 years through the analysis of water intake, urine volume and urine osmolality. Based on their estimates of adequate daily total water intake, the authors confirmed the total water intake of 1.0ml/kcal in children that was recommended earlier (21). The mean intake volumes reported for the age groups 4.0-6.9 and 7.0-10.9 were 1495g and 1834g for boys, and 1318g and 1545g for girls. The trends over the 15-year period of the study revealed an increase in total water intake in 2-13-year-old children irrespective of sex, attributable to increased beverage consumption on the one hand, and decreased milk, coffee/tea and soft-drink consumption on the other (22). The results may be influenced by regional aspects and the high interest in nutrition documented by the longitudinal participation in the trial. Other authors have been rather concerned about consumption of large quantities of squash on the one hand and the low consumption of plain water on the other (23).

IV. THE CONTRIBUTION OF DRINKING WATER TO NUTRIENT MINERAL INTAKE IN INFANCY AND EARLY CHILDHOOD

The potential significance of mineral nutrients in infant nutrition, but especially in human milk, has attracted a great deal of attention over the last few decades (2, 14, 15). The upperbound acceptable mineral concentrations in WHO Drinking Water Guidelines, mature human milk, and the recent recommendations on the composition of infant formulae and follow-on formulae are compared in Table 2. Age-specific nutrient recommendations from different authorities have been compiled by Olivares (24). In view of the specific needs of this age group and recent modifications in the assumptions of adequate intakes in infancy, some aspects of current guidelines or directives for drinking water may be reconsidered. The evaluation of adequacy or optimization potential in terms of feeding infants and young children have preferably to be based at household level to enable the assessment of leaching from water pipes or contamination by other sources.

Public health recommendations should take into account that both desalinated water (25) and infant formula (26) are manufactured products and offer the potential of influencing the final nutrient mineral supply to the infant. Remineralization of processed waters should aim to achieve concentrations that are acceptable when combined with infant formulae but which are potentially useful in early childhood thereafter. However, the assessment may differ for each mineral under consideration.

1. Manganese and Molybdenum.

Manganese and molybdenum concentrations set out in the WHO drinking water guidelines (10, 27) are much higher than those observed in human milk and in the recommendations regarding their respective daily adequate intake (28), Table 2.

Manganese is an essential nutrient in the formation of bone, amino acids, cholesterol and carbohydrate metabolism and present in the metalloenzymes arginase, glutamine synthetase, and manganese superoxide dismutase. The recommended adequate intake (AI) is based on the intake from human milk and set at 0.003mg/day in the first six months (preceding recommendation: 0.3-0.6 mg (21)). Thereafter, an increased intake of 0.6mg/day (7-12 months) and 1.2mg/day for children aged 1-3 years (28) has been recommended. During the last decade other recommendations and directives have also decreased considerably: In Germany, infant formulae for special medical purposes had to provide 0.5 to 2mg manganese per day, equivalent to 0.66 to 2.67mg/L (29), but actual requirements are set at 0.325 to 1.3 mg/L (30). Analytical results of formulae ranged from 0.0 to 7.8mg Mn/L (3), with high concentrations observed in soy-protein-based formulae and formulae for special medical purposes. Subsequently considerably lower results were assessed (0.44 and 0.53mg Mn/L, (31)). Recommendations for infant milk formulae yielded a range of 1-100µg/100kcal (2), equivalent to 0.007 to 0.65mg Mn/L. Therefore, regardless of any change in drinking water composition or respective regulations, the potential significance of the contribution of drinking water supply to infant nutrition has increased considerably.

Table 2. Trace minerals in human milk and in drinking water guidelines

Mineral	Mature human milk (Lawrence and Lawrence 1999, (54))	Drinking water guidelines (WHO 1996 (27))	EC, SCF 2003 (2) (Recommended energy content: 60-70kcal/dl; based on 65kcal/dl, cow's-milk- protein- based formula)
	mg/L	mg/L	mg/L
Calcium (mg/L)	280	n.g.	325-910 (Ca:P=1-2)
Iron (mg/L)	0.40	0.3 ^b	1.95-8.45
Zinc (mg/L)	1.2	3.0 ^b	3.25-9.75
Copper (mg/L)	0.25	1.0 ^b ; 2.0 ^a (P)	0.228-0.65
Selenium (µg/L)	20	10 ^a	20-59
Fluoride (mg/L)	0.016	1.5 (P)	≤ 0.65
Magnesium (mg/L)	30	n.g.	33-98
Sodium (mg/L)	180	200 ^a	130-390
Sulphate (mg/L)	140 (sulphur)	250 ^a	n.g.
Chloride (mg/L)	420 (chlorine)	250 ^a	325-1040
Manganese (µg/L)	6	100 ^b ; 500 ^a (P)	6.5-650
Molybdenum (µg/L)	2µg/d ^c	70 ^a	n.g.

^a Health-based guideline value, (P): provisional; ^b Parameters in drinking water that may give rise to complaints from consumers, ^c FNB (28)

A WHO guideline value has been set at 0.5mg/L based upon aesthetics for manganese in drinking water (27). Powdered infant formula with the mean recommended manganese concentration (0.325mg/L) would exceed the maximum suggested by the SFC (2) if prepared with drinking water in the upper range of WHO guideline values. Animal studies (32, 33) indicate that neonatal manganese metabolism may be of special concern. Balance studies in infants confirmed substantial quantitative manganese retention in infants fed special diets as a result of metabolic disease or in premature infants fed routine mineral supplements and containing manganese by contamination (34, 35). Recommendations concerning drinking water for the reconstitution of infant formulae should take into consideration the low concentration in human milk and concerns about the safety of high manganese intakes. In situations where high manganese concentrations in drinking water cannot be avoided, the potential combination with formulae of high manganese content with high manganese water should be discouraged.

Molybdenum is a co-factor in the enzymes sulphite oxidase, xanthine oxidase and aldehyde oxidase (36). The adequate intake in infants and young children has been set for the age of 0-6 months at 2µg/day, derived from the intake by human milk (preceding recommendation: 15-30µg/day; (21)). In contrast to manganese, however, the AI for the age 7-12 months has been derived from the likely intake with human milk as a sole source (3µg/day) and not from the estimated intake with a mixed diet (28). This value will be greatly exceeded by formula-fed infants and by infants nourished with the recommended complementary feeds (37). The

uncertainty over molybdenum requirements is also reflected in the tolerable upper intake (UL) level defined for adults with 0.01mg/kg body weight (38) or, alternatively, 0.03mg/kg body weight (28). Both were based on the same trial (39), but differed in the assessment of the uncertainty factors. Despite concern about the infant's capacity to deal with excess amounts of Mo, no UL has been set for infancy.

The upper level of Mo intake has been set at 10µg/100kcal for nutrients for special dietary purposes in infancy (30). An infant formula prepared in accordance with this directive may result in a daily Mo uptake > 10µg/kg. In addition, an upper guideline value of 70µg/L has been set for drinking water (10). The maximum Mo intake by water used for powdered formula preparation exceeds 10µg/kg at an intake of 150ml/kg or more. This volume, however, is quite common in premature and young full-term infants (Figure 1) and unsupplemented infant formulae repeatedly contained > 100µg/L (40). The threshold for long-term safe molybdenum intake in infancy may need to be reconsidered and re-evaluated by studies in areas with naturally longterm high molybdenum intakes by drinking water before present WHO guidelines are extended on other processed waters.

2. Copper

Both copper deficiency and excess are of clinical importance: copper deficiency has been described in premature infants and is characterized by oedema, anaemia, leucopenia, neutropenia and osteoporosis (41). On the other hand, concerns have been raised after the description of Non-Indian Childhood Cirrhosis associated with excessive copper concentrations in well waters with a low pH and copper pipes (42). Examinations in a subgroup of infants from households with copper concentrations exceeding 0.8mg/L yielded no indication of a hazard resulting from copper pipes connected to public water supplies (8). The safety of the WHO guideline value (2mg/L) for copper in drinking water during infancy has been supported (43). Leaching from copper pipes and other potential problems of infant formula preparation have raised concern and have been addressed in respective recommendations (44).

Differences in the quality of water available from the tap or directly drawn at the water plant may be important in terms of its suitability in infancy and early childhood. The problems of the use of well water have to be considered in both developed and developing countries, though the proportion of public water supply and the use of home wells may vary considerably. In Germany, in 1998, only 2.5% of the drinking water supply was derived from well supplies. However, more than 58% of the [adult] study population consumed first-draw drinking water for their personal consumption (45). Concerns arise that common awareness of problems caused by the application of this practice to formula preparation cannot be assumed.

The FNB (28) recommendations in the first year for adequate copper intake, based on intake by human milk and adequate complementary feeds thereafter, were set for the first 6 months at 200µg/day and, in the 7-12 months thereafter, at 220µg/day. For young children of 1-3 years, the intake was set at 340µg/day. Regulations for both components of infant formulae - the powdered formula and the water - have declined considerably during the last few years: In the past, national directives for copper intake by foods for medical purposes in infancy were set at 0.5 to 2mg Cu per day, equivalent to 0.670 to 2.7mg/L formula (29). This has been reduced to 0.130 to 0.780mg Cu/L formula (30), slightly lower values are recommended in infant formulae (2). At the same time, the WHO Guideline value of 2mg/L (27) has been transferred into present regulations on drinking water, which formerly set a cut-off value of 3mg/L (46, 47). These changes result in a reduction of the upper margin of daily copper intake from > 4 mg/d to 2 mg/d at an intake of 750 ml/d.

3. Iron and zinc

Iron and zinc drinking guideline values are only set at levels that may give rise to complaints from consumers (27). Recommendations for both elements in infant food other than human milk have to take into consideration the importance of bioavailability (14). Other than for manganese, molybdenum and copper, these are within the concentrations observed in mature (iron) human milk or colostrum and transitional human milk (zinc), Table 2.

The low iron intake recommended in the first six months of life by the FNB (28), 0.27mg Fe/d, is based on the highly bioavailable intake with human milk. Thereafter, however, increased intakes are needed to cover the demands of growth, especially the increase in blood volume (7-12 months: 11mg/d; 1-3 years: 7mg/d). The American Academy of Pediatrics (48) advocated an iron supplementation of 4-12mg/L of infant formulae. The SCF (2) of the European Commission regarded 2-8mg/L as suitable for cow's-milk-based infant formula and 6-15mg/L for soy-based follow-on formula. 2mg/L in drinking water were not considered to present a hazard to health, but it has been assumed that acceptability would usually be affected above 0.3mg Fe/L because of the staining of laundry and plumbing fixtures (27). Therefore, in infants fed iron supplemented formulae, the contribution of drinking water to the iron intake may be of minor importance.

This assessment may be different in infants fed home-prepared, non-supplemented formulae or complementary feeds. In Bangladesh, children drinking water containing > 1mg Fe/L were significantly taller than those drinking < 1mg Fe/L (49). The higher iron concentrations were caused by soil characteristics and high concentrations occurred in clusters of tube wells. However, other minerals present in the soil along with iron may have promoted linear growth too. The effects of other trace elements (e.g. zinc, copper) have not been reported in a trial investigating the fortification of drinking water with iron for pre-school children attending day-care community institutions in Brazil. A reduction of anaemia in preschool children in a day care centre and in families with a low socio-economic status (50, 51) was assessed.

Zinc in colostrum may exceed 10mg/L, declining continuously down to a level of about 1.2mg/L at 4 months (52). Despite the high bioavailability of zinc in human milk, it may be difficult to satisfy needs adequately after a period of four to six months (15). In view of different bioavailability depending on the nutritional source, intake recommendations range from 1.1 to 6.6mg/day during the first six months of life, and 2.5 to 6.4mg/day thereafter (14). The FNB (28) recommended 2 and 3mg respectively, based on the zinc concentration in human milk as the basic source. Recommendations for infant and follow-on formulae (2) in accordance with the current infant formulae EC directive, yield 3.3–10mg/L or, if based on soy protein, 4.9–15.6mg/L. The upper level of tolerable zinc intake for children aged 1-3 years (53) has been set at 7mg/day. Water yielding a zinc concentration within the upper range of WHO guideline values (3mg/L, Table 2) might contribute considerably to the zinc intake in infancy.

4. Calcium and Magnesium

The calcium and magnesium concentrations in cow's milk far exceed those present in human milk (54) and have to be modified e.g. during the manufacturing of infant formulae based on cow's milk protein. The adequate intake (AI) for calcium has been set at 210mg (formula-fed infants: 315mg/d), 270mg and 500mg, respectively, for ages 0-6 and 7-12 months and 1-3 years. The AI for magnesium was set at 30mg (formula-fed infants: 35mg) and 75mg (0-6 and 7-12 months) and for ages 1-3 years the RDA was set at 80mg (11). The ranges recommended for infant formulae by the SCF (2) were 325-910mg/L Ca and 33-88mg/L Mg, respectively. Lesser absorption from infant milk formula compared to human milk has been considered, therefore the minimum requirements are set above that found in human milk.

High mineral concentrations in water increase the renal solute load and, therefore, are of concern in infancy. For natural mineral water labelled as “suitable for the feeding of infants”, the Committee on Nutrition of the German Society of Paediatrics (55) advocated, amongst others, concentrations of < 20mg Na/L and < 200mg/L sulphate. The Committee assessed that, at that time, the cut-off value for sulphate (200mg/L) in mineral waters with a sodium concentration < 20mg/L led to a natural limitation of the calcium and magnesium concentration. In view of this finding, no cut-off value for calcium and magnesium was set. In Switzerland, however, a cut-off value has been set for calcium < 200mg/L and magnesium < 40mg/L (56). Public drinking water in Germany yielded a mean concentration of 66.4 (0-555.9) mg/L calcium and 11.8 (0-555.9) mg/L magnesium (57). Transparency should be provided for the consumer with respect to Ca and Mg concentrations in public drinking water, this may not be achieved by referring to “water hardness”. Older infants fed mainly plant-based complementary foods and young children on vegetarian nutrition may benefit from drinking water contributing to calcium and / or magnesium intake.

5. Sodium

Children, the elderly, and persons with compromised renal systems are more susceptible than healthy adults to harmful effects of high sodium intake. The FNB (28) set the adequate intake (AI) at 120mg, derived from the intake of breast-fed infants. In accordance with respective regulations in the USA, the regulation for infant milk formulae in Europe supports an intake of 20-60mg/100kcal (58), equivalent to 127-292.5mg Na/d. Mineral quantities have been calculated and incorporated into baby formula by the manufacturers in accordance with current directives.

Actual data on sodium metabolism in infants fed modern infant milk formulae, however, are scarce and the interpretation has to take into account specific information: longitudinal sodium balance studies in infancy compared ready-to-feed liquid infant milk formula-fed (269mg Na/L) infants with breast-fed (138mg Na/L) infants (4). The daily retentions observed were 11.5mg/kg and 9.2mg/kg, respectively. The actual composition of this brand marketed as powdered infant formula has a sodium content of 120mg Na/100g powder. In Germany, a mean sodium drinking water concentration of 12.4 (<1.0-210) mg/L has been observed (59), concentrations exceeding 150mg/L are the rare exception. The upper level of the Drinking Water Ordinance is 200mg/L (47), equivalent to respective WHO guidelines. Prepared according to current recommendations for the preparation of infant feeds/formula in the first months of life (12, 44), these factors may theoretically result in a considerable range of sodium intake:

- Up to 186mg/L (formula + bottled mineral water labelled as suitable for the preparation of infant formula, containing ≤ 20mg/L).
- Up to 366mg/L (formula + drinking water with the upper limit of 200mg/L).
- Up to 310 or 490mg/L, respectively, through the components drinking water (as used in 1) or 2) and cow’s milk (580mg/L,(54)) if parents insist on the use of home-prepared formula and prepare it according to Kersting (12).

These examples also highlight potential hazards; drinking water with high sodium concentrations may elevate the sodium concentration of infant formulae considerably. Cow’s-milk-based infant formulae with low sodium content may be either not available, not affordable or inadequately prepared owing to regional or social constraints. The home-prepared replacement formula recipe described by WHO/UNICEF/UNAIDS (13) comprises a 2:1 dilution of cow’s milk with drinking water and added sugar. Cow’s milk contains more sodium than human milk; the sodium concentrations to be expected will range from slightly less than 406mg/L up to 586mg/L

or above (based on the recommendation of $\leq 20\text{mg/L}$ for bottled drinking water for infant use to $\leq 200\text{mg/L}$ for drinking water). Other minerals may potentially increase the renal solute load, too, necessitating higher quantities of water available to form urine in view of the reduced ability of infants to concentrate urine. This may reduce the margin of safety, especially under conditions of stress, such as infections accompanied by fever.

Where definite problems exist with drinking water (e.g. excessive nitrate content), packaged mineral water should be considered as an alternative to drinking water from the public supply. The infant formula industry does try to lower the higher mineral concentration of cow's milk compared to human milk, but with the addition of mineral waters this content may be increased again. Considerable ranges for sodium in packaged mineral waters have been observed in Europe (26.6 (1.3-1723.0) Na, (60)). This necessitates transparency as to whether a product is suitable for the use in infant nutrition or not. Paediatric recommendations for the suitable composition of packaged waters for use in infant feeding were set at $<20\text{mg Na/L}$ (44, 55). A certain contradiction arises in view of the previously mentioned cut-off value for public drinking water (200mg/L). However, these differences may scarcely be of practical relevance for public drinking water in Germany. In other regions, where sodium concentrations exceeding 180mg/L have been repeatedly described in drinking water (9,61), the assessment may differ considerably and preferable choices should be considered if processed water yields options.

Concern has been raised due to higher blood pressure in infants fed infant formulas with a higher sodium content due to the concentration in the drinking water (9). In addition, an association between formula feeding and blood pressure later in life has been found and attributed in part to the difference in sodium content between human milk and infant formula (62). There is a need for epidemiological studies that focus on the potential health effects of different concentrations of sodium and other [trace] mineral supply by drinking water on infants and young children. These should be conducted in regions where "natural experiments" render exposures close to the upper margins of current recommendations and necessitate these evaluations in view of potential consequences for the use of drinking water.

Especially in infancy, fluid consumption may serve nutritional purposes including caloric requirements but it also plays an important role in hydration. This is obvious in cases of diarrhoea, where increased fluid intake is necessary for rehydration and ORS (oral rehydration solution) recommended. A reduction in morbidity has been achieved by lowering the sodium concentration of the former standard solution from 90 to 75 mmol/L (2070mg/L to 1725 mg/L) (63). Oral rehydration solutions containing 60mmol/L (1380 mg/L) have been recommended in Europe in view of the concern over hypernatraemia or the osmotically-driven increase in stool output, especially in infants and young children (64).

Other hazards concerning sodium in infancy may arise from inadequate feeding practices. Hyponatremic seizures resulting from water intoxication have been associated with bottled drinking water used as an inexpensive "supplement" to cow's-milk-based infant formula (65). The labelling had been misinterpreted to indicate that the product had been produced specifically for infants and contained nutrients adequate for use as a feeding supplement. Inadequate dilution of infant foods may result in nutritionally insufficient feeding. Misjudgement, inexperience or poverty may lead to the use of bottled water products marketed specifically for infants as an affordable and appropriate feeding supplement. These aspects underline the importance of adequate labelling and instructions for the parent or carer.

V. CONCLUSIONS

Public health recommendations on infant nutrition should consider the complete process that includes all the stages involved in the source of the nutrients. This refers to the choice of the base ingredients used for the production of manufactured or home-prepared products well as to the provision of adequate instructions for the reconstitution and the feeding to the infant. Water companies should comply with existing guidelines and legislation, however the quality of water available from the tap may differ from that at the water plant and additional water treatment is possible at household level. Transparency in terms of a water's suitability for infant and child nutrition is essential for the consumer to make an informed choice. Depending on region and social status, consumers may have a choice of water (tap, mineral, bottled water) or have to accept what is available. Social inequalities and lack of information may result in non-recommended practices of infant feeding that can cause inadequate mineral intake.

Guidelines that are to be effective in terms of trace element intake in infants and young children have to consider age-specific trace element requirements. It is essential that they take into account the combined effects of general recommendations on infant feeding on a comparable organizational level. For example:

- *The WHO global strategy on infant and young child feeding* and specific recommendations for trace mineral and mineral intakes in infancy;
- The Standards of the Codex Alimentarius for infant formulae;
- The WHO/UNAIDS/UNICEF recommended (and other) recipes for home-prepared replacement milks for infants of HIV-infected mothers, and
- The WHO Guidelines for Drinking Water Quality.

The awareness of potential variations and analysis of the water used by a population is important for epidemiological assessments of the trace element supply in infancy. Intervention studies in communities in the intention of changing water sources or treatment could facilitate the evaluations of the implications of different mineral intakes in infancy and early childhood. The natural occurrence of concentrations in drinking water within the upper range of WHO drinking water guidelines may support the assessment of the safety for some minerals (e.g. molybdenum). In view of the consequences for public health recommendations and potential consequences for infant morbidity on the one hand, and potential economic consequences for governments and households on the other, there is a need for further scientific evidence of the optimal mineral concentrations in drinking water in early childhood.

References

1. WHO, World Health Organization, and UNICEF. Global Strategy for Infant and Young Child Feeding. Geneva: World Health Organization, 2003.
2. SCF, Scientific Committee on Food. Report of the Scientific Committee on Food on the Revision of Essential Requirements of Infant Formulae and Follow-on Formulae. SCF/CS/NUT/IF/65, 2003.
3. Lönnerdal B, Keen CL, Ohtake M, Tamura T. Iron, zinc, copper, and manganese in infant formulas. *Am J Dis Child* 1983;137:433-437.
4. Schulz-Lell G, Dörner K, Oldigs H-D, et al. Sodium and potassium metabolism in infancy. *Monatsschr Kinderheilkd* 1992;40:117-121.
5. Dijkhuizen MA, Wieringa FT, West CE, et al. Effects of iron and zinc supplementation in Indonesian infants on micronutrient status and growth. *J Nutr* 2001;131:2860-2865.
6. Lind T, Lönnerdal B, Stenlund H, et al. A community-based randomized controlled trial of iron and zinc supplementation in Indonesian infants: interactions between iron and zinc. *Am J Clin Nutr* 2003;77:883-890.
7. Penny ME, Marin R, Duran A, et al. Randomized controlled trial of the effect of daily supplementation with zinc or multiple micronutrients on the morbidity, growth, and micronutrient status of young Peruvian children. *Am J Clin Nutr* 2004;79:457-465.
8. Zietz BP, Dieter HH, Lakomed M., et al. Epidemiological investigation on chronic copper toxicity to children exposed via the public drinking water supply. *Sci Total Environ* 2003; 302:127-144.
9. Pomeranz A, Dolfen T, Korzets Z, et al. Increased sodium concentrations in drinking water increase blood pressure in neonates. *J Hypertension* 2002;20:203-207.
10. WHO. Guidelines for drinking-water quality. Recommendations, 2nd edition. Geneva: WHO, 1993.
11. FNB, Food Nutrition Board, Institute of Medicine. Dietary reference intakes for water, potassium, sodium, chloride, and sulphate. Washington, D.C.: The National Academies Press, 2003.
12. Kersting M. Ernährung des gesunden Säuglings: Lebensmittel- und mahlzeitenbezogene Empfehlungen. *Monatsschr Kinderheilkd* 2001;149:4-10.
13. WHO, UNAIDS, UNICEF. HIV and infant feeding counselling: A training course. Geneva: World Health Organization, 2000.
14. FAO, Food and Agriculture Organization and WHO, World Health Organization. Human vitamin and mineral requirements. Report of a Joint FAO/WHO Expert Consultation, Bangkok, Thailand. Rome: FAO/WHO, 2001.
15. Butte NF, Lopez-Alarcon MG, Garza C. Nutrient adequacy of exclusive breastfeeding for the term infant during the first six months of life. Geneva: World Health Organization, 2002.
16. Hofvander Y, Hagman U, Hillervik C, Sjölin S. The amount of milk consumed by 1-3 months old breast- or bottle-fed infants. *Acta Paediatr Scand* 1982;71:953-958.
17. Sievers E, Schleyerbach U, Garbe-Schönberg D, et al. Zinc intakes and plasma concentrations in infancy. *Adv Exp Med Biol* 2000;478:383-4.

18. Sievers E, Schleyerbach U, Arpe T et al. Molybdenum supply of very low-birth-weight premature infants during the first months of life. *Biol Trace Elem Res* 2001;80:97-106.
19. Hilbig A, Kersting M, Sichert-Hellert W. Measured consumption of tap water in German infants and young children as background for potential health risk assessments: data of the DONALD Study. *Food Addit Contam* 2002;19:829-836.
20. Manz F, Wentz A, Sichert-Hellert W. The most essential nutrient: Defining the adequate intake of water. *J Pediatr* 2002;141:587-592.
21. FNB, Food and Nutrition Board, Commission on Life Sciences, National Research Council. Recommended Dietary Allowances. 10th Edition. Washington D.C.: National Academy Press, 1989.
22. Sichert-Hellert W, Kersting M, Manz F. Fifteen year trends in water intake in German children and adolescents: Results of the DONALD Study. *Acta Paediatr* 2001; 90:732-737.
23. Petter LPM, Hourihane JO, Rolles CJ. Is water out of vogue? A survey of the drinking habits of 2-7 year olds. *Arch Dis Child* 1995;72:137-140.
24. Olivares M. Comprehensive Overview Paper: Essential Nutrients in Drinking Water . In: Present publication, in press.
25. Cotruvo J. Desalination Guidelines Development for Drinking Water: Background. In this publication.
26. Tomita M. New technologies and their application in the milk industry. *Asian-Aus. J. Anim. Sci., Animal Production for a Consuming World* 2000;13(suppl):376-382.
27. WHO and World Health Organization. Guidelines for drinking-water quality. Health criteria and other supporting information. Geneva: WHO, 1996.
28. FNB, Food and Nutrition Board, Institute of Medicine. Dietary Reference Intakes: Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium and Zinc. Washington DC: National Academy Press, 2001.
29. BMJFFS, Bundesminister für Jugend Familie Frauen und Gesundheit. Verordnung zur Änderung der Nährwert-Kennzeichnungsverordnung und der Diätverordnung. *Bundesgesetzblatt*, 1988;Teil 1:677-693.
30. EC, European Commission. Commission Directive 1999/21/EC on dietary foods for special medical purposes. *Official Journal of the European Communities* 1999;L 91:29-36.
31. Krachler M, Rossipal E, Irgolic KJ. Trace elements in formulas based on cow and soy milk and in Austrian cow milk determined by inductively coupled plasma mass spectrometry. *Biol. Trace Elem. Res* 1998;65:53-73.
32. Keen CL, Bell JG, Lönnerdal B. The effect of age on manganese uptake and retention from milk and infant formulas in rats. *J Nutr* 1986;116:395-402.
33. Tran TT, Chowanadisai W, Lönnerdal B, et al. Effects of neonatal dietary manganese exposure on brain dopamine levels and neurocognitive functions. *Neurotoxicology* 2002;145:1-7.
34. Sievers E, Oldigs H-D, Dörner K, et al. Trace element excess in PKU diets ? *J. Inher Metab Dis* 1990;13:897-905.
35. Sievers E, Kollmann M, Oldigs H-D, et al. Manganese (Mn) balances in preterm infants. In: Momcilovic B, ed. Trace elements in man and animals 7; 7th International Congress on

- Trace Elements in Man and Animals (TEMA 7) Dubrovnik (Yu) 1990, Nova Gradiska, GTP Gembarovski, 1990:11-14-11-16.
36. Turnlund JR. Molybdenum metabolism and requirements in humans. In: Sigel A et al. eds. Metal ions in biological systems. Molybdenum and tungsten: Their roles in biological processes, New York, Basel: Marcel Dekker, Inc. 2002;39:727-739.
 37. Anke M, Lösch E, Gleis M et al. Der Molybdängehalt der Lebensmittel und Getränke Deutschlands. In: Anke M ed. Mengen- und Spurenelemente, 13. Arbeitstagung. MTV Hammerschmidt GmbH, Gersdorf, 1993:537-553.
 38. European Commission – Opinion of the Scientific Committee on Food on the Tolerable Upper Intake Level of Molybdenum. ACF/CS/NUT/UPPLE/22, 2000.
 39. Fungwe TV, Buddingh F, Demick DS, et al. The role of dietary molybdenum on oestrous activity, fertility, reproduction and molybdenum and copper enzyme activities of female rats. Nutr Res 1990;10:515-524.
 40. Sievers E. Nutrient requirements for preterm infant formulas - molybdenum. J Nutr 2003;133:236-237.
 41. Sutton AM, Harvie A, Cockburn F, et al. Copper deficiency in the preterm infant of very low birthweight. Arch Dis Child 1985;60:644-651.
 42. Dieter HH, Schimmelpfennig E, Meyer E, Tabert M. Early childhood cirrhosis (ECC) in Germany between 1982 and 1994 with special considerations of copper aetiology. Eur. J Med Res 1999;4:233-242.
 43. Olivares M, Pizarro FSH, Lonnerdal B, Uauy R. Copper in infant nutrition: safety of World Health Organization provisional guideline value for copper content of drinking water. J Pediatr Gastroenterol Nutr 1998;26:251-257.
 44. Committee on Nutrition, German Society of Pediatrics. Hinweise zur Zubereitung und Handhabung von Säuglingsnahrungen. Monatsschr Kinderheilkd 2004;152:318-320.
 45. Becker K, Kaus S, Helm D, et al. Umwelt-Survey 1998, Band IV. Trinkwasser, Elementgehalte in Stagnationsproben des häuslichen Trinkwassers der Bevölkerung in Deutschland. Berlin: Fa. Werbung und Vertrieb, 2001.
 46. BMJFG, Bundesminister für Jugend Familie Frauen und Gesundheit. Verordnung zur Änderung der Trinkwasser-Verordnung und der Mineral- und Tafelwasser-Verordnung. Bundesgesetzblatt 1990;I:2600-2629.
 47. BMG, Bundesministerin für Gesundheit, and BMEVL, Bundesministerin für Ernährung Verbraucherschutz und Landwirtschaft. Verordnung zur Novellierung der Trinkwasserverordnung vom 21. Mai 2001. Bundesgesetzblatt 2001;24:959-981.
 48. AAP, American Academy of Pediatrics, Committee on nutrition. Iron fortification of infant formulas. Pediatrics 1999;104:119-23.
 49. Briend A, Hoque BA, Aziz KMA. Iron in tubewell water and linear growth in rural Bangladesh. Arch Dis Child 1990;65, 224-237.
 50. Dutra-de-Oliveira JE, Ferreira JB, Vasconcellos VP, Marchini JS. Drinking water as an iron carrier to control anemia in preschool children in a day-care center. J Am Coll Nutr 1994;13:198-202.

51. Dutra-de-Oliviera JE and Nogueira de Almeida CA. Domestic drinking water - an effective way to prevent anaemia among low socioeconomic families in Brazil. *Food Nutr Bull* 2002;23:213-216.
52. Dorea JG. Zinc in human milk. *Nutr Res* 2000;20:1645-1687.
53. SCF, Scientific Committee on Food . Opinion of the Scientific Committee on Food on the Tolerable Upper Intake Level of Zinc. SCF/CS/NUT/UPPLEV/62, 2003.
54. Lawrence RA, Lawrence RM: Breastfeeding. A guide for the medical profession. St.Louis, Missouri: Mosby Inc., 1999.
55. Committee on Nutrition, German Society of Pediatrics. Zur Zubereitung von Säuglingsnahrung mit Mineralwasser. *Sozialpädiatrie* 1991;13:722-728.
56. Committee on Nutrition, Swiss Society of Pediatrics. Mineralwasser zur Herstellung von Säuglingsnahrung. *SÄZ* 1990;71:487-489.
57. Schimatschek HF. Vorkommen, Bedeutung und Analytik von Calcium und Magnesium. In: Grohmann A, Hässelbarth U, Schwerdtfeger W, ed. *Die Trinkwasserverordnung*, 4th edn., Berlin, Erich Schmidt Verlag, 2003:511-515.
58. EEC, European Commission. Commission Directive 91/321/EEC on infant formulae and follow-on formulae. *Official Journal of the European Communities*, 1991:35-49.
59. Wolter R . Erhebung BIBIDAT 1989, cited in: Großklaus R, Vorkommen, Bedeutung und Bestimmung von Natrium und Chlorid. In: Grohmann A, Hässelbarth U, Schwerdtfeger W, eds. *Die Trinkwasserverordnung* 4th edn. Berlin, Erich Schmidt Verlag 2003:547-555.
60. Misund A., Frengstad B, Siewers U, Reimann C (1999) Variation of 66 elements in European bottled mineral waters. *Sci Total Environ* 1999;243/244 :21-41.
61. Alam IA, Sadiq M. Metal contamination of drinking water from corrosion of distribution pipes. *Environ Pollut* 1989;57:167-178.
62. Martin RM, McCarthy A, Smith GD, et al. Infant nutrition and blood pressure in early adulthood: the Barry Caerphilly growth study. *Am J Clin Nutr* 2003;77:1489-1497.
63. WHO and UNICEF. Reduced osmolarity oral rehydration salts (ORS) formulation report from a meeting of experts. New York: WHO/FCH/CAH/01, 2001.
64. ESPGAN, European Society of Pediatric Gastroenterology and Nutrition. Recommendations for composition of oral rehydration solutions for the children of Europe. *J Ped Gastroenterol Nutr* 1992;14:113-115.
65. CDC, Centres for Disease Control and Prevention: Hyponatraemic seizures among infants fed with commercial bottled drinking water - Wisconsin, 1993. *MMWR* 1994;43:641-643.