



World Health Organization


Mortality Country Fact Sheet 2006

Lithuania



Summary	Year	Males	Females	Both sexes
Population (millions)	2005	2	2	3
Life expectancy (years)	2004	66	78	72
Under-5 mortality (per 1 000 live births)	2004	10	9	10
Adult mortality (per 1 000)	2004	304	102	
Maternal mortality (per 100 000 live births)	2000		19	

Source: World Health Statistics 2006




Legend:


- WHO European Region
- Lithuania

Source: World Health Statistics 2006

Under-5 mortality



Source: i) WHO mortality database ii) World Health Statistics 2006



Source: World Health Report 2005

Under-5 mortality: for highest and lowest quintiles
Lithuania, DHS

Wealth/assets quintiles	Lowest	Highest	Ratio

Sex	Males	Females	Ratio

Urban/Rural	Rural	Urban	Ratio

Mother's education quintiles	None	Higher	Ratio

Note: rate per 1 000 live births for 10-year period preceding the survey
Source: DHS



Note: rate for 10-year period preceding the survey
Source: DHS

Causes of death in children under-5

Distribution of causes of death among children under 5 years of age Lithuania, 2000-2003			Annual estimated proportions of death by cause for neonates Lithuania, 2000		
Causes	Deaths ^b		Causes	Deaths ^c	
	(%)	Regional average (%)		(%)	Regional average ^c (%)
Total neonatal deaths	100	100	Total neonatal deaths	100	100
Neonatal causes ^a	41	44	Neonatal tetanus	0	1
HIV/AIDS	0	0	Severe infection ^a	8	18
Diarrhoeal diseases	0	10	Birth asphyxia	16	18
Measles	0	0	Diarrhoeal diseases	0	1
Malaria	0	0	Congenital anomalies	40	19
Pneumonia	5	13	Preterm birth ^b	28	38
Injuries	17	6	Others	8	6
Others	36	25			

a. Includes diarrhoea during neonatal period
b. Sum of individual proportions may not add up to 100% due to rounding.

a. Includes deaths from pneumonia, meningitis, sepsis/septicaemia and other infections during the neonatal period.
b. Includes only deaths directly attributed to prematurity and to specific complications of preterm birth such as surfactant deficiency, but not all deaths in preterm infants.
c. Sum of individual proportions may not equal 100% due to rounding.

Causes of Death

Top ten causes of death, all ages Lithuania, 2002				Life expectancy at birth among males (years) Lithuania, 2004	
Causes	Deaths		Years of Life Lost	30 40 50 60 70 80 90 100	
	(000)	(%)		[Scale]	
All causes	41	100	100	[Scale]	
Ischaemic heart disease	14	36	22	[Scale]	
Cerebrovascular disease	5	12	8	[Scale]	
Self-inflicted injuries	1	4	8	[Scale]	
Trachea, bronchus, lung cancers	1	4	4	[Scale]	
Chronic obstructive pulmonary disease	0	2	2	[Scale]	
Colon and rectum cancers	0	2	2	[Scale]	
Stomach cancer	0	2	2	[Scale]	
Road traffic accidents	0	2	4	[Scale]	
Poisonings	0	2	3	[Scale]	
Cirrhosis of the liver	0	2	3	[Scale]	

Source: [Death and DALY estimates by cause, 2002](http://www.who.int/entity/healthinfo/statistics/bodgbdeathdalyestimates.xls)
<http://www.who.int/entity/healthinfo/statistics/bodgbdeathdalyestimates.xls>

Life expectancy at birth among females (years) Lithuania, 2004
30 40 50 60 70 80 90 100 [Scale]

Maternal mortality ratio (per 100 000 live births) Lithuania, 2000
0 500 1000 1500 2000 [Scale]

Legend:
 WHO European Region
 Lithuania

Source: World Health Statistics 2006