




World Health Organization


Mortality Country Fact Sheet 2006

Nauru



Summary	Year	Males	Females	Both sexes
Population (millions)	2005	0
Life expectancy (years)	2004	58	65	61
Under-5 mortality (per 1 000 live births)	2004	35	24	30
Adult mortality (per 1 000)	2004	448	303	
Maternal mortality (per 100 000 live births)	

Source: World Health Statistics 2006




Legend:


- WHO Western Pacific Region
- Nauru

Source: World Health Statistics 2006

Under-5 mortality



Source: i) WHO mortality database ii) World Health Statistics 2006



Source: World Health Report 2005

Under-5 mortality: for highest and lowest quintiles
Nauru, DHS

Wealth/assets quintiles	Lowest	Highest	Ratio

Sex	Males	Females	Ratio

Urban/Rural	Rural	Urban	Ratio

Mother's education quintiles	None	Higher	Ratio

Note: rate per 1 000 live births for 10-year period preceding the survey
Source: DHS



Note: rate for 10-year period preceding the survey
Source: DHS



World Health Organization

Mortality Country Fact Sheet 2006

Causes of death in children under-5

Distribution of causes of death among children under 5 years of age Nauru, 2000-2003			Annual estimated proportions of death by cause for neonates Nauru, 2000		
Causes	Deaths ^b		Causes	Deaths ^c	
	(%)	Regional average (%)		(%)	Regional average ^c (%)
Total neonatal deaths	100	100	Total neonatal deaths	100	100
Neonatal causes ^a	7	47	Neonatal tetanus	0	3
HIV/AIDS	0	0	Severe infection ^a	22	21
Diarrhoeal diseases	38	12	Birth asphyxia	12	26
Measles	5	1	Diarrhoeal diseases	0	1
Malaria	0	0	Congenital anomalies	17	8
Pneumonia	30	14	Preterm birth ^b	42	32
Injuries	19	7	Others	6	8
Others	0	18			

a. Includes diarrhoea during neonatal period
b. Sum of individual proportions may not add up to 100% due to rounding.

a. Includes deaths from pneumonia, meningitis, sepsis/septicaemia and other infections during the neonatal period.
b. Includes only deaths directly attributed to prematurity and to specific complications of preterm birth such as surfactant deficiency, but not all deaths in preterm infants.
c. Sum of individual proportions may not equal 100% due to rounding.

Causes of Death

Top ten causes of death, all ages Nauru, 2002				Life expectancy at birth among males (years) Nauru, 2004	
Causes	Deaths		Years of Life Lost (%)	30 40 50 60 70 80 90 100	
	(000)	(%)		[Scale]	
All causes	0	100	100	[Scale]	
Ischaemic heart disease	0	14	11	[Scale]	
Hypertensive heart disease	0	6	5	[Scale]	
Cerebrovascular disease	0	6	4	[Scale]	
Diabetes mellitus	0	5	3	[Scale]	
Nephritis and nephrosis	0	4	5	[Scale]	
Lower respiratory infections	0	4	3	[Scale]	
Cirrhosis of the liver	0	2	2	[Scale]	
Chronic obstructive pulmonary disease	0	2	1	[Scale]	
Hepatitis B	0	2	4	[Scale]	
Oesophagus cancer	0	2	1	[Scale]	

Source: [Death and DALY estimates by cause, 2002](http://www.who.int/entity/healthinfo/statistics/bodgbddeathdalyestimates.xls)
<http://www.who.int/entity/healthinfo/statistics/bodgbddeathdalyestimates.xls>

Life expectancy at birth among females (years) Nauru, 2004	
30 40 50 60 70 80 90 100	[Scale]

Maternal mortality ratio (per 100 000 live births) Nauru, 2000	
0 500 1000 1500 2000	[Scale]

Legend:
■ WHO Western Pacific Region
■ Nauru

Source: World Health Statistics 2006