

response. The data on the gastrointestinal effects of copper must be used with caution, as the effects observed are influenced by the concentration of ingested copper to a greater extent than the total mass or dose ingested in a 24-hour period. Recent studies have delineated the threshold for the effects of copper in drinking-water on the gastrointestinal tract, but there is still some uncertainty regarding the long-term effects of copper on sensitive populations, such as carriers of the gene for Wilson disease and other metabolic disorders of copper homeostasis.

Cyanazine

Cyanazine (CAS No. 21725-46-2) is a member of the triazine family of herbicides. It is used as a pre-emergence and post-emergence herbicide for the control of annual grasses and broadleaf weeds. It can be degraded in soil and water by microorganisms and by hydrolysis.

Guideline value	0.0006 mg/l (0.6 µg/l)
Occurrence	Has been detected in surface water and groundwater, usually at concentrations of a few micrograms per litre, although levels as high as 1.3 and 3.5 mg/l have been measured in surface water and groundwater, respectively
TDI	0.198 µg/kg body weight based on a NOAEL of 0.198 mg/kg body weight for hyperactivity in male rats in a 2-year toxicity/carcinogenicity study, using an uncertainty factor of 1000 (100 for interspecies and intraspecies variation and 10 for limited evidence of carcinogenicity)
Limit of detection	0.01 µg/l by GC-MS
Treatment performance	0.1 µg/l should be achievable using GAC
Guideline value derivation	
• allocation to water	10% of TDI
• weight	60 kg adult
• consumption	2 litres/day
Assessment date	1998
Principal reference	WHO (2003) <i>Cyanazine in drinking-water</i>

On the basis of the available mutagenicity data on cyanazine, evidence for genotoxicity is equivocal. Cyanazine causes mammary gland tumours in rats but not in mice. The mechanism of mammary gland tumour development in rats is currently under investigation and may prove to be hormonal. Cyanazine is also teratogenic in rats at dose levels of 25 mg/kg body weight per day and higher.

Cyanide

Cyanides can be found in some foods, particularly in some developing countries, and they are occasionally found in drinking-water, but usually only at very low concentrations. However, there are occasions on which large spills of cyanide, associated with industry, occur, and these can give rise to very high concentrations in drinking-water source waters, particularly surface waters.