

WHO European Childhood Obesity Surveillance Initiative (COSI)

A brief review of results from round 6 of COSI (2022–2024)



FACT SHEET HIGHLIGHTS 2022-2024



About COSI

Overweight and obesity among children remain one of the major public health challenges facing the WHO European Region. Overweight and obesity affect a child's immediate physical and mental health, educational attainment, and quality of life. Additionally, unhealthy body weight in early life can increase the risk of premature death and disability in adulthood, especially from noncommunicable diseases, the leading risk factor for premature mortality in the Region.

The WHO European Childhood Obesity Surveillance Initiative (COSI), established in 2007, has been measuring trends in overweight and obesity among primary school-aged children for over 15 years. Based on a common protocol, it collects nationally representative data from children aged 6-9 years through standardized measurements of body weight and height, along with additional data on dietary intake,

physical activity, sedentary behaviour, family background and school environments.

Participation in COSI has grown from 13 countries in 2007 to 48 countries in 2024, making it one of the largest childhood obesity surveillance initiatives of its kind globally (Fig.1)1. The first six rounds of data collection have gathered anthropometric data from over 1.7 million children.

The sixth round, conducted between 2022 and 2024, measured around 470 000 children in 37 countries in the Region. The full details of the sixth round of data collection are described in a report 2,3 .

Fig.1. Description of countries participating in the COSI, by round of data collection

(2012 - 2013)

ROUND 3

Belgiuma

Bulgaria

Cyprus

Czechia

Hungary

Ireland

5. Greece

Italy

Latvia

10. Lithuania

12. North Macedonia

11. Malta

13. Norway

14. Portugal

15. Slovenia

16. Spain

8.

9.

| ROUND 1 | | |
|-------------|----------------------|--|
| (2007-2008) | | |
| 1. | Belgium ^a | |
| 2. | Bulgaria | |
| 3. | Cyprus | |
| 4. | Czechia | |
| 5. | Ireland | |
| 6. | Italy | |
| 7. | Latvia | |
| 8. | Lithuania | |
| 9. | Malta | |
| 10. | Norway | |
| 11. | Portugal | |
| 12. | Slovenia | |
| 13. | Sweden | |
| | | |
| | | |
| | | |

(2009-2010)1. Belgium^a Bulgaria Cyprus Czechia 5. Ireland Italy 7. Latvia

4.

6.

ROUND 2

| 8. | Lithuania |
|-----|-----------|
| 9. | Malta |
| 10. | Norway |
| 11. | Portugal |
| 12. | Slovenia |
| 13. | Sweden |
| 14. | Greece |
| | |

16. North Macedonia 17. Spain

15. Hungary

Sweden 18. Albania Republic of Moldova

20. Romania 21. San Marino 22. Türkiye

ROUND 4

(2015-2017)1. Albania Belgium Bulgaria 4. Cyprus 5. Czechia 6. Greece Hungary 7. Ireland 8. Italy 9. 10. Latvia 11. Lithuania 12. Malta 13. North Macedonia 14. Norway 15. Portugal 16. Republic of Moldova

17. Romania 18. San Marino 19. Slovenia 20. Spain

21. Sweden 22. Türkive 23. Austria 24. Croatia

25. Denmark 26. Estonia 27. Finland 28. France

29. Georgia 30. Kazakhstan

31. Kyrgyzstan 32. Montenearo

33. Netherlands (Kingdom of the)

34. Poland

35. Russian Federation^b

36. Serbia 37. Slovakia

38. Taiikistan 39. Turkmenistan

c) Data collection was disrupted by the COVID-19 pandemic and hence could not take place.

d) Data were collected only in the Bremen federal state (GER-BR).

b) Data were collected only in Moscow (rounds 4 and 5, RUS-MS)

e) Data were collected only in Almaty (KAZ-AL)

Text in red: countries that participated

Text in bold: countries that collected

a) Data were collected only in Flanders.

and Yekaterinburg (round 5, RUS-YK).

data during the time period of the

listed data collection round.

in COSI for the first time.

f) Due to difficulties encountered during the data collection process, the number of children measured is insufficient to estimate the prevalence of overweight and obesity in accordance with the COSI Protocol.

ROUND 5

(2018-2020)Albania Austria Belgium Bulgaria 5. Croatia 6. Cyprus Czechia 7. 8. Denmark Estonia 9 10. Finland 11 France 12. Georgia 13. Greece 14. Hungary 15. Ireland 16. Italy 17. Kazakhstan Kyrgyzstan 19. Latvia 20. Lithuania 21. Malta 22. Montenearo 23. Netherlands (Kingdom of the) (NET) 24. North Macedonia 25. Norway 26. Poland 27. Portugal 28. Republic of Moldova

29. Romania 30. Russian Federationb 31. San Marino 32. Serbia

33. Slovakia 34. Slovenia 35. Spain 36. Sweden 37. Taiikistan

38. Türkiye 39. Turkmenistan

40. Armenia 41. Azerbaijan^c

42. Bosnia and Herzegovina 43. Germany^d

44. Israel 45. Uzbekistan^a **ROUND 6**

(2022 - 2024)

1. Albania (ALB) Armenia (ARM) Austria (AUT)

Azerbaijan (AZE)

Belgium (BEL)

6. Bosnia and Hercegovina (BiH) Bulgaria (BUL) 7.

Croatia (CRO) 8. Cyprus (CYP) 9. 10. Czechia (CZH) 11. Denmark (DEN) 12. Estonia (EST) 13. Finland (FIN) 14. France (FRA) 15. Georgia (GEO)

16. Germany (GER-BR)d 17. Greece (GRE) 18. Hungary (HUN) 19. Ireland (IRE) 20. Israel (ISR) 21. Italy (ITA)

22. Kazakhstan (KAZ-AL)º 23. Kyrgyzstan (KGZ)

24. Latvia (LVA) 25. Lithuania (LTU) 26. Malta (MAT) 27. Montenegro (MNE)

28. Netherlands (Kingdom of the) (NET) 29. North Macedonia (MKD)

30. Norway (NOR) 31. Poland (POL)

32. Portugal (POR) 33. Republic of Moldova (MDA)

34. Romania (ROM)

35. Russian Federation (RUS) 36. San Marino (SMR) 37. Serbia (SRB)

38. Slovakia (SVK) 39. Slovenia (SVN) 40. Spain (SPA)

41. Sweden (SWE) 42. Tajikistan (TJK)

43. Türkiye (TUR) 44. Turkmenistan (TKM) 45. Uzbekistan (UZB)

46. Luxemburg (LUX) 47. Switzerland (SWI)

48. Ukraine (UKR)

^{1.} Since 2007, overweight and obesity prevalence has been estimated for 46 of the 48 COSI countries. Data collection could not proceed in Bosnia and Herzegovina or in Switzerland.

^{2.} Report on the sixth round of data collection, 2022–2024: WHO European Childhood Obesity Surveillance Initiative (COSI). The full report is under development and will be published in 2025.

Minor deviations from national results published elsewhere may occur due to slight differences in methods of analysis

OVERWEIGHT and OBESITY

among children aged 7-9 years

in the WHO European Region – latest available data from COSI

Since the first round of COSI data collection in 2007, 46 countries have collected data at least once. When examining the most recent data from all countries, including previous rounds for those that did not participate in the sixth round, the highest prevalence of overweight and obesity in boys and girls were found in southern European countries, such as Cyprus, Greece and Italy (Fig.2).

Conversely, the lowest prevalence estimates of overweight and obesity among children (both genders combined) were observed in Central Asian countries, namely Kyrgyzstan, Tajikistan, Turkmenistan

and Uzbekistan. In the majority of countries (27 of 46), at least one in ten children was living with obesity, while only six countries reported that less than one in five children is affected by overweight or obesity.

The prevalence estimates shown in Fig.2 were calculated by including only one targeted age group per country as described in the footnote below. Differences in prevalence estimates among countries may be partially explained by variations in the targeted age group.

Fig.2. Overweight and obesity prevalence estimates in boys and girls aged 7-9 years according to the most recent available source of COSI.



OVERWEIGHT

among children aged 7-9 years

in the WHO European Region - COSI round 6 (2022-2024)

Overall in this population, 25% of children aged 7–9 years were living with overweight (including obesity) in 2022–2024 according to WHO definitions (Fig.3). Data confirmed important between-country differences, with country-specific prevalence of overweight (including obesity) ranging from 9% in Uzbekistan to 42% in Cyprus. In 11 of the 37 countries that participated in round 6, one third or more of children were living with overweight or obesity.


Results by sex showed a higher prevalence of overweight among boys (26%, overall estimate) compared to girls (23%) (Fig.4). The


country-specific prevalence ranged from 11% to 48% among boys and from 8% to 36% among girls. In most countries, the prevalence of overweight (including obesity) was higher among boys than girls, with a difference of five percentage points or more in 11 countries.


The prevalence estimates shown in Fig.3 and 4 were calculated by including only one targeted age group per country as described in the footnote below. Differences in prevalence estimates between countries may be partially explained by variations in the targeted age group.

Girls

Fig.3. Prevalence of overweight (including obesity) in children aged 7-9 years (%); COSI round 6 (2022-2024)







In Fig.3 and 4, data relate to: (i) 7-year-olds in Bulgaria, Czechia, Denmark, Estonia, Finland, Georgia, Greece, Hungary, Ireland, Israel, Kyrgyzstan, Latvia, Lithuania, Luxemburg, Malta, Montenegro, North Macedonia, Portugal, Republic of Moldova, Romania, Slovakia, Slovenia, Spain, Türkiye, Ukraine and Uzbekistan: (ii) 8-year-olds in Albania, Austria, Azerbaijan, Croatia, Germany (Bremen federal state only), Italy, Kazakhstan (Almaty only), Poland, San Marino and Sweden; and (iii) 9-year-olds in Cyprus. To improve readability, the prevalence estimates are presented as whole numbers, rounded to the necrest integer. Gender differences, measured in percentage points, were calculated by subtracting the estimates for girls from those for boys, without rounding to the necrest whole number. The asterisk indicates a significant difference (P < 0.05). The study population in round 6 of COSI differs from that in round 5, as different groups of countries participated in each round. Therefore, the overalle stimates for round 6 are not directly comparable with those from the previous round, which have been published elsewhere.

OVERWEIGHT

among children aged 6-9 years

in the WHO European Region – comparison between COSI round 6 (2022–2024) and COSI round 5 (2018–2020)

Twenty-eight of the 37 countries that participated in round 6 also collected data in round 5. Fig.5 and 6 show the difference in the prevalence of overweight in boys and girls, respectively, between the two rounds of data collection. The difference was calculated by subtracting the estimates for round 5 from the estimates for round 6.

A statistically significant increase in the prevalence of overweight among boys was observed in three countries (indicated with * in Fig.5), namely Malta, Slovenia and Sweden, with the difference between the two rounds of data collection ranging from +1.1 to +2.7 percentage points. Conversely, a statistically significant decrease emerged in two

countries (Israel and Italy – 1.8 and 2.6 percentage points less, respectively).

The increase in the prevalence of overweight was slightly more frequent among girls, with four countries recording a higher prevalence in round 6 compared to the previous round of data collection – Bulgaria, Malta, Slovenia and Sweden. The difference between the two rounds tended to be more pronounced and ranged from +2.2 to +4.1 percentage points. On the other hand, a statistically significant decline was observed in three countries, namely Greece, Israel and Spain (6.0, 1.5 and 5.2 percentage points less, respectively).

Fig.5. Changes in prevalence of overweight (including obesity) in boys aged 6–9 years between COSI round 5 (2018–2020) and round 6 (2022–2024)





Fig.6. Changes in prevalence of overweight (including obesity) in girls aged 6–9 years between COSI round 5 (2018–2020) and round 6 (2022–2024)



• OBESITY

among children aged 7-9 years

in the WHO European Region - COSI round 6 (2022-2024)

Overall in this population, one in ten children aged 7–9 years was living with obesity (Fig.7). Data show high heterogeneity among countries, with the prevalence of obesity ranging from 3% to 20%. The highest prevalence of obesity among children (both genders combined) was observed in Cyprus, Greece, Italy and Malta, while the prevalence was lowest in Denmark, Kazakhstan (Almaty), Kyrgyzstan and Uzbekistan. These results align with findings from previous rounds of data collection, which reported a north-south gradient in the prevalence of obesity, with the highest values for height in northern Europe, the highest values for weight in southern Europe, and the lowest values for both weight and height in Central Asia.

Overall, the prevalence of obesity among boys was 12%, compared to 8% among girls (Fig.8). At the country level, obesity prevalence ranged from 4% to 25% among boys and from 2% to 15% among girls. Similar to the findings for overweight, boys were living more frequently with obesity compared to girls in most of the countries that collected data in round 6.

The prevalence estimates shown in Fig.7 and 8 were calculated by including only one targeted age group per country described in the footnote below. Differences in prevalence estimates between countries may be partially explained by variations in the targeted age group.

Girls

Fig.7. Prevalence of obesity in children aged 7-9 years (%); COSI round 6 (2022-2024)





Fig.8. Prevalence of obesity in boys and girls aged 7–9 years (%), and difference between genders (boys minus girls, percentage points on horizontal axis); COSI round 6 (2022–2024)



In Fig.7 and 8, data relate to: (i) 7-year-olds in Bulgaria, Czechia, Denmark, Estonia, Finland, Georgia, Greece, Hungary, Ireland, Israel, Kyrgyzstan, Latvia, Lithuania, Luxemburg, Malta, Montenegro, North Macedonia, Portugal, Republic of Moldova, Romania, Slovakia, Slovenia, Spain, Türkiye, Ukraine and Uzbekistan; (ii) 8-year-olds in Albania, Austria, Azerbaijan, Croatia, Germany (Bremen federal state only), Italy, Kazakhstan (Almaty only), Poland, San Marino and Sweden; and (iii) 9-year-olds in Cyprus. To improve readability, the prevalence estimates are presented as whole numbers, rounded to the necrest integer. Gender differences, measured in percentage points, were calculated by subtracting the estimates for girls from those for boys, without rounding to the necrest whole number. The asterisk indicates a significant difference (P < 0.05). The study population in round 6 of COSI differs from that in round 5, as different groups of countries participated in each round. Therefore the overall estimates for round 6 are round for green profiles where

OBESITY

among children aged 6-9 years

in the WHO European Region – comparison between COSI round 6 (2022–2024) and COSI round 5 (2018–2020)

Fig.9 and 10 illustrate the differences in the prevalence of obesity among boys and girls, respectively, between the sixth round of COSI data collection and the fifth round for the 28 countries that participated in both rounds. The difference was calculated by subtracting the estimates for round 5 from the estimates for round 6.

Three countries showed a statistically significant increase in the prevalence of obesity among boys – Malta, Slovenia and Sweden, with differences between the two rounds ranging from +1.4 to +1.9 percentage

points. Only San Marino recorded a statistically significant decrease (4.5 percentage points less).

As with overweight, the rise in the prevalence of obesity was slightly more pronounced among girls compared to boys. In four countries, namely Finland, Malta, Slovenia and Sweden, the round 6 prevalence estimates were significantly higher than those of the previous round of data collection, with differences ranging from 1.4 to 2.8 percentage points, while no country showed a statistically significant decline for girls.

Fig.9. Changes in prevalence of obesity in boys aged 6–9 years between COSI round 5 (2018–2020) and round 6 (2022–2024)







Fig.10. Changes in prevalence of obesity in girls aged 6-9 years between COSI round 5 (2018-2020) and round 6 (2022-2024)





Conclusions and next steps

Data collected in the sixth round of COSI, conducted between 2022 and 2024 in 37 countries and encompassing around 470 000 children, confirm that childhood overweight and obesity remain a major public health challenge in the Region. Overall, in the participating countries, 25% of children aged 7–9 years were living with overweight (including obesity), and 10% were affected by obesity. Important differences continue between countries, with the overall prevalence of overweight ranging from 9% to 42% and obesity from 3% to 20%.

COSI round 6 figures are the first estimates following the COVID-19 pandemic, providing initial insights into how the burden of overweight and obesity among children has changed in the post-pandemic period in the Region. Comparisons with previous round estimates (2018–2020) show different patterns of change in the 28 countries with available data. Despite stability in both overweight and obesity prevalence in several countries, where changes occurred, they were more likely to be increases rather than decreases, suggesting that the COVID-19 pandemic may have further exacerbated the problem in some countries. Future analyses, including the findings from all COSI rounds of data collection, will provide a more comprehensive picture.

Ending childhood obesity requires a comprehensive strategy that includes creating environments which promote healthy diets and ample physical activity for both young children and school-aged children. It also involves preconception and pregnancy care, breastfeeding support, as well as the provision of access to weight management programmes as part of universal health coverage.

Data from COSI play a crucial role in monitoring progress and informing policy-makers as countries look toward the Fourth High-level Meeting of the United Nations General Assembly on the prevention and control of NCDs (noncommunicable diseases) in 2025, aiming to halt the rise in obesity and achieve Sustainable Development Goal Target 3.4, which seeks to reduce premature mortality from NCDs by one third. These data can help promote the policy actions required today to build a future generation that will be resilient to NCDs.

World Health Organization Regional Office for Europe UN City, Marmorvej 51, DK-2100, Copenhagen Ø, Denmark Tel.: +45 45 33 70 00

Fax: +45 45 33 70 01 Email: eurocontact@who.int Website: www.who.int/europe

The WHO Regional Office for Europe

The World Health Organization (WHO) is a specialized agency of the United Nations created in 1948 with the primary responsibility for international health matters and public health. The WHO Regional Office for Europe is one of six regional offices throughout the world, each with its own programme geared to the particular health conditions of the countries it serves.

